


**FIGURE 1. One- and 3-year probabilities of continued opioid use among opioid-naïve patients, by number of days' supply\* of the first opioid prescription — United States, 2006–2015**




\* Days' supply of the first prescription is expressed in days (1–40) in 1-day increments. If a patient had multiple prescriptions on the first day, the prescription with the longest days' supply was considered the first prescription.

data on tramadol for pain management are sparse, with only one trial exceeding 12 weeks in duration (6). Despite this, among patients initiated with tramadol, >64% of patients who continued opioid use beyond 1 year were still on tramadol, suggesting that tramadol might be prescribed intentionally for chronic pain management. A 2016 study in Oregon (7), which did not include tramadol (a predictor of long-term use according to current data), reported similar findings: opioid naïve patients aged <45 years who received two prescription fills (versus one) or a cumulative dose of 400–799 (versus <120) morphine milligram equivalents in their first month of therapy were 2.3 and 3.0 times as likely to be chronic opioid users, respectively. However, that analysis only examined opioid use in the first month after initiation of opioid therapy to characterize risks for long-term use and did not account for the actual duration of therapy.

The findings in this report are subject to at least five limitations. First, although the cumulative dose of the first episode of opioid use is described, the likelihood of long-term use when the prescriber was titrating the dose was not determined. Rather, the total cumulative dose was calculated, which might have been increasing or decreasing over time. Second, the extent to which chronic opioid use was intentional versus the outgrowth of acute use is not known. Less than 1% of patients in this analysis were prescribed Schedule II long-acting opioids at the outset, so intentional chronic opioid prescribing might be uncommon; however, approximately 10% of patients were prescribed tramadol, which might indicate intentional chronic

**FIGURE 2. One- and 3-year probabilities of continued opioid use among opioid-naïve patients, by number of prescriptions\* in the first episode of opioid use — United States, 2006–2015**



\* Number of prescriptions is expressed as 1–15, in increments of one prescription.

opioid prescribing. Third, information on pain intensity or duration were not available, and the etiology of pain, which might influence the duration of opioid use, was not considered in the analysis. Fourth, the frequency of prescriptions having certain days' supply (e.g., prescriptions with a 7-day supply would be more frequently observed than those with an 11- or 13-day supply) was not considered. The variability in the relationships between days' supply, the cumulative dose, and duration of first episode and the probability of long-term use could be affected. Finally, prescriptions that were either paid for out-of-pocket or obtained illicitly were not included in the analysis.

Transitions from acute to long-term therapy can begin to occur quickly: the chances of chronic use begin to increase after the third day supplied and rise rapidly thereafter. Consistent with CDC guidelines, treatment of acute pain with opioids should be for the shortest durations possible. Prescribing <7 days (ideally ≤3 days) of medication when initiating opioids could mitigate the chances of unintentional chronic use. When initiating opioids, caution should be exercised when prescribing >1 week of opioids or when authorizing a refill or a second opioid prescription because these actions approximately double the chances of use 1 year later. In addition, prescribers should discuss the long-term plan for pain management with patients for whom they are prescribing either Schedule II long-acting opioids or tramadol.

<sup>1</sup>Division of Pharmaceutical Evaluation and Policy, College of Pharmacy, University of Arkansas for Medical Sciences; <sup>2</sup>Division of Health Services Research, College of Medicine, University of Arkansas for Medical Services.

Corresponding author: Bradley C. Martin, bmartin@uams.edu, 501-603-1992.