

June 2019

Aquatic Invasive Species Change Ecosystem Services from the World's Largest Wild Sockeye Salmon Fisheries in Alaska

Tobias Schwoerer
University of Alaska Anchorage

Joseph M. Little
University of Alaska Fairbanks

Milo D. Adkison
University of Alaska Fairbanks

Follow this and additional works at: <https://cbe.miis.edu/joce>



Part of the [Aquaculture and Fisheries Commons](#), [Natural Resource Economics Commons](#), [Natural Resources and Conservation Commons](#), [Natural Resources Management and Policy Commons](#), and the [Other Environmental Sciences Commons](#)

Recommended Citation

Schwoerer, Tobias; Little, Joseph M.; and Adkison, Milo D. (2019) "Aquatic Invasive Species Change Ecosystem Services from the World's Largest Wild Sockeye Salmon Fisheries in Alaska," *Journal of Ocean and Coastal Economics*: Vol. 6: Iss. 1, Article 2.

DOI: <https://doi.org/10.15351/2373-8456.1094>

This Research Article is brought to you for free and open access by Digital Commons @ Center for the Blue Economy. It has been accepted for inclusion in Journal of Ocean and Coastal Economics by an authorized editor of Digital Commons @ Center for the Blue Economy. For more information, please contact ccolgan@miis.edu.

Aquatic Invasive Species Change Ecosystem Services from the World's Largest Wild Sockeye Salmon Fisheries in Alaska

Acknowledgments

We thank the 56 experts participating in this study and experts who contributed their knowledge to the design of the elicitation. The study was approved by the Institutional Review Board of the University of Alaska and has been performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. The Alaska Sustainable Salmon Fund funded T.S. and J.L. The Alaska SeaGrant funded T.S. The analysis was conducted by T.S., and the paper was written by T.S., J.L. and M. A.. The authors have no conflict of interest to disclose.

1. INTRODUCTION

Invasive species pose a threat to the health of aquatic ecosystems worldwide, and affect ecosystem services important to economic sectors such as agriculture, forestry, and fisheries. While most bioeconomic research focuses on current invasions, there has been little research examining the risk of invasion to intact ecosystems that have not yet been affected. The absence of information on potential future damages has resulted in inadequate management response as other threats appear more pressing (Perrings et al. 2002). Reasons for hampered policy progress are related to the challenges of valuing the economic benefits and costs associated with mitigating the risk of invasive species, as well as the fact that people remain unaware of the issue when not directly or immediately affected by environmental harm (Doelle 2003). In this context, aquatic invasive species are especially worrisome because their presence below the surface of a waterbody can limit early detection. Compared to terrestrial invasive plants, aquatic invasive species are more likely to be established before being detected. The lag in detection lowers the chance of successful eradication and, ultimately, the minimization of long-term costs to society (Perrings et al. 2002). To effectively address the problem, interdisciplinary solutions are needed to quantify and reduce the risk of long-term ecological and economic damages (Shogren 2000).

Despite the continued development of economic methods to value nonmarket and public goods, established methods often rely on the collection of new data. Particular challenges arise with complex valuation exercises designed to model the changes to ecosystem services as a function of human-ecological feedbacks (Finnoff et al. 2005). For example, damage assessments in fisheries often take a production function approach, explicitly modeling relationships among fish populations, human harvesting, and invasion dynamics (D. Knowler 2005; Frésard and Boncoeur 2006). Deciding which complexities to address and which to set aside is a common conundrum for researchers. Economic valuation is most straightforward when an invader has a direct effect on a harvestable resource because the link between ecology and economics can be established through reliable market data (Barbier, Acreman, and Knowler 1997). Notably, within a rapid response model in which timely action results in long-term cost savings, such a simplified valuation approach can provide critical information to managers.

In the absence of biophysical research establishing a linkage between the invader and a harvestable resource, structured expert judgment (SEJ) can be used to bridge the knowledge gap (Cooke 1991). While SEJ is an incomplete substitute for biophysical experimentation, it provides important insights into management. SEJ allows for explicit treatment of uncertainty in cost-benefit analysis and, as such, can inform managers about the economic value of physical experimentation aimed at reducing uncertainty (Peterman and Anderson 1999). SEJ has been widely applied for decades to estimate the human health impacts of air pollution (Morgan et al. 1984), climate change drivers (Morgan 2011), invasive species impacts (Rothlisberger et al. 2012), and changes in fisheries and marine ecosystems (Rothlisberger et al. 2012; Teck et al.

2010). A common challenge in SEJ is the proper aggregation of opinions across individuals with different levels of expertise. Some approaches use equal weights among experts while others use a performance-weighted average based on seed (test) questions with known answers, commonly known as Cooke's method (Cooke 1991; Grigore et al. 2016). However, in cases with high uncertainty which require a diverse group of experts, seed questions can be difficult to frame appropriately (Grigore et al. 2016).¹ The usefulness of seed questions has been questioned in instances where experts from various specialized fields were needed to quantify uncertainty in the medical field (Fischer, Lewandowski, and Janssen 2013; Soares et al. 2011). Many recent studies suggest that equal weighting is preferable to performance-based aggregation as it avoids overweighing counterintuitive results that can lead to biased expert combinations (Clemen 2008; Morgan 2014). Yet, despite this recent research into Cooke's performance weighting scheme, the question remains how to vet expert opinion.

This study avoids the use of seed questions and applies equal weighting after a multi-method approach is used to assure data quality and elimination of inconsistent experts. Cooke's performance scoring is replaced by a coherence check eliminating illogic judgments. The study contributes a vetting technique to an ongoing area of research that focuses on multi-method approaches to expert elicitation (O'Hagan et al. 2006).

The first method uses a discrete choice model (DCM) which is widely used to measure and predict human behavior but has found little application in expert elicitation (McFadden 1973). DCM uses scenarios to observe experts' choices and does not require them to translate knowledge into probabilistic judgments as such assessments can be derived indirectly from estimated individual-specific utility functions (Schwoerer, Little, and Hayward 2018). Therefore, DCM broadens the expert pool and thus allows for later elimination of incoherent experts. Recent research into the trade-offs between increasing the expert pool and the level of expertise informants in the pool bring to the elicitation shows that pool expansion combined with screening outperforms Cooke's method of post-elicitation weighting (Maestas, Buttice, and Stone 2014).

The second method is an interval judgment in the tradition of (SEJ) but without seed questions (Cooke 1991). The SEJ elicits the range of parameter values for the uncertain quantity of interest. A coherence check between the two methods eliminates illogical judgments before aggregating remaining expert judgments applying equal weights. The joint probability distribution is then integrated into the risk analysis framework. The approach avoids some of the

¹ As discussed, bioeconomic analysis of biological invasions often center on measuring existing impacts. In this context, expert elicitation that informs this analysis relies on experts who are knowledgeable or have witnessed existing effects (Rothlisberger et al. 2012). In contrast, expert elicitation studies aimed at predicting what an invader will likely do in an intact ecosystem are more difficult because the uncertainty is higher. This requires the expert pool to be broad because knowledge needs to be attained from different fields of knowledge (Maestas, Buttice, and Stone 2014).

limitations of Cooke's approach, allowing for a more detailed vetting of expert opinion which is useful given the high level uncertainty and varied expertise.

Economic damage assessments in commercial fisheries have gained attention in recent years as marine and coastal ecosystems face increasing human impact through trade, commerce, and development (D. Knowler 2005; Frésard and Boncoeur 2006). In North America, much of the ecological and economic research on invasion impacts to fisheries has focused on the Great Lakes region (Rothlisberger et al. 2012; Wittmann et al. 2014). Bioeconomic research on aquatic invasions in the Great Lakes quantified damages to ongoing invasions yet studies that quantify the value of preventing intact ecosystem services from being invaded are lacking. The risk from invasive species will likely not be eliminated when invasive species populations have established and irreparably impaired ecosystems (Doelle 2003). The opportunity cost of the continued allocation of resources toward already impaired systems results in the forgone prevention of invasions into intact and highly productive ecosystems.

The last wild salmon runs in the world provide a case study for quantifying the potential risk of aquatic invasive species on ecosystems of global economic significance. In Alaska, Pacific salmon (*Oncorhynchus spp.*) are the economic backbone of many coastal communities (Sethi, Reimer, and Knapp 2014; Knapp, Guettabi, and Goldsmith 2013). Wholesale values of Alaska salmon amounted to \$1.28 billion in 2016, only \$50 million less than Pollock, Alaska's largest first wholesale value fishery (McDowell Group 2017). As human presence and activity in ecosystems in the Arctic and Subarctic increases, the threat of invasive species also increases, particularly for highly productive fisheries in this region (Short, Gross, and Wilkinson 2004). Yet, invasive species protection and prevention have received little attention (Schwörer, Federer, and Ferren 2014).

This study was motivated by the discovery of *Elodea spp.* (*Elodea*), an invasive submersed aquatic plant, in three of Alaska's salmon-producing watersheds. It was also recently found at Anchorage's Lake Hood, the world's largest floatplane base. Floatplanes serve as a pathway to spread the plant to remote freshwater sites, most of which are located in salmon habitats that have not yet been invaded (Carey et al. 2016). Lack of biophysical evidence on the ecological effects of *Elodea* for salmon production in Alaska freshwater habitat is related to an overall gap in research examining how invasive aquatic plants affect food web dynamics and fish as well as macroinvertebrate communities (Schultz and Dibble 2012). While non-native aquatic plants play similar roles compared to native aquatic plants, certain traits are problematic, such as rapid growth, the production of biochemicals that influence the growth and survival of other aquatic plant species in its vicinity, and phenotypic plasticity—the ability of genes to produce more than one trait when interacting with different environments. These are all characteristics found in *Elodea* (Schultz and Dibble 2012; Erhard, Pohnert, and Gross 2007). Information on the potential economic risk presented by the *Elodea* is critical for decision-making; yet, the lack of biophysical evidence relating *Elodea* to salmon abundance and productivity has prevented economic analysis (Carey et al. 2016). Expert elicitation related to *Elodea*'s potential effects on

salmon found that invasions occurring in salmon habitat are believed to possibly lead to both negative and positive outcomes for sockeye salmon (Schwoerer, Little, and Hayward 2018). These results underline the need for a bioeconomic risk analysis which weighs the various possible outcomes of an invasion.

The research objective is to inform statewide management decisions for the treatment of *Elodea* and provide a first estimate of the range of damages related to potential invasions. The first section of this paper provides background regarding commercial sockeye salmon fisheries in Alaska. Next, a bioeconomic market valuation is developed that integrates SEJ-derived growth rates for sockeye salmon (*Oncorhynchus nerka*) with a market demand model which uses published commercial fisheries data (ADFG 2018, 2016). The SEJ approach is justified due to lack of data specifying the biological relationship between *Elodea* and salmon which prevents analysis of the structural changes to the stock recruitment relationship for salmon. The model's primary purpose is to inform invasive species managers about the future costs and benefits of taking action to prevent further environmental and economic damages from the distribution of *Elodea* across Alaska.² The range of outcomes suggests that negative consequences outweigh potential positive net benefits to salmon fisheries over a hundred-year timeframe. The magnitude of the most probable damages indicate that substantial investment is justified to keep productive ecosystems free of aquatic invasive species in Alaska. The paper ends with a discussion of the modeling approach and provides policy implications.

2. BACKGROUND

2.1 *Elodea* ecology, management, and history in Alaska

There are two species of *Elodea* in Alaska that are also hybridized (Les and Tippery 2013; Thum 2015 personal communication). *Elodea canadensis* (Canadian waterweed) and *E. nuttallii* (Nuttall's waterweed) are both native to North America between California (35°N) and British Columbia (55°N), but is not native to Alaska (Cook and Urmi-König 1985). Since the ecology of these two plant species is very similar, the following analysis refers to either of these two species as *Elodea*. The plant prefers sand and small gravel substrate in cold, static or slow-moving water to 9 m depth (Riis and Biggs 2003; Rørslett, Berge, and Johansen 1986). *Elodea* is tolerant of a wide range of environmental conditions and has successfully invaded aquatic ecosystems worldwide (Josefsson 2011)³. Cyclical population dynamics have been observed for *E. canadensis* for isolated populations peaking between three and ten years after invasion and declining or even disappearing thereafter (Heikkinen et al. 2009; Mjelde et al. 2012). Sudden

² Note, the aim of the model is different from common population models developed for fisheries management, where structural changes to the stock recruitment relationship would be specified (D. Knowler 2005).

³ *Elodea* is established in the British Isles and many other parts of Europe (Heikkinen et al. 2009) *Elodea* is established in the British Isles and many other parts of Europe (Heikkinen et al. 2009)

collapses remain unexplained but have been observed throughout Europe (Simberloff and Gibbons 2004; Mjelde et al. 2012). In regulated rivers in its native range, *Elodea* has been found to encroach on spawning sites of Chinook salmon (*Oncorhynchus tshawytscha*) (Merz et al. 2008).

Common human-related pathways of introduction include the aquarium trade, boats, and floatplanes (Sinnott 2013; Strecker, Campbell, and Olden 2011). Natural dispersal includes flooding and wildlife transport (Spicer and Catling 1988; Champion, Winton, and Clayton 2014). In Alaska, *Elodea* reproduces vegetatively with stem fragments surviving desiccation and freeze (Bowmer, Jacobs, and Sainty 1995). *Elodea* has some of the highest fragmentation and regeneration rates among aquatic invasive plants causing rapid dispersal which presents severe challenges for mechanical removal (Redekop, Hofstra, and Hussner 2016). Possible management actions include draining and drying, herbicides, the introduction of herbaceous fish, and mechanical removal through suction dredging or hand pulling (Hussner et al. 2017). For the purpose of eradication, Fluridone and Diquat are herbicides that are most effective while mechanical methods cause populations to rapidly spread (Josefsson 2011). In Alaska, Fluridone and Diquat have eradicated *Elodea* in three waterbodies (Morton 2016). At concentrations around 6 ppb, Fluridone selectively removes *Elodea* with few non-target effects (Kamarianos et al. 1989; Schneider 2000).

In Alaska, *Elodea* was discovered in Fairbanks (Interior Alaska) in 2010, drawing attention to an already established, but until then largely ignored, population in Cordova (Figure 1, Gulf). New infestations have been found every year since 2010. Aquarium dumps are the likely vector near urban locations, while floatplanes are the most likely pathway responsible for long-distance dispersal into remote roadless locations (Hollander 2014). It came as no surprise when, in 2015, *Elodea* was detected in Lake Hood (Figure 1, Cook Inlet), the world's largest seaplane base (Hollander 2015). The discovery of *Elodea* 90 river miles downstream from an unmanaged infestation in Fairbanks was likely caused by flooding (Friedman 2015). The increasing frequency of new discoveries associated with previously unknown pathways underscores the importance of the issue for Alaska.

2.2 Commercial sockeye salmon fisheries

Alaska's commercial sockeye salmon fisheries can be regionally divided into five large watersheds (Figure 1) (USGS 2017). The regions include Bristol Bay and Kuskokwim in western Alaska; Cook Inlet in Southcentral Alaska; Kodiak which encompasses the island of Kodiak and the southern coast of the Alaska Peninsula; and the Gulf which includes the Kenai Peninsula's Gulf coast, Prince William Sound, and watersheds supporting the Copper and Bering River

fishing districts.⁴ These regions have varying seafood processing capacity, run sizes, harvest levels, and prices that depend on global market forces.

Since the 1990s, Alaska salmon prices have experienced downward pressure caused by the rapid and sustained growth of farm raised salmon. Yet over the past decade, prices have recovered due to marketing efforts aimed at wild and sustainably-caught Alaska salmon. Also, disease outbreaks in salmon farms elsewhere have had positive price effects on Alaska salmon (Knapp, Roheim, and Anderson 2007).⁵ In 2014, wild salmon comprised about 30% of global salmon production by volume. Alaska sockeye salmon production is associated with the largest share (65%) of wild salmon sold on global markets. Of this share, 37% of wild Alaska sockeye are caught in Bristol Bay (McDowell Group 2015). With the Bristol Bay sockeye salmon fishery, it can be argued that Alaska sockeye production influences global prices (Knapp, Roheim, and Anderson 2007). Table 1 shows historical wholesale prices for the four main product categories for sockeye salmon—frozen, fresh, canned, and other.⁶ Given a globally traded product, variations in price exist and are correlated across regions (Table 2).




Figure 1. Watersheds supporting commercial sockeye salmon fisheries that were part of this study. This study region was selected because *Elodea* is present in the Cook Inlet and Gulf regions.

⁴ The geographic focus is aimed at currently known *Elodea* infestations, excluding Southeast Alaska's salmon fisheries from the analysis.

⁵ Alaska's constitution prohibits salmon farming in state waters within three nautical miles.

⁶ While there are additional subcategories, the analysis focuses on these four categories for which wholesale prices are published (ADFG 2016b).

Table 1. Fisheries characteristics by region, 2006-2015

Region	Sockeye harvest ('000 lbs)			Sockeye mean (SD) wholesale prices (real \$/lbs) ^{a)}			
	mean	min	max	canned	Frozen	fresh	other ^{b)}
Bristol Bay	154,193	92,000	184,792	\$ 3.66 (2.4)	\$ 4.01 (2.3)	\$ 2.71 (1.1)	\$ 7.54 (2.5)
Cook Inlet	18,920	12,266	36,216	n/a ^{e)}	\$ 4.19 (3.0)	\$ 3.40 (2.5)	\$ 8.24 (6.3)
Gulf	16,386	8,004	24,785	\$ 5.69 (2.9)	\$ 3.79 (2.7)	\$ 4.20 (2.4)	\$ 6.30 (2.9)
Kodiak	11,980	7,692	17,007	n/a ^{e)}	\$ 3.22 (2.8)	\$ 3.12 (1.3)	n/a
Chignik ^{c)}	9,338	4,125	17,889	n/a	\$ 3.22 (2.8)	n/a	n/a
Kuskokwim ^{d)}	746	329	1,379	n/a	\$ 1.11 (1.2)	n/a	n/a

a) Mean (standard deviation) in 2015 USD adjusted for inflation using the U.S. Consumer Price Index. b) Salmon roe products Sujiko in Bristol Bay and mainly Ikura in Gulf. For Cook Inlet: fillets with skin no ribs. c) Assumes Kodiak prices due to lack of data. The analysis treats the Chignik fishing district as a separate region because of available harvest data. However, results are combined with Kodiak. d) Prices reported for the exclusive economic zone (EEZ) were used due to lack of data. e) Region stopped production of this product or production is very inconsistent from year to year due to swings in run size. Source: Alaska Department of Fish and Game Fisheries Management Annual Reports and Commercial Operators Annual Reports.

Table 2. Correlation among regional wholesale prices for sockeye salmon, 2006-2015

	Bristol Bay	Cook Inlet	Kuskokwim	Gulf	Kodiak	Chignik ^{a)}
Bristol Bay	1.00					
Cook Inlet	0.89	1.00				
Kuskokwim	0.06	0.26	1.00			
Gulf	0.78	0.89	0.44	1.00		
Kodiak	0.80	0.90	0.18	0.73	1.00	
Chignik ^{a)}	0.80	0.90	0.18	0.73	1.00	1.00

a) Due to lack of data, assumes prices behave similarly to Kodiak. Note, correlations are based on just one product: frozen headed and gutted sockeye salmon. Prices published for the exclusive economic zone (EEZ) are used for the Kuskokwim due to lack of location-specific price data. Note, Spearman's rank-ordered coefficients are more appropriate for modeling correlation among distributions compared to Pearson's correlation coefficients (Palisade Corporation 2016). Source: ADFG (2016b)

There are also regional differences in seafood processors and the sockeye salmon products they produce. Table 3 shows region-specific production shares and overall processing yields. The latter is an average equal to the ratio of output weight sold over input weight bought by processors (Knapp, Roheim, and Anderson 2007).

Table 3. Production shares and processing yield by region, 2006-2015

Product	Bristol Bay ^{a)}	Cook Inlet ^{b)}	Kuskokwim	Gulf ^{c)}	Kodiak ^{b)}	Chignik ^{a)}
Canned	0.32			0.34		
Fresh	0.02	0.12		0.08	0.12	
Frozen	0.64	0.86	1.00	0.57	0.88	1.00
Other	0.02	0.02		0.01		
Processing yield ^{d)}	0.70	0.78	0.75	0.71	0.78	0.75

a) McDowell Group (2015). b) Author estimates based on observed historic prices (ADFG, 2016b; Knapp et al., 2007).c) Knapp et al. (2007). d) Weighted using product-specific yields: canned 0.59, fresh 0.97, frozen (headed & gutted) 0.75, other 0.75 (Knapp et al., 2007; author assumptions for other).

3. METHODS

The damage assessment approach consists of two components. The first describes the elicitation, evaluation, and aggregation of expert opinion on the impacts of *Elodea* on sockeye salmon productivity in Alaska. Two methods were used to accomplish this, a Discrete Choice Model (DCM) (Schwoerer, Little, and Hayward 2018) and Structured Expert Judgment (SEJ) (Cooke 1991). The second component is the bioeconomic model used to estimate changes in consumer surplus (Freeman 2003). This approach follows the methodology of previous assessments of economic impacts from invasive species in the Great Lakes (Rothlisberger et al. 2012). The section ends with an outline of the biological and economic parameter values used in estimating potential damages to sockeye salmon fisheries.

3.1 Expert judgment

In order to quantify uncertainty about *Elodea*'s effects on salmon habitat and population the approach drew on broad expertise from three areas of knowledge: (1) Pacific salmonids in freshwater habitats, (2) the ecological role of submersed aquatic vegetation, and (3) freshwater aquatic invasive plants. An extensive literature review of nearly 300 sources identified an expert pool of 111 individuals with combined knowledge in all of these areas. Expert selection was based on the number of citations in peer-reviewed publications using Google Scholar. Due to the localized issue of *Elodea* in Alaska and the small number of potential experts, the expert pool was expanded to include fishery biologists, fishery scientists, fish habitat biologists, and invasive species specialists from state and federal resource management agencies. These individuals brought knowledge on localized variability and local observations to the expert pool as all of them had or continue to work with Alaska salmon, aquatic systems, or invasive species.

For a pretest, we initially followed Cooke's method which asks experts questions with known answers, also called seed or calibration questions (Cooke, 1991). Based on experts' performance in these seed questions, their judgment is later weighted giving more weight to well performing

experts. Several attempts to employ Cooke's (1991) method failed because of the difficulty finding questions appropriate to all diverse areas of expertise. Therefore, instead of relying on inappropriate seed questions the DCM and SEJ were used to evaluate experts and test for coherence of opinion.

The DCM asked experts to choose from hypothetical salmon habitat scenarios that they believed would result in the long-term persistence of salmon. The scenarios varied in their description of habitat and invasion characteristics (Schwoerer, Little, and Hayward 2018).⁷ Based on the DCM data, each expert's probability of choosing invaded over uninvaded habitat for persistent salmon populations was estimated (Schwoerer, Little, and Hayward 2018). In a second follow-up exercise, a SEJ was used to ask experts to state intervals for the annual average sockeye growth rates to be expected in *Elodea*-invaded habitat. Both the DCM and SEJ included an extensive background document informing experts about *Elodea*'s known ecological effects and were aimed at bridging knowledge gaps and reducing overconfidence in the interval judgment (Speirs-Bridge et al. 2010).⁸ In the SEJ the annual average sockeye growth rate was referred to as "salmon production over many life cycles, manifesting itself as a long-term trend in abundance" (McElhany et al. 2000). The elicited growth rates apply to the whole population of sockeye salmon and do not specify *Elodea*'s effects on specific age structures. As such, the first question in the elicitation specified the 25th and 75th percentile of the probability distribution related to the annual average sockeye growth rate in *Elodea*-invaded habitat, the second question established the tails of the distribution, and the third question showed the median. The fourth question tested expert's comprehension of the task (Speirs-Bridge et al., 2010) and was also used to further eliminate experts from the pool (see below).

Q1. Imagine Alaska's sockeye salmon systems would be invaded with *Elodea* and you had long-term population records with estimated sockeye growth rates for a random sample of 100 of these sockeye systems. What range of typical sockeye growth rates would you expect to see, that is, rates you would see about half of the time?

Q2. What is the very lowest and very highest sockeye growth rate you would expect to see, if Alaska's sockeye salmon systems would be invaded with *Elodea*? Think about the extreme cases, in other words the tails of the distribution.

Q3. What is your best guess for the sockeye growth rate you would expect to see most often, if Alaska's sockeye salmon systems would be invaded with *Elodea*?

⁷ Environmental characteristics included location of *Elodea* within the salmon system, description of the salmon system, dissolved oxygen levels, predation, prey abundance, and other factors.

⁸ Even though the existing literature describes the reductions in overconfidence relating specifically to the 4-step interval elicitation procedure, the more elaborate nature of the scenario-based approach prior to the interval judgment is believed to have similar overconfidence-reducing effects. While a test of this assumption could be subject to future research, it is outside the scope of this study.

Q4. For sockeye salmon, what sockeye growth rate would cause you to be concerned about extirpation of the population? Please specify in % and use a "-" (minus sign) for decline rates.

Based on the probability of salmon persistence in *Elodea*-invaded habitat estimated through the DCM and the stated sockeye growth rates in the SEJ a coherence check was applied. The purpose of the coherence check was to identify experts whose probability of sockeye salmon persistence in *Elodea*-invaded salmon habitat from the DCM was consistent with the annual average sockeye growth rates they believed would be possible in *Elodea*-invaded salmon habitat stated in the SEJ. A logical and consistent expert either indicated a lower than 50:50 chance of persistence in *Elodea*-invaded habitat in the DCM exercise and stated a negative median growth rate in the SEJ or a higher than 50:50 chance of persistence in the DCM and a positive median growth rate in the SEJ. Inconsistent experts (persistence/negative or extirpation/positive) were excluded from further analysis.

Assuming a standard normal distribution, individual expert's interval judgments from Q1 to Q3 were combined applying equal weights to each expert (Cooke 1991). Figure 2 presents the expert vetting and aggregation process in more detail.




Figure 2. Expert vetting process used for aggregating expert opinion on annual average sockeye growth rates in habitat invaded by *Elodea*.

3.2 Bioeconomic model

Many commercial salmon fisheries in Alaska create significant economic rent indicated by the high permit prices in for example the Bristol Bay salmon fisheries (Knapp, Guettabi, and Goldsmith 2013). Biological invasions in this context could be seen as dissipating rent, therefore, we follow a benefit approach to the economic valuation of ecosystem services impaired by a biological invasion (Freeman 2003).

If *Elodea* changes the provisioning of ecosystem services—the amount of harvestable sockeye salmon—the introduction of *Elodea* changes the benefits consumers derive from the resource. Consumer surplus provides an economic measure of ecosystem benefits and is calculated by taking the difference between the maximum amount consumers are willing to pay for the resource and what they are actually paying. For example, if the consumer only pays \$6 per pound for sockeye salmon, but would be willing to pay up to \$10 per pound, the difference of \$4 is the benefit to that consumer, which aggregated across society is equal to consumer surplus.

The model calculated the change in consumer surplus that resulted from a change in annual harvest and a consequential change in the real price per pound (\$/lbs), assuming a linear demand function (Freeman 2003). Since the SEJ-derived intervals entailed positive and negative sockeye growth rates, this approach allowed for potential positive and negative net changes in consumer surplus. These net changes imposed by quantity changes in annual harvest were equal to the change in the area under the ordinary (Marshallian) demand curve and equal to the consumer surplus in year t minus the consumer surplus in year $t+1$. In mathematical terms, annual damages per region were expressed as follows:

$$\begin{aligned}\Delta CS_{t+1} &= CS_t - CS_{t+1} \\ &= \frac{\gamma}{2}(h_t(p' - p_t) - h_{t+1}(p' - p_{t+1}))\end{aligned}\quad (1)$$

where γ was processing yield, h was sockeye harvest in lbs, p was the real (inflation-adjusted) per lbs wholesale price for sockeye salmon in 2015 USD received by Alaska primary processors. Prices were weighted by sockeye product ratios commonly observed in the Alaska processing sector (Table 3). The choke price at which demand ceases is equal to p . Using the own-price elasticity of demand, ε , the choke price equals $p' = h_t/\varepsilon + p_t$. Further, harvest in period $t+1$ is expressed as a function of SEJ-derived sockeye growth rates θ , thus $h_{t+1} = f(h_t, \theta)$. After substituting and rearranging, Equation 1 becomes

$$\Delta CS_{t+1} = \frac{\gamma}{2} \left[(h_t - f(h_t, \theta)) \left(\left(\frac{h_t}{\varepsilon} + p_t \right) + \frac{f(h_t, \theta) p_t}{h_t \varepsilon} \right) - p_t h_t \right]. \quad (2)$$

The functional response of harvest to an *Elodea* invasion was represented by $h_{t+1} = f(h_t, \theta)$. Consistent with common practice in fisheries modeling, catch was assumed to be proportional to stock size and fishing effort (Haddon 2011).⁹ Year-by-year changes in harvest were modeled using density-dependent population dynamics in logistic form such that harvest levels at $t+1$ equaled $h_{t+1} = h_t(1 + \theta(1 - h_t/K))$, where K is the ecologically limited harvest. Due to the seasonal reproduction of salmon, the discrete time model with an annual time-step is well suited for modeling the growth changes in salmon (Haddon 2011).

⁹ Under this assumption the catchability of the fishing fleet does not change over time or stock size.

The logistic growth model describes the population dynamics for an entire population of salmon irrespective of age-class, making it consistent with the SEJ-derived population growth rates. Due to a number of limitations the logistic growth model is not often used to describe population dynamics in fisheries (Larkin 1977), but it does have the advantage of being useful in situations where data is limited (Beverton and Holt 1957; Haddon 2011). The logistic growth model differs from commonly used population models like the Ricker model in how it describes population change at very high population densities (Ricker 1975). In the logistic growth model, growth at very high densities declines more rapidly, an assumption supported by the encroachment effects of *Elodea* observed on spawning adult salmon (Merz et al. 2008). In addition, the limiting environmental conditions of *Elodea*'s encroachment into salmon spawning beds is captured by the harvest limitation (K) in the logistic model but is lacking in an exponential growth model.

Under exponential growth, the expert-elicited positive effects of *Elodea* for salmon would result in runaway growth, or the believed negative effects would cause short-term extirpation—both biologically unrealistic outcomes. While long-term persistence is not guaranteed under the logistic growth model, the model indicates that, despite environmental perturbation, salmon populations can persist long-term. The invasion of *Elodea* in the British Isles recently reached its ecological limit, after 65 years since introduction (NBN 2015). The NBN data showed that landscape-wide spread of an *Elodea* invasion over a much longer timeframe compared to the 20-year time horizon considered for persistent salmon populations in invaded habitat (Peterson et al. 2008). Research on salmon habitat and *Elodea* occurring in its native range suggests that the effects of *Elodea* encroaching on spawning adults has had incremental rather than catastrophic consequences (Merz et al. 2008). The effects of *Elodea* on salmon in the invasive range may manifest themselves over a longer timeframe without immediate catastrophic outcomes. Moreover, the boom and bust cycle of *Elodea* populations can lead to temporarily more or less pronounced biological effects for different life stages over time (Simberloff and Gibbons 2004). For these reasons, the logistic growth model was chosen to describe the biological relationship between *Elodea* and salmon.

The model was initialized in year zero by the pre-invasion historical sockeye harvest, h_0 , and the pre-invasion historical wholesale prices per region, p_0 . Potential economic damages to commercial sockeye salmon fisheries were expressed over a 100-year time horizon in the following two ways. First, the annual changes in consumer surplus estimated by Equation 2 are converted into net present value terms (NPV)

$$NPV = \sum_{t=0}^{100} \Delta CS_t (1+d)^{-t} \quad (3)$$

where d is the real social discount rate and takes into account the public's time preference related to natural capital. NPV is a measure for natural capital, from which a constant flow of ecosystem services can be calculated such that

$$NPV_{annual} = NPV \frac{d}{1 - (1 + d)^{-100}}. \quad (4)$$

NPV measures the value of the stock of natural capital (Equation 3) at time t , annualized NPV is the constant flow of ecosystem services associated with natural capital (Equation 4). Put simply, natural capital is wealth, ecosystem services are income. The regional estimates were summed to show combined statewide damages.

Several simplifying assumptions were made relating to the economic and environmental conditions of the commercial salmon fisheries and the invasion by *Elodea*. First, the analysis estimated potential damages to fisheries should the regions become invaded by *Elodea* in the first year of a 100-year time horizon and remain unmanaged. Therefore, the estimated damages were hypothetical for regions that have not yet been colonized by *Elodea*. Second, the predicted changes caused by *Elodea* only change the weight of fish landed and do not alter the consumer demand function. Third, market conditions were assumed to be in equilibrium so that there were no incentives for harvesters and processors to enter or exit the market. Similarly, participation by fishing permit holders did not change over time. Fourth, wholesale prices were assumed to proxy prices for end consumers. Analysis based on retail prices would have been more difficult, complicated by exchange rates and a number of other issues. Retail prices exhibit variations which reflect factors, such store location, parking, and the availability of other products, which can't be attributed to salmon (Knapp et al., 2007). Lastly, from an economic perspective, the fishery was assumed to be optimally managed, meaning the ecological and economic systems were in equilibrium throughout the 100-year time horizon. This assumption ignores various management inefficiencies such as over-capitalization which remains an issue for Alaska salmon fisheries due to regulations resulting in a derby-style "race for fish" (Knapp and Murphy 2010). Alaska commercial salmon fisheries, however, are sustainably managed as certified by the Marine Stewardship Council (MSC) (MSC 2017; Clark et al. 2006).

3.3 Model simulation and parameter assumptions

The deterministic nature of the economic valuation did not require a simulation approach. However, Monte Carlo simulation was used to estimate NPV for a range of parameter inputs. Historical data was used to determine the range of salmon harvests and prices. Due to the deterministic nature of the model and contrary to common stochastic modeling approaches, the simulation held these uncertain parameters fixed over the model's time horizon. The simulation tested up to 10,000 possible input assumptions for each uncertain parameter, generating a distribution for Equations 3 and 4. The simulation stopped when there was a 95% chance that the mean NPV was within $\pm 3\%$ tolerance of its true value (Palisade Corporation, 2016b).¹⁰

¹⁰ Sampling type: Latin Hypercube, random number generator: Mersenne Twister.

Uncertain parameters include the expert-elicited annual average growth rate for sockeye salmon, θ , the pre-invasion sockeye harvest in year zero, h_0 , pre-invasion wholesale prices, p_0 , own-price elasticity of sockeye demand, ε , and the social discount rate, d . For the annual average growth rate for sockeye salmon, θ , the equally-weighted percentiles were used to fit a normal distribution describing the joint probability density function for θ . A normal distribution is suitable for this purpose because many unknown ecological processes are likely at play in *Elodea*-invaded habitat and average out over a large sample (Hilborn and Mangel 1997). For example, long-term variation of salmon returns is also driven by Pacific climate variability and other factors (Hare, Mantua, and Francis 1999). Since the return of salmon from different populations can vary within the same year, each harvest distribution is assumed to be independent of all others (Schindler et al. 2010).

To describe the variation of historical harvest, region-specific commercial sockeye harvest records in pounds landed from 2006 to 2015 were used to fit a uniform distribution (Table 4). For the purpose of testing different model assumptions surrounding historical harvest, this non-informative distribution was found to best accommodate this purpose across regions. The lognormal distribution is commonly used in economics to describe the distribution of income, wealth, and prices and was used to specify the pre-invasion wholesale price in each region (Table 1 and Table 4) (Aitchison and Brown 1976). The correlation of prices among regions was modeled based on estimated Spearman's rank-order correlation coefficients observed between 2000 and 2015 (Table 1). To derive this correlation, the model generated rank-correlated pairs of prices for two regions at a time following a distribution-free approach to induced rank correlation (Iman and Conover, 1982; Palisade Corporation, 2016a).

Reliable market data on prices and quantities was then used to derive estimates of economic benefit. In order to measure changes in consumer surplus, the approach requires an estimate of the own-price elasticity of demand. The elasticity describes how responsive consumer demand for salmon is to changes in the price of salmon (Freeman 2003). With this information, associated changes in consumer surplus can be estimated. Unfortunately, there are no specific estimates of own-price elasticities for Alaska sockeye salmon. Estimates from elsewhere in North America serve as a proxy. A variety of sources were consulted that estimated the elasticity of demand for fresh and frozen wild sockeye salmon in the Pacific Northwest, Oregon, or Canada (DeVoretz 1982; Wang 1976; Johnston and Wood 1974; Swartz 1978). Due to farmed salmon dominating world markets, more recent demand investigations have focused on farmed instead of wild salmon demand (Asche, Bard, and Atle 2019; Andersen, Roll, and Tveteras 2008). All estimates indicated elastic demand, $|\varepsilon| > 1$, and ranged between a minimum of -12.78 and a maximum of -1.472 (DeVoretz 1982; Wang 1976). A uniform distribution was applied using maximum and minimum values related to elasticity estimates as bounds to reflect the uncertainty in these historical estimates (Table 4). There are several arguments that would support higher elasticities $|\varepsilon| > 1$; the most important of which is the availability of very close substitutes to wild sockeye salmon, such as coho, pink, or chum salmon. Wild sockeye is also considered a normal good where demand increases with rising income and vice versa. To the

contrary, brand loyalty to a wild and sustainably harvested product is an argument for more inelastic demand if current marketing efforts and consumer awareness continue (McDowell Group 2015).

Since our analysis is focused on changes to consumer welfare and the benefits to society from public investments in cleanup that are more or less uncertain for current and future generations, a triangular distribution of discount rates was used to account for such uncertainty. Much uncertainty does not surround the cost of capital due to Alaska's Aa2 credit rating, but rather relates to whether successful statewide eradication of elodea can be achieved. Therefore, the distribution of the marginal opportunity cost of capital chosen for the analysis included the risk free rate as well as a rate much higher than the risk free rate, comparable to the 6% used by Rothlisberger et al. (2012) for their analysis of economic damages caused by the invasion of *Dreissena* mussels in the Great Lakes. Also, the real 30-year social discount rate recommended by OMB ranges between 1% and 6% (OMB 2016). The triangular distribution with a most likely rate of 3% is consistent with best practices in financial valuation (Winston and Albright 2016). A distribution also reflects varying time preference rates observed across society. This approach is suitable for intergenerational time horizons and in cases where damages accrue in the private as well as public sectors as in the *Elodea* case suggests the use of multiple discount rates (Arrow et al. 2013; Baumgärtner et al. 2015).¹¹ Table 4 summarizes the model parameter assumptions used in the analysis.

¹¹ The upper bound of 6% reflect real annual rates of return for Alaska's commercial salmon fisheries (Huppert, Ellis, and Noble 1996). The lower bound is consistent with recent research which suggests that impacts to ecosystem services should be discounted at much lower rates compared to impacts related to manufactured capital (Baumgärtner et al. 2015). A reduction in harvest due to an *Elodea* invasion could result in fishing vessels being on dry dock rather than fishing with private opportunity costs to capital.

Table 4. Bioeconomic model parameters

Parameter	Units	Region-specific	Specification	Source
Annual average sockeye growth rate of salmon occupying <i>Elodea</i> -invaded salmon habitat, θ	decimal	no	Normal (-0.0522 , SD: 0.1388)	This study
Pre-invasion harvest, h_0	lbs	yes	Normal (see Table 1)	ADFG 2016a
Pre-invasion wholesale price ^{a)} , p_0	2015 USD	yes	Lognormal (see Table 1)	ADFG 2016b
Own-price elasticity of demand, ϵ	decimal	no	Uniform (-12.78 , -1.472)	Wang 1976; DeVoretz 1982
Ecological limit of sockeye harvest, K	lbs	yes	Max. hist. harvest (see Table 1)	ADFG 2016a
Processing yield, γ	decimal	yes	See Table 3	Knapp et al. 2007
Real social discount rate, d	decimal	no	Tri (0.01 , 0.03 , 0.06)	Rothlisberger et al. 2012, OMB 2016

a) Weighted by the region-specific sockeye product amounts for frozen, canned, fresh, and other product categories (see Table 3).

2.4 Sensitivity analysis

A sensitivity analysis was used to test the robustness of the estimate and contained two parts. First, Latin Hypercube sampling was used to randomly draw input parameter values from the distributions outlined in Table 4 (Palisade Corporation 2016). Second, the combined uncertainty of all the input parameters was then measured by assessing the variance of the loss distribution using 100,000 iterations.

4. RESULTS

4.1 Coherence check of expert judgment

A total of 56 experts participated in the DCM and 44 experts took part in the SEJ focused on the judgment of annual average sockeye growth in *Elodea*-invaded habitat. Five of the remaining experts were unreachable or had retired by the time of the follow-up interval judgment. Six of the remaining experts were unwilling to provide interval judgments and stated lack of knowledge or unfamiliarity with sockeye growth rates as reasons for not responding. One of the remaining experts did not complete the full interval judgment. Of the 44 participating experts in the interval judgment, five experts stated positive sockeye growth rates when asked about growth rates of populations that would cause them to be concerned about extirpation. Consequently, these experts were eliminated. The scatterplot in Figure 3 A shows how the 39 remaining experts varied in their opinion between the DCM and the SEJ. The vertical axis indicates each expert's probability of sockeye persistence as estimated by the DCM and the horizontal axis represents

the expert's best estimate for the annual average sockeye growth rate elicited in the SEJ. A total of eight experts provided illogical estimates in the SEJ and are illustrated by triangular markers (Figure 3 A). The eight experts were eliminated before aggregating their interval judgments to form a joint and normal probability distribution (Figure 3 B). This normal probability distribution depicted a 37% chance of observing positive annual average sockeye growth rates in *Elodea*-invaded habitat (Mean: -0.0522 , SD: 0.1388). This distribution of the combined expert opinion incorporated both *Elodea*'s negative and positive potential growth effects on sockeye salmon.




Figure 3. Exhibit A) presents the coherence check comparing expert's probability of sockeye persistence as calculated by the DCM with annual average growth rates for sockeye populations in *Elodea*-invaded salmon habitat from as stated in the SEJ. Exhibit B) shows each coherent expert's believed distribution of average annual sockeye growth rates in *Elodea*-invaded salmon habitat (25th, mean, and 75th percentile).

4.2 Potential economic loss

Figure 4 illustrates the non-discounted annual loss in consumer surplus (in 2015 US dollars) over the 100-year time period for Bristol Bay. Annual loss is not only increasing, it becomes increasingly more uncertain in future years. Consistent with expert opinion identifying some potential for positive sockeye growth and positive net benefits in *Elodea*-invaded salmon habitat, the 90% uncertainty bars extend below the zero-damage line.

Equation 3 was used to calculate the net present value of potentially lost natural capital which amounted to a mean of \$5.1 billion for all five regions combined (90% CI: $-\$4.6$ billion in net benefits, \$20.0 billion in damages) (Table 6). A more detailed look at the NPV distribution and 90% uncertainty range show that despite the 35% probability of positive ecosystem services (negative damages), the upper bounds of damages were much larger than the potential benefits. The associated constant annual loss in ecosystem services for all five regions combined amount

to a mean loss of \$159 million annually (90% CI: -\$144.4 million in net benefits, \$577.3 million in damages) (Table 6). Across the five regions, this estimate ranged between a mean of \$0.2 million in annual damages in the Kuskokwim to \$111.9 million in annual damages in Bristol Bay (Table 6). Considering that the annual real (inflation-adjusted) wholesale value of Bristol Bay sockeye ranged between \$221 and \$458 million in the past ten years, the estimated annual damages from *Elodea* would be equivalent to a third to two thirds reduction in wholesale value (McDowell Group 2015).




Figure 4. Annual non-discounted consumer surplus loss over the next 100 years assuming Bristol Bay’s sockeye salmon habitat is invaded by *Elodea* and remains unmanaged.

Table 1. Potential damages to commercial sockeye fisheries by region (\$ million)

Region	Change in ecosystem services (NPV _{annual})			Change in natural capital (NPV)		
	Mean	5%	95%	Mean	5%	95%
Bristol Bay	111.9	-101.0	401.6	3,558.6	-3,175.8	13,957.5
Cook Inlet	23.3	-26.1	95.0	739.4	-834.4	3,196.9
Gulf	15.0	-17.3	56.3	474.8	-547.8	1,920.7
Kodiak	9.0	-8.5	38.8	287.5	-267.7	1,283.5
Kuskokwim ^{d)}	0.2	-0.3	1.0	6.8	-7.8	34.3
Total	159.4	-144.4	577.3	5,067.1	-4,589.5	20,029.7

Reasons for the differences are linked to the varying regional characteristics discussed above. Bristol Bay shows the largest economic impact followed by Cook Inlet, Gulf, Kodiak, and

Kuskokwim regions (Table 6). Important to note is that across all regions, the most probable outcomes are showing loss.

5. SENSITIVITY ANALYSIS

The SEJ-derived annual average sockeye growth rates in *Elodea*-invaded salmon habitat contributed to more than half of the variance observed in the simulated NPV distribution. This result is not surprising considering the large uncertainty in this parameter. Some of the lowest growth rate assumptions increased the mean loss by over \$8 billion, while some of the largest growth rates decreased the mean loss by \$8 billion, resulting in a mean of \$3.4 billion in net benefits. The discount rate and Bristol Bay wholesale price for frozen sockeye product contributed much less to the variance than the growth rate (Table 7). A Bristol Bay price assumption of \$18.59/lbs increased losses by \$4.4 billion, whereas \$0.82/lbs reduced losses by \$2.2 billion (Table 2.7). Sensitivity of model outcomes to assumptions surrounding the own-price elasticity of demand were insignificant and contributed less than 1% to variance in NPV, less than wholesale prices in other regions (not shown).¹² As expected, the sockeye growth rate and discount rate both were negatively correlated with damages whereas wholesale prices were positively correlated (Figure 5). The effect of positive sockeye growth rates in *Elodea*-invaded salmon habitat (θ) on the mean NPV was reduced by the harvest constraint, creating a convexity of the solid line in Figure 5 above the 70th input percentile. Initial harvest assumptions did not significantly contribute to variance in NPV.

Table 2. Sensitivity of annualized damage estimates to parameter assumptions with the largest influence on simulation outcomes

	% Contribution to variance	Change in mean NPV (billions 2015 USD) ^{a)}	
		Lowest input assumption	Highest input assumption
Annual average sockeye growth rate, θ	52.8%	8.8	-8.7
Discount rate, d	5.6%	5.0	-2.9
Price for frozen product in Bristol Bay, p_0 ^{b)}	6.6%	-2.2	4.4
Own-price elasticity, ϵ	0.22%	-0.7	0.9

a) Mean NPV equal to \$5.188 billion. Changes are calculated holding all other parameters constant at their mean levels. b) Similar changes relate to other frozen product prices in the Gulf, Chignik, Cook Inlet, and Kodiak regions.

¹² A test using a triangular distribution with a peak of -4.82 lead to mean damages within 0.1% of the shown result.an




Figure 5. Change in mean NPV for all regions combined dependent on percentile changes related to the three parameters contributing the most to variance in the NPV distribution.

6. DISCUSSION

This study offers a damage forecast related to *Elodea*'s effect on sockeye salmon fisheries that stands in stark contrast to comparable bioeconomic damage assessments of aquatic invasive species elsewhere. This study followed methodology used in estimating fisheries damages of aquatic invasive species in the Great Lakes and as such offers a direct comparison of damages (Rothlisberger et al. 2012; Rosaen, Grover, and Spencer 2012). In their highest of four scenarios, Rothlisberger et al. (2012) estimated the cumulative damages related to *Dreissena* mussels invading the Great Lakes through marine shipping at \$2.16 billion over the next fifty years. Weighing the damages from biological invasions in the Great Lakes to the transportation savings associated with shipping, the primary pathway for invasions, the damages outweigh the transportation cost savings by \$750 million. In contrast, this study estimates the cumulative mean damages to commercial salmon fisheries in Alaska at \$5.1 billion over the next 100 years.¹³

There are a number of contributions made by this examination. First, the analysis explicitly addresses uncertainty in the predicted damage distributions by applying a unique coherence check for vetting expert opinion. The approach taken is more conducive to a larger expert pool because it does not require probabilistic statements and avoids the use of seed questions, which

¹³ Rothlisberger et al. (2012) presented cumulative damages, thus, the annual damages were recalculated to provide comparison. Also note, the economic benefits of *Elodea*, in specific the non-market values related to use of *Elodea* in aquariums is insignificant. In addition, the positive effects of *Elodea* for salmon fisheries were accounted for by this study.

are often difficult or impossible to frame for experts from different fields. Thus, the multi-method approach to expert elicitation offers wider application across specialized fields. Additional options for combining expert opinion include Bayesian approaches or the use of geometric means which are limited to positive values (logarithmic opinion pool) (O'Hagan et al. 2006). While these methods vary in ease of application, there is disagreement on how well each one performs in specific situations (Hammit and Zhang 2013; O'Hagan et al. 2006; Morgan 2014; Clemen 2008). Besides the simple application of the equal weights method for post-elicitation, the multi-method approach performed well but will require further performance testing in a broader range of applications (Morgan 2014).

Second, the study informs invasive species management not only about the potential negative economic consequences of an invader but also accounts for the invader's ability to aid in the growth of a harvestable resource. Few studies assessing the economic impacts of biological invasions account for multi-directional effects of an invasion (Gleditsch and Carlo 2011). The observed range of expert opinion on *Elodea*'s growth effects on sockeye salmon is consistent with research pointing towards various positive and negative effects of aquatic invasive plants on fish species and supports the expert elicitation approach taken (Schultz and Dibble 2012). Third, the benefit approach to valuation used publicly available data on fish harvest and wholesale prices, providing a more reliable method when compared to economic valuation techniques based on stated preference approaches (e.g., contingent valuation) (Freeman 2003). Fourth, the study is in contrast to most economic invasive species assessments that estimate damages only after substantial and often irreversible injury has occurred to native ecosystems (Rothlisberger et al. 2012; Lodge et al. 2016).

The sensitivity analysis showed that results are robust considering the assumptions. The mean damages could be underestimated for the following reasons. First, the study only included sockeye salmon, which since 1984 amounted to half of the wholesale value of Alaska salmon (ADFG 2016)¹⁴. Second, the analysis leaves out potential effects on other sectors such as recreational or subsistence fisheries. For example, in Bristol Bay other research found that recreational and subsistence fisheries can amount to more than twice the net economic value attributable to the commercial salmon fishery (Duffield et al. 2013). Third, the study does not quantify the effects of *Elodea* on other ecosystem services. For example, there is evidence that *Elodea* affects nutrient cycling (Ozimek, Donk, and Gulati 1993), reduces lakefront property values by up to 16% (Zhang and Boyle 2010), and has severe impacts on biodiversity (Mjelde et al. 2012). In addition, *Elodea* invasions of remote waterbodies can also affect floatplane access and lead to recreation losses. Some of these limitations also underline that the true value of ecosystem services that are affected by *Elodea* are likely higher and cannot solely be expressed in monetary units. However, damages would likely be smaller if the model would account for the varying spatial dispersal across Alaska by for example the floatplane pathway. Additionally, a

¹⁴ Sockeye salmon amount to 26% of Alaska's commercial salmon catch in volume.

more detailed model would also account for the probability that isolated elodea populations collapse naturally.

Fourth, potential damages to salmon harvesters as expressed by changes to the producer surplus are not accounted for in the framework. A production function approach, where habitat quality is a direct input into salmon production, could measure such welfare changes (D. J. Knowler et al. 2003). Fifth, the assumption of ordinary Marshallian demand prevents a more detailed analysis of how the underlying individual consumer preferences could change given that the invasion of *Elodea* changes the quality of fish or consumers' perceived changes to environmental quality. For example, consumers may be hesitant to buy wild Alaska salmon knowing that the species' existence is threatened by aquatic invasive species. In instances where the quality of the ecosystem service is of concern, measures of compensating or equivalent surplus would be more appropriate. Last, the estimated damages do not include the potential cost of managing *Elodea* and, thus, do not provide a full accounting of the social costs of a potential invasion.

Damage assessments are usually based on empirical evidence of economic and ecological changes after invasions while controlling for different drivers of ecosystem and human system conditions. Our early-stage assessment of potential damage lacks empirical data on changes resulting from the invasion. Obtaining expert knowledge provided a feasible solution to data limitation, while being able to explicitly quantify uncertainty in the estimates. Some simplifications were needed. The use of the logistic growth model ignores specific age-structure effects of *Elodea* on salmon and also ignores correlation between growth rates and carrying capacity. Low levels of dissolved oxygen associated with crashing *Elodea* populations are a concern for freshwater life stages (Schwoerer, Little, and Hayward 2018) and encroachment of *Elodea* in salmon spawning beds is a concern for spawning adults in other locations outside Alaska where *Elodea* occurs in its natural range (Merz et al. 2008). These density-dependent effects would likely have higher impacts on sockeye populations spawning in slow moving streams and shallow water depth that is more suitable to *Elodea* compared to lake spawners in deeper waters (Braun and Reynolds 2014; Dodds and Biggs 2002).

Detailed age-structure data would allow analysis of fisheries management actions under an invasion scenario. The application of the logistic growth model instead focused on explaining the effects of *Elodea* on an entire population of salmon irrespective of age classes and is not suitable to provide fisheries management advice (Larkin 1977). Recognizing that expert elicitation is no panacea for biophysical research that establishes the ecological relationship between the invader and the harvestable resource, expert elicitation does however, enable researchers to quantify a first damage estimate from which further research can be expanded.

Not surprisingly, the expert-derived growth rates for sockeye salmon contribute the most to variance in the damage estimate. This result suggests that investments into biophysical research on the effects of *Elodea* on salmon are warranted to improve the precision of damage estimates.

7. CONCLUSION

Even though the range of potential damages estimated are large, the mean annual damage of \$159 million suggests that investment in preventing aquatic invasive species from gaining a foothold in Alaska is justified. Considering the attention and investment the invasive species threat in the Great Lakes has received in the past decade, the much larger damage estimate presented here raises the question as to whether these investments are optimally allocated. The results presented here suggest that future invasive species investments are better directed towards preventing damage to some of the most productive and intact ecosystems of national and global significance, especially as still in their pristine state are rare (Pinsky et al. 2009). With the invasive species problem still in its infancy in Alaska, society still has the opportunity to take prevention seriously

Directions for future research should be aimed at first reducing uncertainty associated with damage estimate by conducting biophysical research into *Elodea*'s growth effects on sockeye and possible other salmonids and second accounting for *Elodea*'s landscape-wide distribution pathways. The sensitivity analysis of this study clearly showed that the believed distribution of annual average growth rates for sockeye salmon in *Elodea*-infested salmon habitat is the primary driver of uncertainty in the damage estimate. Since the study does not account for the landscape-wide distribution of *Elodea* into salmon-bearing watersheds, accounting for the primary human pathways of long-distance dispersal through floatplane and boat traffic could further refine and potentially reduce damage estimates. Similarly, accounting for the natural rate of *Elodea* collapse would have the similar effects.

REFERENCES

- ADFG. 2016. "Commercial Operator's Annual Reports (COAR)." Anchorage, AK: Alaska Department of Fish and Game. 2016. <http://www.adfg.alaska.gov>.
- . 2018. "Alaska Commercial Salmon Harvest and Exvessel Value." Anchorage, AK: Alaska Department of Fish and Game. 2018. <http://www.adfg.alaska.gov>.
- Aitchison, J, and JAC Brown. 1976. *The Lognormal Distribution with Special Reference to Its Uses in Economics*. Cambridge, UK: Cambridge University Press.
- Andersen, Trude B., Kristin H. Roll, and Sigbjorn Tveteras. 2008. "The Price Responsiveness of Salmon Supply in the Short and Long Run." *Marine Resource Economics* 23: 425–37.
- Arrow, KJ, Maureen L. Cropper, Christian Gollier, Ben Groom, Geoffrey M. Heal, Richard G. Newell, William D. Nordhaus, et al. 2013. "How Should Benefits and Costs Be Discounted in an Intergenerational Context?" Washington, DC. <https://pdfs.semanticscholar.org/fc63/f9c324376ba68d866d6f6fb6725e44c604a6.pdf>.
- Asche, Frank, Misund Bard, and Oglend Atle. 2019. "The Case and Cause of Salmon Price Volatility." *Marine Resource Economics* 34 (1): 23–38.
- Barbier, Edward B., Mike Acreman, and Duncan Knowler. 1997. "Economic Valuation of Wetlands." Gland, Switzerland.
- Baumgärtner, Stefan, Alexandra M Klein, Denise Thiel, and Klara Winkler. 2015. "Ramsey Discounting of Ecosystem Services." *Environmental and Resource Economics* 61: 273–96. <https://doi.org/10.1007/s10640-014-9792-x>.
- Beverton, RJH, and SJ Holt. 1957. "On the Dynamics of Exploited Fish Populations." *U.K. Ministry of Agriculture and Fisheries, Fisheries Investigations (Series 2)* 19: 1–533.
- Bowmer, Kathleen H., S.W.L. Jacobs, and G.R. Sainty. 1995. "Identification, Biology and Management of *Elodea Canadensis*, Hydrocharitaceae." *Journal of Aquatic Plant Management* 33: 13–19.
- Braun, Douglas C, and John D Reynolds. 2014. "Life History and Environmental Influences on Population Dynamics in Sockeye Salmon." *Canadian Journal of Fisheries and Aquatic Sciences* 71: 1198–1208.
- Carey, Michael, Suresh A. Sethi, Sabrina Larsen, and Cecil Rich. 2016. "A Primer on Potential Impacts, Management Priorities, and Future Directions for *Elodea* Spp. in High Latitude Systems: Learning from the Alaska Experience." *Hydrobiologia* 777 (1): 1–19. <https://doi.org/DOI:10.1007/s10750-016-2767-x>.
- Champion, Paul D, Mary D De Winton, and John S Clayton. 2014. "A Risk Assessment Based Proactive Management Strategy for Aquatic Weeds in New Zealand." *Management of Biological Invasions* 5 (3). http://www.reabic.net/journals/mbi/2014/3/MBI_2014_Champion_etal_correctedproof.pdf.
- Clark, John H, Andrew McGregor, Robert D Mecum, Paul Krasnowski, and Amy M Carroll. 2006. "The Commercial Salmon Fishery in Alaska." *Alaska Fishery Research Bulletin* 12 (1): 1–146.

- Clemen, Robert T. 2008. "Comment on Cooke's Classical Method." *Reliability Engineering & System Safety* 93: 760–65. <https://doi.org/10.1016/j.ress.2008.02.003>.
- Cook, Christopher D.K., and Katharina Urmi-König. 1985. "A Revision of the Genus *Elodea* (Hydrocharitaceae)." *Aquatic Botany* 21: 111–56.
- Cooke, Roger M. 1991. *Experts in Uncertainty*. New York: Oxford University Press.
- DeVoretz, D. 1982. "An Econometric Demand Model for Canadian Salmon." *Canadian Journal of Agricultural Economics* 30 (1): 49–60.
- Dodds, Walter K., and Barry J.F. Biggs. 2002. "Water Velocity Attenuation by Stream Periphyton and Macrophytes in Relation to Growth Form and Architecture." *Journal of the North American Benthological Society* 21 (1): 2–15.
- Doelle, Meinhard. 2003. "The Quiet Invasion : Legal and Policy Responses to Aquatic Invasive Species in North America." *International Journal of Marine and Coastal Law* 18 (2): 261–95.
- Duffield, John, Chris Neher, David Patterson, Gunnar Knapp, Tobias Schwörer, Ginny Fay, and Oliver Scott Goldsmith. 2013. "Bristol Bay Wild Salmon Ecosystem: Baseline Levels of Economic Activity and Values." *An Assessment of Potential Mining Impacts on Salmon Ecosystems of Bristol Bay, Alaska*. U.S. Environmental Protection Agency. <http://cfpub.epa.gov/ncea/bristolbay/recordisplay.cfm?deid=253500#Download>.
- Erhard, Daniela, Georg Pohnert, and Elisabeth M. Gross. 2007. "Chemical Defense in *Elodea Nuttallii* Reduces Feeding and Growth of Aquatic Herbivorous Lepidoptera." *Journal of Chemical Ecology* 33 (8): 1646–61. <https://doi.org/10.1007/s10886-007-9307-0>.
- Finnoff, David, Jason F. Shogren, Brian Leung, and David Lodge. 2005. "The Importance of Bioeconomic Feedback in Invasive Species Management." *Ecological Economics* 52 (3): 367–81. <https://doi.org/10.1016/j.ecolecon.2004.06.020>.
- Fischer, K, D Lewandowski, and M Janssen. 2013. "Estimating Unknown Parameters in *Haemophilia* Using Expert Judgement Elicitation." *Haemophilia* 19 (5): 282–88.
- Freeman, A Myrick. 2003. *The Measurement of Environmental and Resource Values*. 2nd ed. Washington, D.C.: Resources for the Future.
- Frésard, Marjolaine, and Jean Boncoeur. 2006. "Controlling the Biological Invasion of a Commercial Fishery by Space Competitor: A Bioeconomic Model with Reference to the Bay of St-Brieuc Scallop Fishery." *Agricultural and Resource Economics Review* 35 (1): 78–97.
- Friedman, Sam. 2015. "Invasive *Elodea* Found North of Nenana; First Interior Sighting beyond Fairbanks Area." *Fairbanks Daily News Miner*, September 1, 2015. http://www.newsminer.com/news/local_news/invasive-elodea-found-north-of-nenana-first-interior-sighting-beyond/article_cd427cf0-511d-11e5-8c60-c733aa6798ec.html.
- Gleditsch, Jason M, and Tomas A. Carlo. 2011. "Frugivores in Central Pennsylvania." *Diversity and Distributions* 17: 244–53. <https://doi.org/10.1111/j.1472-4642.2010.00733.x>.
- Grigore, Bogdan, Jaime Peters, Christopher Hyde, and Ken Stein. 2016. "A Comparison of Two Methods for Expert Elicitation in Health Technology Assessments." *BMC Medical*

- Research Methodology* 16 (85): 1–11. <https://doi.org/10.1186/s12874-016-0186-3>.
- Haddon, Malcolm. 2011. *Modelling and Quantitative Methods in Fisheries*. 2nd ed. Boca Raton, FL: Chapman & Hall/CRC.
- Hammit, James K, and Yifan Zhang. 2013. “Combining Experts’ Judgments: Comparison of Algorithmic Methods Using Synthetic Data.” *Risk Analysis* 33 (1): 109–20. <https://doi.org/10.1111/j.1539-6924.2012.01833.x>.
- Hare, S.R., N.J. Mantua, and R.C. Francis. 1999. “Inverse Production Regimes: Alaska and West Coast Pacific Salmon.” *Fisheries* 24: 6–14.
- Heikkinen, Risto, Niko Leikola, Stefan Fronzek, Raino Lampinen, and Heikki Toivonen. 2009. “Predicting Distribution Patterns and Recent Northward Range Shift of an Invasive Aquatic Plant: *Elodea Canadensis* in Europe.” *BioRisk* 2 (November): 1–32. <https://doi.org/10.3897/biorisk.2.4>.
- Hilborn, Ray, and Marc Mangel. 1997. *The Ecological Detective*. Princeton, NJ: Princeton University Press.
- Hollander, Zaz. 2014. “Lake-Choking Invasive Weed Makes It to Mat-Su.” *Alaska Dispatch News*, October 16, 2014. <http://www.adn.com/article/20141016/lake-choking-invasive-weed-makes-it-mat-su>.
- . 2015. “Pilots Push for Weed Killer to Combat Lake Hood’s Plane-Fouling Plants.” *Alaska Dispatch News*, July 10, 2015. <http://www.adn.com/article/20150710/pilots-push-weed-killer-combat-lake-hood-s-plane-fouling-plants>.
- Hussner, A, I Stiers, M J J M Verhofstad, E S Bakker, B M C Grutters, J Haury, G van Valkenburg, J L CH Brundu, et al. 2017. “Management and Control Methods of Invasive Alien Freshwater Aquatic Plants: A Review.” *Aquatic Botany* 136: 112–37. <https://doi.org/10.1016/j.aquabot.2016.08.002>.
- Iman, Ronald L., and WJ Conover. 1982. “A Distribution-Free Approach to Inducing Rank Correlation among Input Variables.” *Communications in Statistics - Simulation and Computation* 11 (3): 311–34.
- Johnston, Richard S., and W. Robert Wood. 1974. “Demand Analysis for Canned Red (Sockeye) Salmon at Wholesale.” Corvallis, OR. <http://ir.library.oregonstate.edu/xmlui/bitstream/handle/1957/4522/?sequence=1>.
- Josefsson, Melanie. 2011. “Invasive Alien Species Fact Sheet *Elodea Canadensis*, *Elodea Nuttallii* and *Elodea Callitrichoides*.” Stockholm, Sweden: Department of Natural Resources, Swedish Environmental Protection Agency. <http://www.nobanis.org/files/factsheets/Elodea.pdf>.
- Kamarianos, A., J. Altiparmakis, X. Karamanlis, D. Kufidis, T. Kousouris, G. Fotis, and S. Kilikidis. 1989. “Experimental Evaluation of Fluridone Effectiveness on Fish Productive Aquatic Ecosystems.” *Journal of Aquatic Plant Management* 27: 24–26.
- Knapp, Gunnar, Mouhcine Guettabi, and Oliver Scott Goldsmith. 2013. “The Economic Importance of the Bristol Bay Salmon Industry.” Anchorage, AK.
- Knapp, Gunnar, and James J Murphy. 2010. “Voluntary Approaches to Transitioning from

- Competitive Fisheries to Rights- Based Management : Bringing the Field into the Lab.” *Agricultural and Resource Economics Review* 39 (2): 245–61.
- Knapp, Gunnar, Cathy A. Roheim, and James L. Anderson. 2007. *The Great Salmon Run: Competition between Wild and Farmed Salmon*. Washington, D.C.: TRAFFIC North America World Wildlife Fund.
- Knowler, D. 2005. “Reassessing the Costs of Biological Invasion: *Mnemiopsis leidyi* in the Black Sea.” *Ecological Economics* 52 (2): 187–99. <https://doi.org/10.1016/j.ecolecon.2004.06.013>.
- Knowler, Duncan J., Brice W. MacGregor, Michael J. Bradford, and Randall M. Peterman. 2003. “Valuing Freshwater Salmon Habitat on the West Coast of Canada.” *Journal of Environmental Management* 69 (3): 261–73. <https://doi.org/10.1016/j.jenvman.2003.09.001>.
- Larkin, PA. 1977. “An Epitaph for the Concept of Maximum Sustainable Yield.” *Transactions of the American Fisheries Society* 106: 1–11.
- Les, Donald H, and Nicholas P Tippery. 2013. “In Time and with Water . . . the Systematics of Alismatid Monocotyledons.” In *Early Events in Monocot Evolution*, edited by Paul Wilkin and Simon J. Mayo, 118–64. Cambridge, UK: Cambridge University Press.
- Lodge, David M, Paul W Simonin, Stanley W Burgiel, Reuben P Keller, Jonathan M Bossenbroek, Christopher L Jerde, Andrew M Kramer, et al. 2016. “Risk Analysis and Bioeconomics of Invasive Species to Inform Policy and Management.” *Annual Review of Environment and Resources* 41: 453–88. <https://doi.org/10.1146/annurev-environ-110615-085532>.
- Maestas, C. D., M. K. Buttice, and W. J. Stone. 2014. “Extracting Wisdom from Experts and Small Crowds: Strategies for Improving Informant-Based Measures of Political Concepts.” *Political Analysis* 22 (3): 354–73. <https://doi.org/10.1093/pan/mpt050>.
- McDowell Group. 2015. “Sockeye Market Analysis.” Juneau, AK. <http://www.mcdowellgroup.net/wp-content/uploads/2015/12/Fall-2015-Sockeye-Market-Report-Final.pdf>.
- . 2017. “The Economic Value of Alaska’s Seafood Industry.” Juneau, AK. <http://www.mcdowellgroup.net/wp-content/uploads/2017/10/ak-seafood-impacts-sep2017-final-digital-copy.pdf>.
- McElhany, P., Mary H. Ruckelshaus, Michael J. Ford, Thomas C. Wainwright, and Eric Bjorkstedt. 2000. “Viable Salmonid Populations and the Recovery of Evolutionarily Significant Units.” *NOAA Technical Memorandum NMFS-NWFSC*; 42., no. June: 156. http://www.nwfsc.noaa.gov/assets/25/6190_06162004_143739_tm42.pdf.
- McFadden, Daniel. 1973. “Conditional Logit Analysis of Qualitative Choice Behavior.” In *Frontiers in Econometrics*, edited by P. Zarembka, 105–42. New York: Academic Press. <https://doi.org/10.1108/eb028592>.
- Merz, Joseph E., James R. Smith, Michelle L. Workman, Jose D. Setka, and Bert Mulchaey. 2008. “Aquatic Macrophyte Encroachment in Chinook Salmon Spawning Beds: Lessons Learned from Gravel Enhancement Monitoring in the Lower Mokolumne River,

- California.” *North American Journal of Fisheries Management* 28 (5): 1568–77.
<https://doi.org/10.1577/M07-043.1>.
- Mjelde, Marit, Paola Lombardo, Dag Berge, and Stein W. Johansen. 2012. “Mass Invasion of Non-Native *Elodea Canadensis* Michx. in a Large, Clear-Water, Species-Rich Norwegian Lake – Impact on Macrophyte Biodiversity.” *Annales de Limnologie - International Journal of Limnology* 48 (2): 225–40. <https://doi.org/10.1051/limn/2012016>.
- Morgan, M. Granger. 2011. “Certainty, Uncertainty, and Climate Change.” *Climatic Change* 108 (4): 707–21. <https://doi.org/10.1007/s10584-011-0184-8>.
- Morgan, M. Granger, Samuel C. Morris, Max Henrion, Deborah A. L. Amaral, and William R. Rish. 1984. “Technical Uncertainty in Quantitative Policy Analysis - A Sulfur Air Pollution Example.” *Risk Analysis* 4 (3): 201–16. <https://doi.org/10.1111/j.1539-6924.1984.tb00139.x>.
- Morgan, M Granger. 2014. “Use (and Abuse) of Expert Elicitation in Support of Decision Making for Public Policy.” *Proceedings of the National Academy of Sciences of the United States of America* 111 (20): 7176–84. <https://doi.org/10.1073/pnas.1319946111>.
- Morton, John. 2016. “Refuge Notebook : *Elodea* Gone from the Kenai Peninsula ?” *Peninsula Clarion*, June 2, 2016.
- MSC. 2017. “Alaska Salmon Fishery - Certified Sustainable Seafood.” Seattle, WA: Marine Stewardship Council (MSC). <https://fisheries.msc.org/en/fisheries/alaska-salmon/@@certificates>.
- NBN. 2015. “NBN Gateway.” Nottingham, UK: National Biodiversity Network Great Britain. <https://data.nbn.org.uk/Taxa>.
- O’Hagan, Anthony, Caitlin E. Buck, Alireza Daneshkhah, J. Richard Eiser, Paul H. Garthwaite, David J. Jenkinson, Jeremy E. Oakley, and Tim Rakow. 2006. *Uncertain Judgments: Eliciting Experts’ Probabilities*. West Sussex, UK: Wiley.
- OMB. 2016. “2017 Discount Rates for OMB Circular No. A-94.” Washington, DC: U.S. Executive Office of the President Office of Management and Budget. <https://www.whitehouse.gov/sites/default/files/omb/memoranda/2017/m-17-10.pdf>.
- Ozimek, Teresa, Ellen Van Donk, and Ramesh D Gulati. 1993. “Growth and Nutrient Uptake by Two Species of *Elodea* in Experimental Conditions and Their Role in Nutrient Accumulation in a Macrophyte- Dominated Lake.” *Hydrobiologia* 251 (1–3): 13–18.
- Palisade. 2016. “How @Risk Correlates Inputs.” 2016. <http://kb.palisade.com/index.php?pg=kb.page&id=71>.
- Palisade Corporation. 2016. “@RISK: Risk Analysis and Simulation Add-in for Microsoft® Excel.” Ithaca, NY: Palisade Corporation.
- Perrings, Charles, Mark Williamson, Edward B Barbier, Doriana Delfino, Silvana Dalmazzone, Peter Simmons, and Andrew Watkinson. 2002. “Biological Invasion Risks and the Public Good : An Economic Perspective.” *Conservation Ecology* 6 (1): 7.
- Peterman, Randall M., and Judith L. Anderson. 1999. “Decision Analysis: A Method for Taking Uncertainties into Account in Risk-Based Decision Making.” *Human and Ecological Risk*

Assessment: An International Journal 5 (2): 231–44.
<https://doi.org/10.1080/10807039991289383>.

- Peterson, Douglas P, Bruce E Rieman, Jason B Dunham, Kurt D Fausch, and Michael K Young. 2008. “Analysis of Trade-Offs between Threats of Invasion by Nonnative Brook Trout (*Salvelinus Fontinalis*) and Intentional Isolation for Native Westslope Cutthroat Trout (*Oncorhynchus Clarkii Lewisi*).” *Canadian Journal of Fisheries and Aquatic Sciences* 65 (4): 557–73. <https://doi.org/10.1139/f07-184>.
- Pinsky, Malin L, Dane B Springmeyer, Matthew N Goslin, and Xanthippe Augerot. 2009. “Range-Wide Selection of Catchments for Pacific Salmon Conservation.” *Conservation Biology* 23 (3): 680–91. <https://doi.org/10.1111/j.1523-1739.2008.01156.x>.
- Redekop, Petra, Deborah Hofstra, and Andreas Hussner. 2016. “Elodea Canadensis Shows a Higher Dispersal Capacity via Fragmentation than Egeria Densa and Lagarosiphon Major.” *Aquatic Botany* 130: 45–49. <https://doi.org/10.1016/j.aquabot.2016.01.004>.
- Ricker, W.E. 1975. “Computation and Interpretation of Biological Statistics of Fish Populations.” 191. Vol. 108. *Bulletin*. Ottawa. <https://doi.org/10.1038/108070b0>.
- Riis, Tenna, and Barry J. F. Biggs. 2003. “Hydrologic and Hydraulic Control of Macrophyte Establishment and Performance in Streams.” *Limnology and Oceanography* 48 (4): 1488–97. <https://doi.org/10.4319/lo.2003.48.4.1488>.
- Rørslett, Bjørn, D A G Berge, and Steine W. Johansen. 1986. “Lake Enrichment by Submersed Macrophytes: A Norwegian Whole-Lake Experience with Elodea Canadensis.” *Aquatic Botany* 26: 325–40.
- Rosaen, Alex L, Erin Grover, and Colby W. Spencer. 2012. “The Costs of Aquatic Invasive Species to Great Lakes States.”
- Rothlisberger, John D, David C Finnoff, Roger M Cooke, and David M Lodge. 2012. “Ship-Borne Nonindigenous Species Diminish Great Lakes Ecosystem Services.” *Ecosystems* 15: 462–76. <https://doi.org/10.1007/s10021-012-9522-6>.
- Schindler, Daniel E, Ray Hilborn, Brandon Chasco, Christopher P Boatright, Thomas P Quinn, Lauren a Rogers, and Michael S Webster. 2010. “Population Diversity and the Portfolio Effect in an Exploited Species.” *Nature* 465 (June): 609–12. <https://doi.org/10.1038/nature09060>.
- Schneider, James C. 2000. “Evaluation of the Effects of the Herbicide Sonar on Sport Fish Populations in Michigan Lakes.” 2000–2. Michigan Department of Natural Resources Fisheries Technical Report. Ann Arbor, MI. <http://www.aswcd.org/MDEQstudy-Sonar-on-fish-population.pdf>.
- Schultz, Rachel, and Eric Dibble. 2012. “Effects of Invasive Macrophytes on Freshwater Fish and Macroinvertebrate Communities: The Role of Invasive Plant Traits.” *Hydrobiologia* 684 (1): 1–14. <https://doi.org/10.1007/s10750-011-0978-8>.
- Schwoerer, Tobias, Joseph M. Little, and Greg Hayward. 2018. “Quantifying Expert Opinion Using a Discrete Choice Model: Will Invasive Elodea Spp. Threaten Wild Salmonids in Alaska?” <https://doi.org/10.13140/RG.2.2.29993.44646>.
- Schwörer, Tobias, Rebekka N Federer, and Howard J Ferren. 2014. “Invasive Species

- Management Programs in Alaska : A Survey of Statewide Expenditures , 2007 – 11.” *Arctic* 67 (1): 20–27. <https://doi.org/http://dx.doi.org/10.14430/arctic4359>.
- Sethi, Suresh Andrew, Matthew Reimer, and Gunnar Knapp. 2014. “Alaskan Fishing Community Revenues and the Stabilizing Role of Fishing Portfolios.” *Marine Policy* 48: 134–41. <https://doi.org/10.1016/j.marpol.2014.03.027>.
- Shogren, Jason F. 2000. “Risk Reduction Strategies against the ‘Explosive Invader.’” In *The Economics of Biological Invasions*, edited by Charles Perrings, Mark Williamson, and Silvana Dalmazzone, 56–69. Northampton, MA: Edward Elgar.
- Short, C.I. I, S K Gross, and D Wilkinson. 2004. “Preventing, Controlling, and Managing Alien Species Introduction for the Health of Aquatic and Marine Ecosystems.” In *Sustainable Management of North American Fisheries*, 43:109–25. Bethesda, MD: American Fisheries Society.
- Simberloff, Daniel, and Leah Gibbons. 2004. “Now You See Them, Now You Don’t! – Population Crashes of Established Introduced Species.” *Biological Invasions* 6 (2): 161–72. <https://doi.org/10.1023/B:BINV.0000022133.49752.46>.
- Sinnott, Rick. 2013. “With a Noxious Aquatic Weed Threatening Alaska’s Prime Fishing Waters, Science and Local Politics Are at Odds.” *Alaska Dispatch*, November 28, 2013. <http://www.adn.com/article/20131128/noxious-aquatic-weed-threatening-alaskas-prime-fishing-waters-science-and-local>.
- Soares, MO, L Bojke, J Dumville, C Iglesias, N Cullum, and K Claxton. 2011. “Methods to Elicit Experts’ Beliefs over Uncertain Quantities: Application to a Cost Effectiveness Transition Model of Negative Pressure Wound Therapy for Severe Pressure Ulceration.” *Statistics in Medicine* 30 (19): 2363–80.
- Speirs-Bridge, Andrew, Fiona Fidler, Marissa McBride, Louisa Flander, Geoff Cumming, and Mark Burgman. 2010. “Reducing Overconfidence in the Interval Judgments of Experts.” *Risk Analysis* 30 (3): 512–23. <https://doi.org/10.1111/j.1539-6924.2009.01337.x>.
- Spicer, K.W., and Paul M. Catling. 1988. “The Biology of Canadian Weeds. 88.” *Canadian Journal of Plant Science* 68 (4): 1035–51.
- Strecker, Angela L., Philip M. Campbell, and Julian D. Olden. 2011. “The Aquarium Trade as an Invasion Pathway in the Pacific Northwest.” *Fisheries* 36 (2): 74–85. <https://doi.org/10.1577/03632415.2011.10389070>.
- Swartz, A. Nelson. 1978. “The Pacific Coho Salmon Fishery: An Intraseasonal and Interregional Economic Analysis of the Ex-Vessel Market.” Oregon State University. <https://ir.library.oregonstate.edu/xmlui/bitstream/handle/1957/25951/SWARTZANELSON1979.pdf?sequence=1>.
- Teck, Sarah J, Benjamin S Halpern, Carrie V Kappel, Fiorenza Micheli, Kimberly A Selkoe, Caitlin M Crain, Rebecca Martone, et al. 2010. “Using Expert Judgment to Estimate Marine Ecosystem Vulnerability in the California Current.” *Ecological Applications* 20 (5): 1402–16.
- USGS. 2017. “National Hydrography Dataset.” 2017. <http://nhd.usgs.gov/data.html>.
- Wang, Hsiung Der. 1976. “An Econometric Study of the Canadian Sockeye Salmon Market.”

Oregon State University.

Winston, Wayne L., and S. Christian Albright. 2016. *Practical Management Science*. 5th ed. Boston, MA: Cengage Learning.

Wittmann, Marion E, Roger M Cooke, John D Rothlisberger, and David M Lodge. 2014. “Using Structured Expert Judgment to Assess Invasive Species Prevention: Asian Carp and the Mississippi-Great Lakes Hydrologic Connection.” *Environmental Science & Technology* 48 (4): 2150–56. <https://doi.org/10.1021/es4043098>.

Zhang, Congwen, and Kevin J. Boyle. 2010. “The Effect of an Aquatic Invasive Species (Eurasian Watermilfoil) on Lakefront Property Values.” *Ecological Economics* 70 (December): 394–404. <https://doi.org/10.1016/j.ecolecon.2010.09.011>.

U.S. State Invasive Species Councils

Prepared by the Government Relations Committee, Alaska Invasive Species Partnership

December 2024

State Invasive Species Councils at a Glance

State	Name ¹	Mech.	Year	Website	Funded	# Seats	Administrative Department
Arizona	Advisory Council	E.O.	2005	N	UNK	21	Game and Fish, Agriculture
California	Council	Law	2019	link	True	6	Food and Agriculture
Connecticut	Plants Council	Law	2003	link	False	9	Energy and Environmental Protection
Guam	Council	Law	2011	N	True	9	Agriculture
Hawai'i	Council	Law	2003	link	True	6	Lands and Natural Resources
Idaho	Council	Law	2017	link	True	12	Agriculture
Indiana	Council	E.O.	2009	link	UNK	11	Purdue University College of Agriculture
Massachusetts	Advisory Council		Bill		True	14	Agricultural Resources
Montana	Council	Law	2017	link	True	22	Natural Resources and Conservation
Nebraska	Council	Law	2012	link	UNK	12	Game and Parks Commission
New Hampshire	Committee	Law	2000	N	False	11	Agriculture, Markets, and Food
New Jersey	Task Force		Bill		False	9	Agriculture
New York	Council	Law	2007	link	True	9	Environmental Conservation
Oregon	Council	Law	2009	link	True	22	Agriculture
Texas	Coordinating Committee	Law	2009	link	False	6	State Soil & Water Conservation Board
Virginia	Advisory Group	Law	2023	N	False	15	Conservation and Recreation
Washington	Council	Law	2006	link	True	8	Recreation and Conservation
Wisconsin	Council	Law	2017	link	False	12	Natural Resources

¹ All names preceded by "Invasive Species" except Connecticut, which is the "Invasive Plants Council"

Council Composition

St	State Representatives	Other Representatives	Notes
AZ	<p>Game & Fish Agriculture Transportation Water Resources State Land State Parks Health Services Office of the Governor</p>	<p>University of Arizona Native American community Agricultural interests Conservation community Forest health management interests Ranching industry Nursery & landscape industry Pet industry Cooperative Weed Management Areas or similar Federal agencies Academic community / University Cooperative Extension</p>	
CA	<p>Food & Agriculture Natural Resources Environmental Protection Transportation Health & Human Services Emergency Services</p>		<p>One dedicated staff member serves as executive director</p> <p>Law also creates 19-member Invasive Species Advisory Committee</p>
CT	<p>Agriculture Energy & Environmental Protection CT Agricultural Experiment Station</p>	<p>UC College of Agriculture & Natural Resources Invasive Plant Atlas of New England Nonprofit environment association Nonprofit concerned with growers & retailers of plants Nonprofit associated concerned with oceans, lakes, & rivers Company that grows or sells plants</p>	<p>Members outside state agencies & the University of CT are appointed by congressional leaders (Senate minority leader, House speaker, Senate president pro tempore, House minority leader) & Governor</p>
GU	<p>Agriculture Customs & Quarantine International Airport Authority Port Authority Parks & Recreation Northern SWCD Southern SWCD Bureau of Statistics & Plans (non-voting) Homeland Security (non-voting)</p>	<p>University of Guam Mayors Council of Guam USDA (non-voting)</p>	<p><u>Staff</u>. Territorial Invasive Species Coordinator (funded through Inspection Fee Fund, see next section)</p>
HI	<p>Lands & Natural Resources Agriculture Health Business, Economic Development, & Tourism Transportation</p>	<p>University of Hawaii</p>	<p>There are 8 non-voting legislative members—4 senators & 4 representatives from each of the four counties</p>

St	State Representatives	Other Representatives	Notes
ID	Governor's office Agriculture Environmental Quality Parks & Recreation Fish & Game Lands Water Resources Transportation Species Conservation State Senate State House	Idaho Outfitters & Guides Association <i>In addition: "Representatives & members of federal entities, local government organizations, tribal governments, Idaho universities & private & not-for-profit organizations with an interest in the well-being of Idaho pertaining to invasive species shall be invited to participate by the director of the Department of Agriculture."</i>	
IN	Agriculture Transportation Division of Fish & Wildlife Division of Entomology of Plant Pathology State veterinarian	Purdue University College of Agriculture individual representing invasive species research two individuals that represent any of the following industries (a) hardwood tree; (b) horticulture; (c) agriculture; (d) aquaculture Two individuals primarily concerned with any of the following (a) land trusts; (b) biodiversity conservation; (c) aquatic conservation; (d) local parks & recreation	
MA	Natural Heritage & Endangered Species Program Division of Ecological Restoration Agricultural Resources	Conservation organization River association MA Invasive Plant Advisory Group Land trust Native plant organization Association of conservation commissions Plant nursery association Landscape association MA Congress of Lake & Ponds Associations MA biodiversity research institute or university Veterinarian or biologist studying wildlife & plant diseases	Note this is an advisory committee Law creates new position of statewide invasive species coordinator in Ag. Res. chairs committee
MT	Fish, Wildlife, & Parks Natural Resources & Conservation Transportation Agriculture Commerce	County weed districts Conservation districts MT State University Extension Service Agriculture Conservation organizations Wildlife organizations Fishing organizations Hydropower utility industry Private landowners Each of the tribal governments in MT (8)	States that council shall seek "active input & participation" from Army Corps of Engineers, Bureau of Reclamation, BLM, APHIS, FWS, USFS, & NPS Council also has Scientific Advisory Panel for specific topics

St	State Representatives	Other Representatives	Notes
NE	Agriculture Game & Parks Commission	Electric generating utility NE Forest Service, University of NE Institute of Ag. & Nat. Res. University of Nebraska-Lincoln BE Weed Control Association NE Association of Resources Districts Up to 5 at-large members representing public interests (at least 3 represent agricultural landowner interests)	<u>Nonvoting, Ex officio Representatives.</u> US DOI USDA APHIS USDA NRCS USGS The Nature Conservancy, Nebraska Field Office
NH	State entomologist Environmental Services Natural & Cultural Resources Transportation Fish & Game	University of NH College of Life Sciences & Agriculture University of NH cooperative extension Person representing horticultural interests Person representing environmental interests Person representing livestock owners' & growers' interests Person representing general public	
NJ	Agriculture Environmental Protection Health State Forester	Two public members appointed by Governor One public member appointed by Senate President One public member appointed by Assembly Speaker NJ Agricultural Experiment Station at Rutgers	Department of Environmental Protection shall also provide staff support to the task force
NY	Environmental Conservation Agriculture & Markets Transportation Office of Parks, Recreation & Historic Preservation Education State State Thruway Authority Adirondack Park Agency		Law also creates NY invasive species advisory council up to 25 at-large members
OR	Agriculture Fish & Wildlife State Forester Environmental Quality Marine Director Parks & Recreation State Invasive Species Coordinator <i>nonvoting</i> Governor's Office <i>nonvoting</i> Senate member <i>nonvoting</i> House member <i>nonvoting</i>	Center for Lakes & Reservoirs, Portland State University Sea Grant Environmental stewardship advocacy group Private industry advocacy group Native American or Indian tribe or association of tribes 6 members of entities related to invasive species, each located in one of six groups of counties Member representing the public	The wording for members representing groups of counties: "A member who represents an entity, regardless of the form of the entity, with a headquarters or principal operations in x, y, z... Counties & with a purpose of responding to invasive species concerns

St	State Representatives	Other Representatives	Notes
TX	Agriculture Parks & Wildlife State Soil & Water Conservation Board TX Forest Service Water Development Board	Texas AgriLife Extensive Service (Texas A&M)	law requiring one full time committee coordinator repealed in 2023 Other state agency representatives may be added by unanimous committee vote
VA	Agriculture & Consumer Services Conservation & Recreation Environmental Quality Forestry Health Transportation Wildlife Resources Marine Resources Commission Secretary of Natural & Historic Resources	Virginia Cooperative Extension VA Institute of Marine Science Representative for agriculture & forestry Industries Representative for conservation community Representative for academic institutions Representative for commercial interests	
WA	Agriculture Fish & Wildlife Ecology Natural Resources Transportation State Noxious Weed & Control Board	County east of the Cascade Mountains crest County west of the Cascade Mountains crest	Members may each invite one non-voting representative from USDA, USFWS, US EPA, & US Coast Guard to participate
WI	Natural Resources Administration Agriculture, Trade, & Consumer Protection Tourism Transportation	7 other members appointed by the governor representing public & private interests	

Council Funding

St	Funding	Budget	Notes
AZ	UNK		
CA	True	\$5,000,000 appropriated to Invasive Species Account (Budget Act of 2021)	Law establishes Invasive Species Account
CT	False		They state in annual reports from 2021-2023 that “The lack of funding for the coordinator position has negatively impacted the activities and public outreach of the Invasive Plants Council since 2014”
GU	True	Revenue: \$755,328 for FY 2025, \$654,864 for FY 2024 from Inspection Fee (see notes)	Law establishes Invasive Species Inspection Fee Fund appropriated to Dept of Agriculture for use by council (Fees are associated with freight inspection, quarantine, and eradication)
HI	True	Departmental appropriations request for FY 24 & 25: \$1,500,000 FY 2022 legislature appropriated \$5,750,000	Additional state funding for specific programs allocated to council (e.g., invasive ant eradication program)
ID	True	Appropriations from SB 1372 (2024 session): \$6,791,100 for Invasive Species Fund (see notes). This includes \$5,000,000 for detection of and response to quagga mussels.	The Idaho Invasive Species Law (2008) establishes Invasive Species Fund
IN	UNK	Cannot find balance of council fund or appropriations in recent budget bills	Law establishes invasive species council fund
MA	True	NA; Funding proposed in bill but not passed	Bill establishes Invasive Species Trust Fund
MT	True	\$537,445 appropriated for FY2024: \$208,865 administration; \$50,000 summit costs, travel, projects \$278,580 aquatic invasive species grants	
NE	UNK		In their 2022 report, they state that they worked to fill a council coordinator position with funds from Nebraska Games and Parks Commission. The position is housed at University of Nebraska Lincoln.
NH	False		Department of Ag. has a full-time invasive species coordinator. Coordinator’s involvement with council is unclear.
NJ	NA		Bill mentions that funds may be appropriated to task force
NY	UNK		There is a NY DEC Invasive Species Grant Program (2019: \$2.8 million. 2024: \$3 million). Unsure if connected with council. According to meeting minutes for the advisory council in 2022, DEC budget for invasive species was \$15,974,000. However, no line item for council specifically and could not find any in NYS budgets.

St	Funding	Budget	Notes
		\$168,858.00 from state lottery (thought to be annual)	
		\$1,470,000 one-time General Fund in 2024 HB 4148 (in Ways & Means upon adjournment):	
		- \$220,000 for the Invasive Species Council to conduct a program review.	Law establishes two funds: Invasive Species Council Account in the general fund and the Invasive Species Control Account outside the general fund. Law also states that Dept of Agriculture is responsible for ensuring payment of administrative expenses, though then can be reimbursed through Council Account.
OR	True	- \$500,000 for deposit into the Invasive Species Control Account to support outreach and education on invasive species.	
		- \$750,000 for deposit into the Invasive Species Control Account to fund activities to mitigate negative impacts from, and establishment of, invasive species.	Law also creates the Invasive Species Coordinator position
TX	False		
VA	False		
		\$851,000 for 2012-2023 (~71,000/year).	
WA	True	\$328,000 appropriated in 2023 senate bill 5187 (no appropriations in 2024 bill, though there is funding for aquatic invasive species program)	Section 79A.25.370 of law created Invasive Species Council fund, but this section expired in 2022, while remaining sections were renewed.
WI	False		While there is no funding for the council, the law describing council duties also enumerates appropriations to the Nat. Res. Dept for invasive species control

Council Reporting and Accomplishments

<i>Council Requirements</i>			
St	Reporting	Meetings	Notable Accomplishments
AZ	Develop prevention response plan Prepare invasive species management plan Prepare list of invasive species and update every 3 years Summit annual report to the governor and legislature	NA	AZ iMapInvasives project Invasive Species Management Plan (2008) Aquatic Invasive Species Management Plan (2011)
CA	Develop and maintain statewide surveys Host annual CA Invasive Species summit Develop and submit report of activities Develop statewide action plans	Annually	Strategic initiative for Invasive Shot Hole Borer Keeps Invasive Species List of CA updated and creates associated Impact Scorecards Pest Reporting Web App Organizes annual CA Invasive Species Action Week Rapid Respond to Emerging Invasive Species Threats funding grant
CT	Annually publish and update list of invasive or potentially invasive plants Develop and conduct education program Annual reports to General Assembly	2/Year	Produces annual reports (as mandated in Law) Oversees list of invasive and potentially invasive plants
GU	Develop invasive species management plan Annual progress reports	Quarterly	Meet monthly 2017-2019 management plan publicly available
HI	Create and implement prevention, early detection, rapid response, control, enforcement, and education plan Annual progress reports to governor and legislature Biennial multi-departmental budget proposal	2/Year	From 2023 annual report: Raised \$3,830,870 to support project and programs; initiated 68% of Biosecurity Plan actions; 610 new weed risk assessments; 10 interceptions of high-risk pests at ports of entry through monitoring program; 1230 public pest reports
ID	Annual report to governor and legislature	2/Year	2022-2026 strategic plan Watercraft inspections site improvements for relevant infrastructure (e.g., decontamination units) Watercraft Inspection Station dashboard and data analysis Annual interagency activity updates
IN	Convene/support invasive species meeting biennially Apply for and provide grants Submit a progress report every other year	Annually	Task force reports biennial reports

<i>Council Requirements</i>			
St	Reporting	Meetings	Notable Accomplishments
MA	Develop data relevant to invasive species Assist in implementation and revision of strategic management plan Annual report to legislature Create and maintain list of all present and potential invasive species Incorporate and expand existing MA Invasive Plant Advisory Group invasive plant assessment Review and establish fines and penalties for invasive species related crimes	Annually	NA – Proposed Bill
MT	Annual Reports to Governor and Department	2/Year	Holds biennial summit Developed MT Invasive Species Framework Issues Aquatic Invasive Species Grants Have many reports, newsletters, publications, etc. (seem very active)
NE	Develop invasive species adaptive management plan updated every three years Annual reports to Governor and Legislature	NA	Co-hosted 2023 annual North American Invasive Species Management Association (NAISMA) Other actions in their reports are accomplishments of agency programs. They seem to serve a supporting role in many of these programs
NH	With commissioner, prepare and publish by April 1 annually a list of invasive species deemed to present an immediate danger	NA	The invasive species list is kept up to date Contribution to 2023 New Hampshire Guide to Upland Invasive Species 2023 Invasive Species Strategic Plan
NJ	Develop plan to prevent and limit spread of invasive species Develop plan to restore ecosystems and repair damage caused by invasive species Prepare comprehensive invasive species management plan Prepare comprehensive list of all invasive species in NJ and classify based on threat level, updated every 3 years Annual written reports to the Governor and Legislature	Quarterly	NA – Proposed Bill

<i>Council Requirements</i>			
St	Reporting	Meetings	Notable Accomplishments
NY	development of a comprehensive plan for invasive species management organize and convene a biennial invasive species summit 2010 report to governor and legislature	Quarterly	Rapid Response Framework published in 2016 New York State Invasive Species Comprehensive Management Plan finalized 2018 Developed 2016-2021 Education and Outreach Strategic Plan Coordinates NY Invasive Species Awareness Week Held 2023 NY Invasive Species Expo, planning for 2025
OR	Biennial reports to Legislature	NA	The Invasive Species Threats & Opportunities: A Primer for Oregon Policymakers published Collaborated with students from Portland State University on invasive species projects Coordinated cross-agency Committees Maintained Oregon's Invasive Species Hotline Issues grants for Invasive Species Education & Outreach Hosted 2024 summit on invasive insect pests
TX	None	Annually	Meeting minutes published online and show updates from member's agencies
VA	Create invasive species plan and update every four years	2/Year	Invasive species management plan published in 2018
WA	Develop statewide strategic plan and update at least once every 3 years Biennial report to governor, senate, and house	NA	Website, reports, and meetings are current and active. In 2024, established 3 annual awards Hold regular events, including participation in National Invasive Species Awareness Week Hosts and maintains website, social media accounts, and reporting hub 2024 report outlines agency-level accomplishments. Council general accomplishments include "Improving Communication to Diverse Audiences, Understanding Tribal and Municipal Capabilities, Adjusting the Language of Invasive Species, Integrating Invasive Species Awareness into Other Plans"
WI	system for classifying invasive species procedure for awarding cost-sharing grants to control invasive species.	NA	Annual invasive species action awards While law does not specify meeting frequency, meeting minutes on website show multiple executive committee and public meetings for 2023 and 2024 Hold WI Invasive Species Awareness Day The Nat. Res. Department is required to submit biennial report (not the council, though the council seems to be heavily involved in reviewing the report)



LET'S KEEP ALASKA WILD AND FREE FROM INVASIVE SPECIES

Alaskans depend on their natural surroundings for food, work, and fun. The far-reaching effects of **INVASIVE SPECIES** threaten the very resources that are fundamental to life in Alaska.

WHO WE ARE

The mission of the Alaska Invasive Species Partnership (AKISP) is to prevent, detect, and manage invasive species in Alaska by facilitating and fostering invasive species awareness, knowledge and information sharing, and collaboration among diverse stakeholders.

WHAT WE DO

We strive to provide credible and unbiased scientific information to promote better management decisions. To share the latest information, research, management practices, and technologies related to invasive species concerns affecting Alaska, we host monthly statewide virtual meetings and an annual workshop for approximately 200 participants. We manage an active social media presence and regularly coordinate outreach campaigns.

WHAT'S THE PROBLEM?

Invasive species are not native to an area and cause harm to the environment, economy, and/or health. The cumulative effects of invasive species threaten Alaska's natural resources and economy. Previous annual economic costs of invasives exceeded **\$26 billion**¹ in the United States and **\$423 billion** globally². Like groceries and gas, the cost to prevent and manage invasive species is higher today than it was in years past. But, the cost of responding to these threats must not be a reason to turn a blind eye. **BIOSECURITY** should be given the same focus and attention as national security, cyber security, or healthcare: we must invest in prevention, detection, and response to protect what we value, including Alaska fisheries, agriculture, recreation, and tourism.

¹ Crystal-Ornelas et al. 2021. NeoBiota 67: 485-510.

² IPBES 2023. <https://www.ipbes.net/IASmediarelease>

WHAT YOU CAN DO

- Allocate funding and pass legislation to create an Alaska Invasive Species Council
- Enact legislation to facilitate coordination and collaboration among state agencies on invasive species prevention and management
- Contact the AKISP Board and your local invasive species experts to learn about species causing problems in your district and how you can help address management challenges
- Support and participate in community efforts to control invasive species
- Get involved by joining AKISP monthly meetings or attending our annual workshop. To join our listserv and receive our biannual newsletter visit <https://alaskainvasives.org/>



The AKISP held its annual workshop in Sitka, November 2023, with the theme "Invaders on the Edge". This event rotates location around the state between Anchorage, Fairbanks, and a remote community. Photo: Ashley Lutto, USFVWS

Thanks to collaboration among AKISP partners, Alaska has had some success managing invasive species; however, programs continue to be severely underfunded and are operating with limited staff.


CASE EXAMPLE: ELODEA IN THE MAT-SU BOROUGH

The aquatic invasive plant Elodea was discovered in Alexander Lake in 2014. Initially sparse, it rapidly spread, including to nearby Sucker Lake, thanks to its easy dispersal by boats and floatplanes.

Elodea's aggressive growth disrupts native plants, harms fish habitat, and blocks movement of boats and floatplanes. In 2019, agencies collaborated to restrict floatplane traffic in the infested lakes to prevent further spread and then secured grants to apply herbicide in both lakes.

Despite initial success, Elodea was subsequently found in Big Lake, which resulted in limited personnel and resources diverted to the new infestation in a high use lake. In the absence of continued treatment, a resurgence of Elodea in Alexander and Sucker lakes occurred.

Today, Elodea covers both lakes and is spreading downstream into Alexander Creek. This example highlights how limited resources delay action, hindering invasive species management. Yet, eradication remains feasible in several Alaska water bodies.



The invasion curve illustrates the increasing impacts and costs of invasive species over time and options for management at each stage of the invasion. Alaska is currently dealing with invaders at each stage of the invasion curve. Photos courtesy of ADF&G.

AQUATIC ASSETS NEED ATTENTION

A lack of resources is currently impeding the state's ability to respond to the spread of Elodea, on-going management of invasive northern pike in Southcentral, and the new European green crab infestation in Southeast Alaska.

The impacts from looming invasive species, such as zebra and quagga mussels, can be prevented in Alaska if surveillance, decontamination, and early detection efforts are adequately funded. Rapid response activities focused on southern Southeast Alaska can help prevent green crabs from destroying eelgrass nursery habitat. Early detection, monitoring, and functional eradication

methods to contain populations of invasive green crabs will reduce the impacts to areas where they occur and un-invaded areas. With additional resources to respond to new invasions and address management of existing invasive species, we can protect salmon fisheries and minimize degradation of aquatic habitats.

Alaska was once considered safe from invasive species, but times have changed. **Establishing a state invasive species council** would provide additional financial and administrative support to address invasions. You can help keep Alaska wild and free of invasive species.

AKISP BOARD OF DIRECTORS

Summer Nay, Chair

Salcha-Delta Soil and Water
Conservation District

Gino Graziano, Vice Chair

University of Alaska Fairbanks
Cooperative Extension Service

John Hudson, Secretary

Southeast Alaska Watershed Coalition

Ben Wishnek, Vice Secretary

United States Fish and Wildlife Service

Parker Bradley

Alaska Department of Fish & Game

Jen Chauvet

Homer Soil and Water Conservation
District

Amanda Jackson

Copper River Watershed Project

Cody Jacobson

Alaska Department of Natural Resources

Jess Parks

Alaska Department of Transportation &
Public Facilities

Nicole Swenson

Tyonek Tribal Conservation District

Hannah Thompson

Anchorage Soil and Water Conservation
District

Contact us at: akispboard@gmail.com

HOW DO INVADERS MOVE?

Invasive species are a threat to Alaska food security, recreation, and economic viability. Common means of transporting people, goods, and services to and around Alaska can be vectors of invasive species.

In freshwater, watercrafts and floatplanes can transport zebra and quagga mussels and Elodea, an invasive aquatic weed. A recent study by economists at the University of Alaska estimated potential damages from Elodea to the sockeye salmon fishery in Alaska at \$159 million dollars per year³.

On land, gravel pits, roadside brushing, and construction equipment can transport seeds of invasive plants, such as white sweetclover, along miles of roadsides. The nursery trade can deliver and sell invasive ornamental plants, including popular bird cherry trees. The ADNRC Alaska Community Forest Program has awarded over \$325,000 in grants to eradicate bird cherry trees in Southcentral and Southeast Alaska from 2020 to 2023.

Along the coast, commercial ships can disperse invasive crabs and sea squirts across continents in

³ Schwoerer et al. 2019. *Journal of Ocean and Coastal Economics* 6:2.

ballast water and on hulls. Mariculture equipment transported from out of state can be a haven for invasive parasites and hitchhiking organisms.

Even some unwanted pets can become invasive when released. Recently, ADF&G eradicated a population of spawning goldfish from Cuddy Pond in Anchorage, some as long as a woman's size 8 Xtratuf boot. In Fairbanks, Elodea was likely introduced into Chena Slough when dumped from an aquarium, then quickly spread.

PREVENTION continues to be the first and best line of defense against invasive species. It is also the most **COST-EFFECTIVE** way to limit introductions.

When prevention doesn't work, **EARLY DETECTION** is critical, followed by **RAPID RESPONSE** to control or remove the species. Whenever possible, **ERADICATION** is the best-case scenario. If that is infeasible, long-term **MANAGEMENT** of the population is often the next step. The goal is always to limit harm to environmental, cultural, and economic resources.



The USFWS, in coordination with US Customs & Border Protection and ADF&G, maintains a seasonal watercraft inspection and decontamination station at the Alcan Land Port of Entry, located on the Alaska-Yukon border, to intercept boats and other conveyances infested with zebra and quagga mussels or other aquatic invasive species. Thousands of boats have entered Alaska via this checkpoint in recent years with over half not inspected prior to arriving⁴.

⁴ Schwoerer et al. 2023. *Marine Policy* 148:105448 Photo: Ashley Lutto, USFWS



Look Out for Invasive Crab!

The green crab is invading from the west coast and has recently reached Alaska.

What is a green crab?


The green crab is considered one of the most invasive species in the marine environment. It has few predators, aggressively hunts and eats its prey, destroys seagrass, and outcompetes local species for food and habitat. It has been documented that green crab devour juvenile king crab as well as juvenile salmon. They also destroy eelgrass habitat that larval fish use to hide from predators, and outcompete Dungeness crabs for food and habitat. Green crab could potentially damage Alaska's multi-billion dollar fisheries industries, especially for salmon, crab, and mariculture operations. Resource managers in Alaska have been keeping an eye on the invasive crab's northward movement for years.



Invasive Green Crab. Photo: Linda Shaw, NOAA Fisheries

How did they get here?

Green crabs were first introduced to North America in the 1800s, likely hitching a ride in the ballast water of merchant ships from Europe. Experts believe the invasive crab was transported to the West Coast in ballast water as well. They may also be transported with shellfish, equipment, or packing materials in aquaculture operations. Larval green crabs can also spread from one invaded area to another in ocean currents. On July 19, 2022, the Metlakatla Indian Community Department of Fish & Wildlife confirmed the first observation of the invasive green crabs in the State of Alaska on Annette Islands Reserve. This map shows the range of green crab throughout the world with their native range in blue and all other colors indicating areas they invaded or have the potential to spread.



Range map of invasive crab (Carcinus maenas). Source: NOAA Alaska Region Fisheries

An aerial photograph of a stream flowing through a dense, green forest. The stream is the central focus, winding from the top right towards the bottom left. The surrounding vegetation is a mix of various shades of green, indicating a healthy, diverse ecosystem. The text is overlaid on the upper left portion of the image.

Emerging Invasive Species Concerns and Solutions in Alaska

Emily Reed

**Regional Invasive Plant Coordinator
Southeast Alaska Watershed Coalition**



**NC STATE
UNIVERSITY**



**Invasive Species
COLLABORATIVE AT VIRGINIA TECH**

Southeast Alaska Watershed Coalition






Panel: Grass Mix 1

✓	04/21/2022	Timothy	100,000 BAU	Prick	3/30,3+
✓	04/21/2022	Orchard	100,000 BAU	Prick	40/45,4+
✓	04/21/2022	Sweet Vernal	100,000 BAU	Prick	40/45,4+
✓	04/21/2022	Meadow Fescue	100,000 BAU	Prick	25/45,4+
✓	04/21/2022	June (KY Blue)	100,000 BAU	Prick	25/45,4+
✓	04/21/2022	Redtop	100,000 BAU	Prick	20/50,4+
✓	04/21/2022	Rye, Perennial	100,000 BAU	Prick	25/45,4+




What makes a non-native species **INVASIVE?**



Invasive species means a **nonnative species** whose introduction does or is likely to cause **economic** or **environmental harm** or **harm to human health**

Alaska Administrative Code Title 5, section 92.990



BIOSECURITY IMPACTS

Human Health

Animal Health

Infrastructure

Military Readiness

Forest Health

Salmon Habitat



Tiger Mosquito

BIOSECURITY IMPACTS

Human Health
Animal Health
Infrastructure
Military Readiness
Forest Health
Salmon Habitat



Photo: Free Range American

Brucellosis

BIOSECURITY IMPACTS

Human Health
Animal Health
Infrastructure
Military Readiness
Forest Health
Salmon Habitat



Photo: Japanese Knotweed Solutions Ltd

Knotweed

BIOSECURITY IMPACTS

Human Health

Animal Health

Infrastructure

Military Readiness

Forest Health

Salmon Habitat



Pigeon

BIOSECURITY IMPACTS

Human Health
Animal Health
Infrastructure
Military Readiness
Forest Health
Salmon Habitat



Rusty Tussock Moth

BIOSECURITY IMPACTS


Human Health
Animal Health
Infrastructure
Military Readiness
Forest Health
Salmon Habitat



Reed Canarygrass


Invasive Species in the US

iNaturalist: Observed Non-Native Species




Invasive Species in the US

iNaturalist: Observed Non-Native Species




Invasive Species in the US

USGS Non-Indigenous Aquatic Species




Invasive Species in the US


USGS Non-Indigenous Aquatic Species



STAGES of INVASION




STAGES of INVASION




STAGES of INVASION

ABSENT



LOCALIZED




Introduction


IMPACT & COST

TIME


AREA INVADED




STAGES of INVASION




STAGES of INVASION




PREVENTION




ERADICATION



CONTAINMENT



PROTECTION



PREVENT

Statewide



Quagga Mussel



Zebra & Quagga Mussels



Zebra & Quagga Mussels


Damage infrastructure
Impair power & water facilities
Reduce fish populations
.....
Spread and reproduce quickly
Difficult and costly to remove
.....
High Costs: \$300-\$500 million
per year in Great Lakes

Utah Division of Wildlife Resources



US Army Corps. Of Engineers

Zebra & Quagga Mussels




Reported Last State of Use for Watercraft Inspected at the Alcan Port of Entry (2024)



Zebra & Quagga Mussels




Annually, more than 1,000 watercraft from outside enter Alaska through multiple unprotected / not monitored ports of entry



Reported Last State of Use for Watercraft Inspected at the Alcan Port of Entry (2024)



ERADICATE/CONTAIN



ELODEA



ELODEA

Clog waterways

Decrease water quality

Threaten salmon habitat

.....

Regrow from small fragments

Difficult and costly to remove


.....

**Potential to reduce revenue by est.
\$97 million annually.** (Schwoerer et al.
2019)


Ashley Lutto, USFWS




ELODEA



ELODEA



Action on Invasive Species



State Policies

Effective: June 11, 2008

West's F.S.A. § 369.251

369.251. Invasive nonnative plants; prohibitions; study; removal; rules

Effective: July 1, 2019

Ga. Code Ann., § 27-3-24

§ 27-3-24. Hunting of feral hogs

§19-16A-1. Short title.

This article shall be known as the "West Virginia Pesticide Control Act of 1990".

Ark. Admin. Code 007.10.33-XI

007.10.33-XI. VECTOR AND RODENT CONTROL.

N.C.G.S.A. § 113-292

§ 113-292. Authority of the Wildlife Resources Commission in regulation of inland fishing and the introduction of exotic species

S.C. Code of Regulations R. 27-5

27-58. Asian Longhorned Beetle Quarantine.

Regs. Conn. State Agencies § 22-84-5g

Sec. 22-84-5g. Regulation of firewood transported into and within Connecticut

Example: Montana Mussel Program

Montana Aquatic Invasive Species Boat Inspections - 2024 Season

AIS Mussels Fouled Boats

42

Statewide Total

*Data does not filter by month

High Risk Boats

37,913

*Boats last launched in states with known AIS infestations

Total Watercraft Inspections

100,064


updated every Monday afternoon

Inspection Station Locations



Esri, HERE, Garmin, NGA, USGS

Out-of-State Boats Inspected



AIS Fouled Boats

Inspection Station - Anaconda
Date: 10/23/2024, 4:00 PM
Water Last Launched: Lake Michigan
State of Origin: IL

Inspection Station - Anaconda
Date: 10/23/2024, 4:00 PM
Water Last Launched: Lake Michigan
State of Origin: IL

Inspection Station - Hardin
Date: 10/14/2024, 4:00 PM
Water Last Launched: Lake Michigan
State of Origin: WI

Inspection Station - Wibaux
Date: 10/11/2024, 4:00 PM
Water Last Launched: Lake Michigan
State of Origin: MI

Inspection Station - Anaconda
Date: 10/7/2024, 4:00 PM
Water Last Launched: Lake Oahe
State of Origin: SD

*Data does not filter by month

National & Regional Coordination

NISC

NATIONAL INVASIVE SPECIES COUNCIL



Coordination

Alaska Invasive Species Partnership



The AKISP is an informal affiliation of interested individuals representing agencies, organizations, and members of the public.

Our goal is to heighten *awareness* of the problems associated with non-native invasive species and to bring about greater statewide *coordination*, *cooperation* and *action* to halt the introduction and spread of these invasive species.

Coordination

Alaska Invasive Species Partnership




Alaska Exotic Plant Information Clearinghouse

Alaska Invasive Plant Ranking System

Alaska Aquatic Invasive Species Clearinghouse


Annual Workshops

State Invasive Species Councils



State Invasive Species Councils


Fish & Game



Transportation



Agriculture




Natural
Resources



Economic
Development



Environmental
Quality




Health &
Human Services



Non-state members: county/municipal governments, industry, non-profits, large landowners, Universities, Tribal Governments, interest groups

State Invasive Species Councils

Fish & Game



Transportation



Agriculture



Natural
Resources



Economic
Development



Environmental
Quality



Health &
Human Services



Accountability

Transparency

Coordination


Authority

Responsiveness

Efficiency



Alaska Center for
Conservation Science
UNIVERSITY of ALASKA ANCHORAGE



Acknowledgements

Dr. Tobias Schwoerer, International Arctic Research Center

Danielle Verna, Prince William Sound Regional Citizens' Advisory Council

Tammy Davis, Alaska Department of Fish & Game

Alaska Invasive Species Partnership

Southeast Alaska Watershed Coalition

Representative Himschoot

Senator Dunbar



Questions?

Emily Reed: emily@sawcak.org | Tobias Schwoerer: tschwoerer@alaska.edu |
AKISP: akispboard@gmail.com

Preserving Alaska's resources by managing invasive species.



Aaron Martin
Alaska Regional Invasive Species
Program Coordinator
US Fish and Wildlife Service


March 11th, 2021



What is at stake?

- Infrastructure:
 - Hydropower facilities:
 - 21% of the state's power
 - Watercraft:
 - 68,616 registered watercraft (1/every 9 Alaskans)
 - **\$587M** annual economic impact
 - Floatplanes:
 - 114 floatplane bases (40% of all towns)
 - Lake Hood floatplane base
 - ✓ **\$56 Million** (labor + economic activity)
- Food security and industries
 - Sport fishing - \$2.4B
 - Salmon commercial fisheries - \$4.2B
 - 229 Federally recognized tribes, 12 ANCs
 - Hunting and wildlife viewing - \$7.5B





Legend

Non-native plant occurrence

1 45 80 132 175

Number of species present

Map tools

Visualize on Polygon tool

Location types

- Non-native plant population detected in most recent survey
- No high-priority non-native plant populations detected in most recent survey
- No non-native plant population detected in most recent survey

Download all data


Species

Search Invasiveness Category

Species	Invasiveness	Downloads
<input checked="" type="checkbox"/> <i>Acer negundo</i> L. boxelder		

69 5362 -172 5195

Credit: State of Victoria, Department of Primary Industries, 2010.



First step in prevention is to understand risk = vectors and suitability.

Prevention: Quagga and Zebra Mussels



Zebra Mussels



Quagga Mussels

Zebra and Quagga Mussel Sightings Distribution


Dreissena polymorpha and *D. rostriformis bugensis*



- Zebra mussel occurrences
- Quagga mussel occurrences
- ▲ Zebra or Quagga mussel occurrences
- Both species occurrences
- Zebra/Quagga mussels eradicated
- Zebra/Quagga mussels failed



Western Watercraft Inspection and Decontamination Stations




**USFWS Role:
Lacey Act of 1900
(amended 2008)**

Results



- Research suggests high habitat suitability for invasive mollusks.
- No confirmed reports of invasive mollusks in the wild.
 - Smith et al. 2005
 - D. Bogan 2012 – AKISP presentation Kodiak 2012



- Alaska has 3 native mussels and 21% of state's power comes from hydropower facilities.
- 226 watercraft inspected (2017-19) at Alcan Port of Entry:
 - **70%** not inspected in route, 30% inspected in-route
 - **38%** coming from a state with Q/Z mussels, 62% coming from states without
- No live mussels detected yet, but...
- Currently assessing other critical control points in BC, WA, and OR.

Sailboat being decontaminated, Tok AK July 2019



PSA – Zebra Mussels in moss balls

March 3, 2021



U.S. Fish & Wildlife Service

Fish and Aquatic Conservation

Search



[Home](#) [About Us](#) [National Fish Hatchery System](#) [Species & Habitat Conservation](#) [Invasive Species](#) [Recreation & Education](#) [In The News](#) [You Can Help](#)

You Are Here: [Fisheries Home](#) » [Aquatic Invasive Species](#) » [Zebra Mussel Disposal](#)

Destroy! Don't Dump!

Invasive zebra mussels have been found in "moss balls," an aquarium plant product sold at aquarium and pet supply stores. Zebra mussels are regarded as one of the most destructive invasive species in North America.

Zebra mussels can quickly take over once they get established in a waterbody and cause significant damage including disrupting the food chain, changing the chemistry of the water (which can cause more blue green algae outbreaks or offensive taste), and clogging water intake and delivery systems. The concern is that live mussels released into a storm drain or flushed could be introduced into a waterway.

Quick Links

- [Aquatic Nuisance Species Task Force](#)
- [Habitattitude](#)
- [Stop Aquatic Hitchhikers](#)
- [Aquatic Invasive Species Coordinators](#)



Moss ball and Zebra mussel. Photo by IUH-W




Zebra mussels can quickly take over once they get established in a waterbody and cause significant damage

Originating locations of watercraft crossing into AK, summer 2017-2019



Legend

Number of watercraft



Alcan border

Infected states



STOP AQUATIC HITCHHIKERS!
Be A Good Steward.
Clean. Drain. Dry.
StopAquaticHitchhikers.org

Keep Alaska Wild & Free of Invasive Species

Green leaves in clusters of 3, occasionally 4 →



Long trailing stem →

Elodea

Grows in a tangled mass
Can survive when frozen in ice
Endangers safe float plane operation
Degrades fish habitat and displaces native plants
By law, cannot be brought into or moved around the state
Can form new plants from broken segments, roots, and seeds
Makes boat travel difficult and reduces recreation opportunities



ELODEA HAS BEEN FOUND
IN THESE AREAS



ELODEA COULD COST
RECREATIONAL FLOAT PLANE
PILOTS AND COMMERCIAL
FISHERIES \$100-\$500M / YEAR
IF NOT ADDRESSED NOW



STOP AQUATIC HITCHHIKERS!

Be A Good Steward. Clean. Drain. Dry.

StopAquaticHitchhikers.org

✓ CLEAN	✓ DRAIN	✓ DRY
Remove all visible mud, plants, and fish/animals from equipment	Eliminate water from all equipment before transporting; pull the plug	Dry everything for at least five days OR dry thoroughly before next launch

Report anything suspected to be an **invasive species**:

📍 Note its location: Get GPS coordinates

📝 Describe its habitat

📷 Snap some photos

📞 INVASIVE HOTLINE: 1-877-INVASIV (468-2748)



Prevention and Eradication: Elodea

- Background:
 - Alaska's 1st submerged aquatic invasive plant
 - Native to the Pacific Northwest and New England
 - Introduced through aquarium dumps
- Impacts:
 - Habitat degradation/ loss for fish and wetland obligate species
 - Reduced biodiversity, fishing opportunities, floatplane and watercraft safety
 - Increased sedimentation



FAIRBANKS SOIL & WATER
CONSERVATION DISTRICT



Homer Soil & Water
CONSERVATION DISTRICT

Keep Alaska Wild & Free of Invasive Species

Green leaves in clusters of 3, occasionally 4 →



Elodea

Long trailing stem →

- Grows in a tangled mass
- Can survive when frozen in ice
- Endangers safe float plane operation
- Degrades fish habitat and displaces native plants
- By law, cannot be brought into or moved around the state
- Can form new plants from broken segments, roots, and seeds
- Makes boat travel difficult and reduces recreation opportunities

ELODEA HAS BEEN FOUND IN THESE AREAS

ELODEA COULD COST RECREATIONAL FLOAT PLANE PILOTS AND COMMERCIAL FISHERIES \$100-\$500M / YEAR IF NOT ADDRESSED NOW

STOP AQUATIC HITCHHIKERS!
Be A Good Steward. Clean. Drain. Dry.
StopAquaticHitchhikers.org

CLEAN	DRAIN	DRY
Remove all visible mud, plants, and fish/animals from equipment	Eliminate water from all equipment before transporting; pull the plug	Dry everything for at least five days OR dry thoroughly before next launch

Report anything suspected to be an **invasive species**:

- Note its location: Get GPS coordinates
- Describe its habitat
- Snap some photos

INVASIVE HOTLINE: 1-877-INVASIV (468-2748)

Prevention and Eradication: Elodea

- Economic analysis:
 - Impact Potential: Annual loss of \$159M to the sockeye salmon fisheries if not stopped (Schwoerer et al 2019).
 - Ship-borne AIS impacts in the Great Lakes: Annual cumulative loss of \$138M/year to sportfishing, commercial fishing and water use. (Rothlisberger et al 2019).

Photo credit: Jason Ching



Photo credit: USFWS



Known Elodea Infestations in Alaska

Eradication
in progress

Manley Slough
Bathing Beauty Pond
Birch Lake
Chislom Lake
Harding Lake
Piledriver Slough

Fairbanks:
Chena Slough
Chena River
Chena Lake
Totchaket Slough

Anchorage:
~~Lake Hood~~
~~Sand Lake~~
~~Little Campbell Lake~~
~~Delong Lake~~
Potters Marsh
Jewell Lake

Eradication
in progress

Mat-Su:
Alexander Lake
Sucker Lake Complex
Big Lake

Eradication
beginning
2020

Cordova:
Eyak Lake
Eyak River System
Bering Lake
McKinley Lake
Martin Lakes
Odiak Lagoon
Wooded Pond
~~Wrong Way Pond~~
Alaganik Slough System


Eradication
in progress
by USFS

Kenai Peninsula:
~~Stormy Lake~~
~~Daniels Lake~~
~~Beck Lake~~
North South Lakes
~~Sport Lake~~

Sandpiper Lake
Eradication
beginning
2020




(a)



Freshwater Vectors – Floatplanes

Floatplanes' first-leg flight paths between freshwater start and destination locations. Data from a survey with pilots about their 2015 flights. Schwoerer et al. 2017.





Prevention and Preparedness: European Green Crabs

- Introduced through ballast water and now currents
- Considered one of world's worst 100
- Impacts:
 - Eats native oysters, clams, mussels
 - Damages estuaries and bays

European green crab in Puget Sound

© 2011 Oregon Coast, Monitoring, Fish, Oyster



Green crabs can be identified by their unique shell shape. Adults can have shells up to four inches across in width.


Credit: [NOAA](#)



Prevention and Preparedness: European Green Crabs

Commercial boating traffic in the North Pacific (2015).


Droghini et al. 2020





Prevention and Preparedness: European Green Crabs

Receivers



Contributors

Transit origins outside of the Bering Sea and the Bering Sea destination, 2014-2016 for **A) all marine vessels** and **B) commercial fishing vessels**.

Key Takeaways

- Alaska's way of life and industries are at risk.
- Keep invasive species diversity and distribution low.
- Solidify foundation:
 - Increase awareness and prevention
 - Build on organization and capacity among partnerships



KEEP ALASKA WILD AND FREE... OF INVASIVE PLANTS AND ANIMALS

TO LEARN MORE ABOUT INVASIVE SPECIES AND THE DAMAGE THEY DO IN YOUR AREA: GO TO THE INVASIVE SPECIES PAGE AT WWW.ADFG.ALASKA.GOV




Aaron Martin, Regional Invasive Species Program Coordinator

Aaron_e_martin@fws.gov

907-378-0568 (cell), 907-786-3510 (Desk)

FOUND IN SOUTHEAST ALASKA INVASIVE EUROPEAN GREEN CRAB

How to identify a European green crab?



Native crabs found in Southeast Alaska

GREEN SHORE CRAB



HAIRY HELMET CRAB



SPOTTED A EUROPEAN GREEN CRAB? NOW WHAT?

FIND IT

If you find a European green crab, do not throw it back alive!

CONTAIN IT

Keep it in a container with the date and location you found it.

PHOTOGRAPH IT

Photos help with identification. Take a photo of the top of the crab shell then submit the photo to ADF&G.

FREEZE IT

Freeze it or preserve it with rubbing alcohol.

REPORT IT

Call the free statewide invasive species hotline at 1 (877) INVASIV. We need the collected crab to confirm its identification.

**SALMON AND SHELLFISH HABITAT IS
DESTROYED BY EUROPEAN GREEN CRABS.**



Scan the QR Code or call the invasive species hotline at 1-877-INVASIV to report observations of any invasive, nonnative species, or unusual plants or animals.



I Spy a Changing Ecosystem

IMPACTS OF INVASIVE EUROPEAN GREEN CRAB IN ALASKA

What do you spy?

Before EGC Invasion

1. Healthy marine ecosystems support subsistence, recreation, and commercial fishing activities
2. Shorebirds feed on abundant fish and invertebrates
3. Vibrant intertidal zone with *fucus* seaweed, mussels, clam beds, fishes, and eelgrass beds
4. Native crustaceans, such as juvenile Dungeness and shore crabs, live among rocks and eelgrass beds
5. Juvenile fish, such as salmon, use eelgrass beds for protection as the grow

After EGC Invasion

6. EGC arrive in the ecosystem and compete with native crabs
7. EGC dig up eelgrass beds while foraging for food and digging burrows
8. EGC eat native shellfish species at high rates
9. EGC females release nearly 200,000 eggs per year
10. EGC eggs hatch into planktonic larvae that float on currents and settle in new habitats as juvenile crabs
11. Once EGC have become established, the best management tool is intensive trapping and removal

What are European Green Crab?


The European Green Crab (EGC) is considered one of the most detrimental invasive species in the world due to its harmful impacts on nearshore ecosystems. These invasive crabs thrive in protected estuaries and bays with muddy intertidal zones and eelgrass beds. They can survive in a wide range of environmental conditions.

How to Find Them

- Look for carapaces (molts) while walking on the beach
- Look for them in your crab traps

How to Identify Them

- Five spines (marginal teeth) on the outer edge of the shell, with three bumps on the shell between the eyes
- Color varies widely - they can appear green, yellow, brown, or reddish
- Adults can grow up to 4 inches wide across the carapace, but juveniles are much smaller



How do they Impact Ecosystems?

EGC are aggressive and can compete with native crab species like juvenile Dungeness and smaller shore crabs for the best hiding spots and food. While looking for food such as juvenile clams and crabs, EGC destroy eelgrass beds which are critical habitat for many other species, including salmon. Their juvenile life stages (planktonic larvae) travel on ocean currents and settle in new habitats, allowing EGC to spread rapidly along the Pacific coast and into Alaska's nearshore waters.

To protect the fishes, shellfish, birds, and mammals that call our estuaries home, we need all Alaskans to help by keeping an eye out for EGC and reporting any sightings to the Alaska Department of Fish and Game (ADF&G). You can participate in early detection of EGC by searching for molts when walking beaches or by participating in the Alaska community-based monitoring network. Call ADF&G for more information.

Scan the QR code to learn how you can help protect Alaska's ecosystems and communities from the impacts of invasive EGC. Call 1-877 - INVASIV to report sightings of EGC or other invasive species.



RATS!

What to know before buying, selling, or breeding rats in Alaska

Not all pet rats are legal

State regulations prohibit the sale and ownership of all rats that are not albino. Albino rats must have fur that is completely white *and* pink eyes. Some municipalities have stricter regulations than the state. For example, in Anchorage, all rats are prohibited as pets, even albino rats. Know what is legal where you live.

Rats carry consequences

Rats can carry diseases transmissible to humans, wildlife, and pets. Rats can also compete with or predate upon native species. If buying or breeding albino rats, make sure that they are kept enclosed. Releasing rats, even by accident, is illegal and may result in a citation and fines. Rats can have negative consequences for human and wildlife health, wildlife populations, and wildlife habitat.

Help prevent outbreaks

Female rats can give birth to more than a dozen pups every 4-5 weeks, which can result in hundreds of offspring per year. Rat "outbreaks" can start with a single pregnant female.




Use of traps and/or rodenticide are allowed to remove rats and prevent the spread of a harmful invasive species. Please report sightings of released or wild rats to ADF&G.

Releasing any pets is illegal

Releasing any pets outdoors to the "wild" is against the law in the state of Alaska. If you encounter released pets, please report them to ADF&G.



HELP KEEP ALASKA RAT FREE!

-  Only albino rats may be kept as pets.
-  Do not release rats or any other pets.
-  Report rat sightings to ADF&G.

Contact us!

For more information, or to report sightings of rats or other harmful invasive species:

Call: 1-877-INVASIV

Email: dfg.dsf.InvasiveSpecies@alaska.gov

Visit online: adfg.alaska.gov, and search "invasive species"



You can help stop the spread of Elodea



Elodea is an invasive plant that easily spreads and ruins fish habitat and clogs waterways.

Don't let Elodea hitch a ride on you or your equipment. Check and clean your gear.



For more information or to report an invasive plant sighting, please contact the Alaska Department of Natural Resources at (907) 745-8721.
For more information or to report an invasive animal sighting, please contact the Alaska Department of Fish and Game at (877) 468-2748.



ELODEA



spread the word,
not the weed!

Elodea is a highly invasive submerged aquatic plant. It survives freezing & spreads by tiny fragments introduced by float planes, boats, trailers, school/home aquariums, and equipment.

Elodea is an unwanted invasive species Alaska.

Why no one wants *Elodea* in Alaska because it:

- Fouls float plane rudders & boat propellers causing a hazard,
- Impedes boat & float plane launching, navigation, and fishing,
- Degrades salmon spawning habitat,
- Reduces property and recreational value by fouling launch sites, fly-in hunting and fishing sites, and aquatic habitat.

You can make a difference by inspecting and cleaning your watercraft & gear

- Before launching/takeoff inspect watercraft; remove visible plants and/or debris and pump water from floats
- Before takeoff: raise and lower rudders to free plant fragments. Avoid taxiing through heavy plant growth. After take off: free plant fragments over the waters you just left or dry land
- Always inspect watercraft and boat trailers; remove all plant debris and mud.

Report sightings & help identify new infestations

- Note location (GPS or mark on map) and water depth/clarity
- Take a photo of the plant
- Off base, call 1-877-INVASIV (468-2748) to report sightings!
- On base, call JBER Natural Resources at 907-384-3212
- Report sightings online to Alaska Department of Fish and Game



**STOP AQUATIC
HITCHHIKERS!™**

Be A Good Steward.
Clean. Drain. Dry.

StopAquaticHitchhikers.org

Support eradication efforts!



Attention Anglers!

If you catch a northern pike in this lake, please harvest the fish and report it to ADF&G.



To report northern pike or any other potential invasive species, scan the QR code, or call 1-877-INVASIV.

- Northern pike are an invasive species in Southcentral Alaska.
- It is illegal to introduce fish or any other organisms into state waters.

Northern pike

Rarely seen under 6"; large bodied; elongate body shape with light mottling on a greenish background; forked tail fin.



Fish in this lake that could be misidentified as northern pike:

Alaska Blackfish:

(Nonnative to lakes in Southcentral)

Less than 10"; Blunt snout with rounded tail fin, dark brown mottling on a tan background.



Sticklebacks:

(Native to lakes in Southcentral)

Less than 4"; Often seen in schools; Very common.



DON'T LET IT LOOSE!



Releasing an unwanted pet into the wild is never the right thing to do. Your released pet can cause unanticipated ecological damage, carry pathogens that harm fish and wildlife, and it's illegal.



***Give your unwanted pets another chance,
contact local vendors to find them a new home.***

- ***Alaska Coral and Fish: (907) 929-833***
- ***Petzoo: (907) 344-2966***
- ***PetSmart: (907) 522-7387***
- ***Petco: (907) 365-5500***

If unable to find a new home, it is more humane to euthanize than release.

www.adfg.alaska.gov