

MIND, DISRUPTED

How toxic chemicals may change how we think and who we are



Polybrominated Diphenyl Ethers (PBDEs)

What Are PBDEs?

Polybrominated diphenyl ethers, or PBDEs, are a class of flame retardant chemicals added to many consumer products found in the home, office, automobiles, and airplanes.¹ Three mixtures used widely—penta-BDE, octa-BDE, and deca-BDE—made up 14%, 6%, and 80% of the 1999 worldwide production of PBDEs, respectively.² Usually found in electronics, such as TVs, and used in some furniture foams, fabrics, and kitchen appliances, the industry voluntarily ended production in the United States of the formulations of penta and octa in 2004 after high levels were found in breast milk. However, North America made up a large percentage of global demand for these PBDEs, 97.5 % for penta and 35.9% for octa, and there is still a large reservoir of products containing these PBDEs in the US.³ The deca formulation is currently produced and used primarily in plastic electronics such as televisions and computer casings. It is also used in the upholstery covers of items such as furniture and car seats.^{4,5}

PBDEs are part of a larger chemical class called polyhalogenated aromatic hydrocarbons (PHAHs) which include other highly toxic chemicals such as polychlorinated biphenyls (PCBs) and dioxins.² PHAHs are intrinsically hazardous because of their chemical make-up: (1) they are stable, meaning they persist in the environment and do not break down easily; (2) they are lipophilic, meaning they build up in fatty tissues of living organisms; and (3) they have toxic properties, including the potential to act as endocrine disruptors.⁶ Their persistence and fat solubility allow them to both biomagnify and bioaccumulate, meaning they build up in the bodies of animals and humans as they move through the food chain. While not all PBDEs meet these criteria, a significant portion have been found to exhibit these characteristics. Although the main component in deca-BDE, BDE-209, has a relatively short half-life in people,¹ animal studies show that the liver breaks down BDE-209 into the more persistent and bioaccumulative forms of hepta-BDE, octa-BDE, and nona-BDE.^{7,8} There are relatively few data available concerning the extent to which this occurs in humans.

How Are We Exposed?

There are many ways that humans are exposed to PBDEs, including ingestion of contaminated foods and incidental intake of PBDE-contaminated dust.⁸ PBDEs are not permanently bound to the products in which they are used. Some PBDEs are semi-volatile and may be released from PBDE containing products. Other PBDEs are released from furniture, electronics and other products as they physically degrade. It is not entirely clear how deca is released from products, though some think it might be released from physical abrasion or deterioration of the product. Deca-BDE breaks down into more toxic by-products and may volatize when exposed to ultraviolet (UV) light.⁹ Although penta-BDE and octa-BDE are no longer produced in the United States, exposure continues from old computers, furniture, fabrics, and other consumer sources that were made before the discontinuation.¹⁰

Occupational exposure may also occur for workers in electronics recycling, computer repair, and rubber manufacturing facilities.¹¹ PBDEs are found in sewage sludge used as agricultural fertilizers and may be a source of exposure for workers applying contaminated fertilizers.¹¹

Ingestion of PBDEs in fatty foods such as meat and dairy products is thought to be a major route of exposure,¹² as PBDEs are fat-seeking and build up in the food web. Indoor exposures to dust and contaminated air are also thought to be major routes, as well as direct absorption through the skin from dust or products.¹³ Relatively little is known about the routes of deca-BDE exposure. It is estimated that the highest intake of deca-BDE results from household dust.^{13,14,15}

People may also be exposed through consumption of fish and other marine animals, because these organisms have been found to contain high levels of deca-BDE.¹⁶ The high levels in marine environments may be due to atmospheric deposition (when contaminants from the air come down to the earth's surface through rain, snow, falling particles, and the absorption of gas) of deca-BDE as well as sewage discharge into the oceans.¹⁶

For infants, breast milk is a major source of PBDEs. The PBDEs that have accumulated in women may be passed to children through breastfeeding.¹⁷ It is important to note that breastfeeding is still considered best for a baby's health.

PBDEs in Our Bodies

Due to their widespread use, persistence and bioaccumulative properties, PBDEs have been found in humans at high levels. A recent study by scientists from the U.S. Centers for Disease Control and Prevention (CDC) found PBDEs in nearly all 2,040 participants in a sample representative of the 2003–2004 U.S. population.¹⁸ In this study, 3 congeners were found in over 90% of the participants: 97% had levels of BDE-47 (2,2',4,4'-hexabromodiphenyl ether), 93% had levels of BDE-100 (2,2',4,4'-hexabromodiphenyl ether), and 93% had levels of BDE-153 (2,2',4,4',5,5'-hexabromodiphenyl ether) above the limit of detection.¹⁸ Another study showed that 5 percent of American women have levels of PBDEs that are close to the levels linked to reproductive problems in animals.¹ PBDEs have been found in mothers' breast milk and in the blood of mothers and their babies.^{19,20} People of the Arctic may experience an even higher risk of exposure due to their traditional subsistence diet rich in fat from marine mammals.²¹ Concentrations of PBDEs have grown over the years in marine mammals due to atmospheric transport of chemicals into the north (transported long distances from areas of production and use via air and ocean currents) and bioaccumulation.²

Reducing Your Exposure

You can minimize your exposure to PBDEs by taking the following steps:

PBDE-free Furniture

- Before purchasing furniture, find out which companies offer PBDE-free products. The following websites can help:
 - Clean Production Action: www.cleanproduction.org/FlameAlternatives.php
 - Pollution in People: www.pollutioninpeople.org/safer/products
- Contact the company directly if you cannot figure out if the manufacturer uses PBDEs.
- Choose furniture made with less flammable fabrics like leather, wool and cotton.

PBDEs in Foam Padding

- Foam items purchased before 2005 can potentially contain PBDEs. Completely covering these items in fabric can limit potential exposure.
- Avoid reupholstering foam furniture.
- Be very careful when removing old carpet. Try to keep your work area separated from the rest of the house and thoroughly clean up the area.

Reduce Your Dust Exposure

- Wash your hands often to remove dust particles that your hands pick up throughout the day on everything you touch.

What Does Exposure to PBDEs Mean for Our Health?

The presence of environmental chemicals in the human body does not necessarily imply that they are causing adverse health effects; however, environmental chemical exposures can and do affect human health. It is important to note that both the dosage and the timing of exposure have significant effects on any potential health outcome.

The following information is intended to inform the reader about the current state of knowledge on the health effects of PBDEs, including both human and animals studies.

Relatively little is known about the health effects of PBDEs. Overall there are few studies on health effects in the general population; however, health effects have been detected in rigorous animal based studies. Nevertheless, research has found associations between PBDEs and many adverse health effects, including:

Neurodevelopmental Effects

There are few published human studies on the health effects of environmental exposure to PBDEs. One study on Dutch children found that prenatal PBDE exposure affected childhood development. Exposed children had worse fine motor skills and

- Use a wet rag or cloth while dusting to avoid kicking up the dust in the air.
- If possible, use a vacuum fitted with a HEPA filter. These vacuums can trap smaller particles of dust and will be more likely to remove contaminants from your home.
- Vehicles have been exempt from recent PBDE laws and high levels have been found inside cars.⁵ Removing dust with a wet cloth and keeping car seat cushions in good repair will help to reduce your exposure.

PBDE-free Electronics

- Many companies are beginning to make electronics with alternatives to PBDEs. Certain PBDE-free products are available from Canon, Dell, HP, Intel, Erickson, Apple, Acer, Nokia, Motorola, LG Electronics, and Sony.

Consider Eating Less Fat:

- Consider choosing leaner meat and poultry cuts.
- Consider removing fat that you see on meat and fish whenever possible.
- Choose cooking methods that remove excess fat such as broiling, grilling, and roasting.

attention, but better coordination, visual perception and behavior. The mothers in this study had serum PBDE concentrations in the low ppb range.²² Another study of children prenatally exposed to PBDEs showed decreases in mental and physical development. PBDEs 47, 99 and 100 were correlated with lower developmental scores, while PBDE 153 was not. The most highly exposed children had significantly lower developmental scores than the less exposed children.²³

A recent peer-reviewed scientific consensus statement by the Collaborative on Health and the Environment's Learning and Developmental Disabilities Initiative states, "Recent studies have left little doubt that PBDEs are developmental neurotoxicants in animals and lead to changes in motor activity and reduced performance on learning and memory tests."^{24,25,26} Numerous studies on rodents suggest that neonatal exposure to PBDEs permanently affects learning and memory functions, impairs motor activity, and is linked to aberrations in spontaneous motor behavior and hyperactivity.^{27,28,29,30,31,32,33}

Researchers have also found that some of the developmental effects of exposure become worse over time. In a study of PBDE exposure in mice, spontaneous behavior and habituation capability (both indicators of the ability of individuals and the species to survive in the wild) were permanently damaged by neonatal exposure to PBDEs and worsened with age.²⁷ Exposure to hexa-BDE has been linked to other developmental neurotoxic effects in mice, including changes in spontaneous behavior and impairments in learning and memory, which also appear to worsen with age.³⁴ A recent study in adult mice found consistent results which show that deca-BDE can be just as toxic to neurodevelopment as other PBDEs. In this study, exposure to deca-BDE was linked to hyperactivity and reduced or lack of habituation, also worsening with age.³⁵

Thyroid Problems

Some PBDE congeners are structurally similar to thyroid hormone and have been shown to disrupt (by decreasing, increasing, or mimicking) the biological action of thyroid hormone.³⁶ In a study of newborn babies, high PBDE levels in cord blood were associated with decreased levels of thyroid hormones. PBDE concentrations ranged from below 10 ppb to several hundred ppb in the blood.³⁷ In a recent study of adult men who consume sport fish, researchers found that exposure to PBDEs at levels similar to those found in the general US population were associated with increased thyroid hormone T₄ (thyroxine).³⁸ Men in this study with the highest body burdens of PBDEs also showed increased thyroid antibodies, a risk factor for thyroid diseases such as chronic autoimmune thyroiditis and Graves' disease.³⁸ PBDE exposure has also been linked to hypothyroidism in adults. In an occupational exposure study at a deca-BDE and deca-bromobiphenyl manufacturing plant, 4 out of 35 exposed workers showed clinical hypothyroidism, while none of the 89 unexposed workers exhibited thyroid dysfunction.³⁹

Corresponding animal studies have also shown that PBDE exposure is linked to decreased circulating concentrations

of thyroid hormone^{14,40} and decreased thyroid weight in adult rodent offspring.¹² Exposure to deca-BDE has been linked to thyroid hyperplasia in rodent studies. In a 2-year study by the National Toxicology Program, deca-BDE was found to increase the incidence of follicular cell hyperplasia (proliferation of follicular cells, which are responsible for making thyroid hormones) in the thyroid gland in male and female mice.⁴¹

Reproductive Effects

PBDEs can be both mildly estrogenic⁴² and anti-androgenic compounds.^{43,44} They have been correlated to cryptorchidism (undescended testes), in newborn boys exposed through breast milk with a median concentration of 4.16 ppb(lw) vs. 3.16 ppb(lw) for controls.⁴⁵ PBDEs have been shown to permanently impair sperm development in rodent studies.¹⁴ Exposure to BDE-153, has been associated with a decrease in testicle size and the sperm concentration in humans.⁴⁶ Exposure to PBDEs in household dust at levels commonly encountered in the US has been linked with lower levels of androgens (male hormones) in adult men.⁴⁷ PBDEs have also been associated with delay of puberty in both male and female rodents and changes in sexual development and gender-specific sexual behavior.⁴⁸ Exposure to PBDEs has been linked to low birth weight, birth defects, reduced weight gain during pregnancy, changes in ovary cells and reduced sperm count.^{49,50} The breakdown products of PBDEs may inhibit human aromatase, an enzyme important in the formation of androgens and estrogens (male and female hormones), and in skeletal development. However, these results were found at concentrations above current human exposure levels.⁵¹

Cancer

One study suggests that *in utero* exposure to PBDE concentrations is associated with an increased risk of testicular cancer in men, although the study has limitations and further studies would help to shed light on this association.⁵² The Agency for Toxic Substances and Disease Registry (ATSDR) lists deca-BDE as a possible human carcinogen based on the development of liver tumors in rats.⁸

Regulation of PBDEs

There have been many steps forward in the regulation of PBDEs in recent years, but much more is needed to ensure the safety of public health. Internationally, Sweden was the first to initiate a phase-out of PBDEs in the late 1990s, followed by the European Union—first phasing out penta-BDE and octa-BDE in 2003 and then deca-BDE in 2006.⁵³ In 2008, Norway banned the use of deca-BDE in new consumer products.⁵⁴ Although no production occurred there, Canada prohibited the future manufacture of all PBDEs and placed restrictions on the import of certain PBDEs.⁵⁵

The Stockholm Convention on Persistent Organic Pollutants, a United Nations treaty on persistent bioaccumulative substances, placed tetra-, penta-, hexa- and hepta-PBDEs under provisions

of the Convention for global phase-out after determining they posed a risk to environmental and human health.

In the United States, industry voluntarily ended production of penta-BDE and octa-BDE in 2004, but deca-BDE is still permitted for use in consumer products.²⁰ In December, 2009 two U.S. producers of deca-BDE and the largest U.S. importer announced

commitments to end production, importation, and sales of deca-BDE for most uses by December, 2012.⁵⁶ Washington, Maine, Oregon, and Vermont are currently discontinuing the use and manufacture of deca-BDE,^{57,58} and 12 states have banned penta- and octa-BDE, with other states across the nation aiming for similar legislation.

Endnotes

- 1 McDonald TA. 2005. Polybrominated diphenylether levels among United States residents: Daily intake and risk of harm to the developing brain and reproductive organs." *Integrated Environmental Assessment and Management* 1(4):343-354.
- 2 Ikonomou MG, Rayne S, Addison RF. 2002. Exponential increases of the brominated flame retardants, polybrominated diphenyl ethers, in the Canadian Arctic from 1981 to 2000. *Environmental Science & Technology* 36(6):1886-1892.
- 3 Hale RC, Alae M, Manchester-Neesvig JB, Stapleton HM, Ikonomou MG. 2003. Polybrominated diphenyl ether flame retardant in the North American environment. *Environmental International* 29:771-779.
- 4 Toxic-Free Legacy Coalition Report. 2006. Pollution in People. Available: www.toxicfreelegacy.org.
- 5 Lagalante AF, Oswald TD, Calvosa FC. 2008. Polybrominated diphenyl ether (PBDE) levels in dust from previously owned automobiles at United States dealerships. *Environment International* 35(3):539-44.
- 6 Hooper K, McDonald TA. 2000. The PBDEs: An emerging environmental challenge and another reason for breast-milk monitoring programs. *Environmental Health Perspectives* 108(5):387-392.
- 7 Stapleton HM, Brazil B, Holbrook RD, Michelmore CL, Benedict R, Konstantinov A, Potter D. 2006. In vivo and in vitro debromination of decabromodiphenyl ether (BDE 209) by juvenile rainbow trout and common carp. *Environmental Science and Technology* 40(15): 4653-4658.
- 8 Van der ven LT, van de Kuil T, Leonards PE, Slob W, Canton RF, Germer S, Visser TJ, Litens S, Hakansson H, Schrenk D, van der Berg M, Piersma AH, Vos JG, Opperhuizen A. 2008. A 28-day oral dose toxicity study in Wistar rats enhanced to detect endocrine effects of decabromodiphenyl ether (decaBDE). *Toxicology Letters* 179(1): 6-14.
- 9 Washington State Department of Health and Washington State Department of Ecology. 2006. Washington State Polybrominated Diphenyl Ether (PBDE) Chemical Action Plan: Final Plan. Available: www.ecy.wa.gov/pubs/0507048.pdf
- 10 Northwest Watch. 2004. Flame retardants in the bodies of pacific northwest residents: A study of toxic body burdens. Available: www.sightline.org/
- 11 Sjödin A, Patterson Jr DG, Bergman Å. 2003. A review on human exposure to brominated flame retardants—particularly polybrominated diphenyl ethers. *Environment International* 29(6): 829-40.
- 12 Agency for Toxic Substances and Disease Registry (ATSDR). 2004. Toxicological Profile for Polybrominated Biphenyls and Polybrominated Diphenyl Ethers. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry. Available: www.atsdr.cdc.gov/toxprofiles/tp68.html.
- 13 Frederiksen M, Vorkamp K, Thomsen M, Knudsen LE. 2009. Human internal and external exposure to PBDEs – a review of levels and sources. *International Journal of Hygiene and Environmental Health* 212(2): 109-134.
- 14 Wilford BH, Shocib M, Harmer T, Zhu J, Jones KC. 2005. Polybrominated diphenyl ethers in indoor dust in Ottawa, Canada: implications for sources and exposure. *Environmental Science and Technology* 39(18): 7027-7035.
- 15 Huwe JK, Hakk H, Smith DJ, Dilberto JJ, Richardson V, Stapleton HM, Birnbaum LS. 2008. Comparative absorption and bioaccumulation of polybrominated diphenyl ethers following ingestion via dust and oil in male rats. *Environmental Science and Technology* 42(7): 2694-2700.
- 16 Ross PS, Couillard CM, Ikonomou MG, Johannessen SC, Lebeuf M, Macdonald RW, Tomy GT. 2009. Large and growing environmental reservoirs of Deca-BDE present an emerging health risk for fish and marine mammals. *Marine Pollution Bulletin* 58(1): 7-10.
- 17 Frederiksen M, Vorkamp K, Thomsen M, Knudsen LE. 2009. Human internal and external exposure to PBDEs-A review of levels and sources. *Int J Hyg Environ Health* 212:109-134
- 18 Sjödin A, Wong L-Y, Jones RS, Park A, Zhang Y, Hodge C, Dipietro E, McClure C, Turner W, Needham LL, Patterson Jr. DG. 2008. Serum Concentrations of Polybrominated Diphenyl Ethers (PBDEs) and Polybrominated Biphenyl (PBB) in the United States Population: 2003–2004. *Environ Sci Technol* 42(4): 1377-84.
- 19 Schecter A, Pavuk M, Papke O, Ryan JJ, Birnbaum L, Rosen R. 2003. Polybrominated diphenyl ether (PBDEs) in U.S. mother's milk. *Environmental Health Perspectives* 111(14):1723-1729.
- 20 Mazdai A, Dodder NG, Abernathy MP, Hites RA, Biggs RM. 2003. Polybrominated diphenyl ethers in maternal and fetal blood samples. *Environmental Health Perspectives* 111(9):1249-1252.
- 21 Letcher RJ, Bustnes JO, Dietz R, Janssen BM, Jorgensen EH, Sonne C, Verreault J, Vijayan MM, Gabrielsen GW. 2009. Exposure and effects assessment of persistent organohalogen contaminants in arctic wildlife and fish. *Science of the Total Environment* doi:10.1016/j.scitotenv.2009.10.038
- 22 Roze E, Meijer L, Bakker A, Van Braeckel KNJA, Sauer PJJ, Bos AF. 2009. Prenatal exposure to organohalogens, including brominated flame retardants, influences motor, cognitive, and behavioral performance at school age. *Environmental Health Perspectives* doi:10.1289/ehp.0901015.
- 23 Herbstman JB, Sjödin A, Jones R, Kurzon M, Lederman SA, Rauh VA, Needham LL, Wang R, Perera FP. 2008. Prenatal Exposure to PBDEs and Neurodevelopment. *Environmental Health Perspectives* doi:10.1289/ehp.0901340
- 24 Collaborative on Health & the Environment, Learning and Developmental Disabilities Initiative. 2008, February 20. *Scientific Consensus Statement on Environmental Agents Associated with Neurodevelopmental Disorders*. Available: www.iceh.org/pdfs/LDDI/LDDIStatement.pdf
- 25 Costa LG, Giordano G. 2007. Developmental neurotoxicity of polybrominated diphenyl ether (PBDE) flame retardants. *Neurotoxicology* 28 (6):1047-1067

26 Rice DC, Reeve EA, Herlihy A, Zoeller RT, Thompson WD, Markowski VP. 2007. Developmental delays and locomotor activity in the C57BL6/J mouse following neonatal exposure to the fullybrominated PBDE, decabromodiphenyl ether. *Neurotoxicol Teratol* 29(4):511-20.

27 Eriksson P, Jakobsson E, Fredriksson A. 2001. Brominated flame retardants: A novel class of developmental neurotoxicants in our environment? *Environmental Health Perspectives* 109(9):903-908.

28 Kuriyama SN, Talsness CE, Grote K, Chahoud I. 2005. Developmental exposure to low-dose PBDE-99: Effects on male fertility and neurobehavior in rat offspring. *Environmental Health Perspectives* 13(2):149-154.

29 Johansson N, Viberg H, Fredriksson A, Eriksson P. 2008. Neonatal exposure to deca-brominated diphenyl ether (PBDE 209) causes dose-response changes in spontaneous behavior and cholinergic susceptibility in adult mice. *Neurotoxicology* 29(6): 911-919.

30 US EPA. 2008a. Reference Dose for PBDE-47. 2,2',4,4'-Tetrabromodiphenyl ether (BDE-47) Quickview (CASRN 5436-43-1). Integrated Risk Information System. Available: http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.showQuickView&substance_nmbr=1010 [NEED COPY]

31 US EPA. 2008b. Reference Dose for PBDE-99. 2,2',4,4',5-Pentabromodiphenyl ether (BDE-99) Quickview (CASRN 60348-60-9). Integrated Risk Information System. Available: http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.showQuickView&substance_nmbr=1008 [NEED COPY]

32 US EPA. 2008c. Reference Dose for PBDE-153. 2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE-153) Quickview (CASRN 68631-49-2). Integrated Risk Information System. Available: http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.showQuickView&substance_nmbr=1009 [NEED COPY]

33 US EPA. 2008d. Reference Dose for PBDE-209. 2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (BDE-209) Quickview (CASRN 1163-19-5). Integrated Risk Information System. Available: http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.showQuickView&substance_nmbr=0035 [NEED COPY]

34 Viberg H, Fredriksson A, Eriksson P. 2003. Neonatal exposure to polybrominated diphenyl ether (PBDE 153) disrupts spontaneous behaviour, impairs learning and memory, and decreases hippocampal cholinergic receptors in adult mice. *Toxicol Appl Pharmacol* 192(2): 95-106. [NEED COPY]

35 Viberg H, Fredriksson A, Eriksson P. 2007. Changes in spontaneous behaviour and altered response to nicotine in the adult rat, after neonatal exposure to the brominated flame retardant, decabrominated diphenyl ether (PBDE 209). *Neurotoxicology* 28(1): 136-42. [NEED COPY]

36 Siddiqi MA, Laessig RH, Reed KD. 2003. Polybrominated diphenyl ethers (PBDEs): New pollutants-old diseases. *Clinical Medicine & Research* 1(4): 281-90.

37 Herbstman JB, Sjodin A, Apelberg BJ, Witter FR, Halden RU, Patterson DG, Panny SR, Needham LL, Goldman LR. 2008. Birth delivery mode modifies the associations between prenatal polychlorinated biphenyl (PCB) and polybrominated diphenyl ether (PBDE) and neonatal thyroid hormone levels. *Environmental Health Perspectives* 116(10): 1376-1382.

38 Turyk ME, Persky VW, Imm P, Knobeloch L, Chatterton R, Anderson HA. 2008. Hormone disruption by PBDEs in adult male sport fish consumers. *Environmental Health Perspectives* 116(12). Available: www.ehponline.org/members/2008/11707/11707.html

39 Bahn AK, Mills JL, Synder PJ, Gann PH, Houten L, Bialik O, Hollmann L, Utiger RD. 1980. Hypothyroidism in workers exposed to polybrominated biphenyls. *N Engl J Med* 302:31-33. [NEED COPY]

40 Tseng LH, Li MH, Tsai SS, Lee CW, Pan MH, Yao WJ, Hsu PC. 2008. Developmental exposure to decabromodiphenyl ether (PBDE 209): Effects on thyroid hormone and hepatic enzyme activity in male mouse offspring. *Chemosphere* 70(4): 640-647.

41 National Toxicology Program (NTP). 1986. Toxicology and carcinogenesis studies of decabromodiphenyl oxide (Case No. 1163-19-5) in F344/N rats and B6C3F1 mice (feed studies). Technical Report 309. Research Triangle Park, NC: National Toxicology Program. Available: http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr309.pdf p. 47

42 Mercado-Feliciano M, Bigsby RM. 2008. The polybrominated diphenyl ether mixture DE-71 is mildly estrogenic. *Environmental Health Perspectives* 116(5): 605-611.

43 Yang W, Mu Y, Giesy JP, Zhang A, Yu H. 2009. Anti-androgen activity of polybrominated diphenyl ethers determined by comparative molecular similarity indices and molecular docking. *Chemosphere* 75:1159-1164

44 Stoker TE, Cooper RL, Lambright CS, Wilson VS, Furr J, Gray LE. 2005. In vivo and in vitro anti-androgenic effects of DE-71, a commercial polybrominated diphenyl ether (PBDE) mixture. *Toxicology and Applied Pharmacology* 207:78-88

45 Main KM, Kiviranta H, Virtanen HE, Sundqvist E, Tuomisto J, Vartiainen T, Skakkebaek NE, Toppari J. 2007. Flame retardants in placenta and breast milk and cryptorchidism in newborn boys. *Environmental Health Perspectives* 115(10): 1519-26.

46 Akutsu K, Takatori S, Nozawa S, Yoshiike M, Nakazawa H, Hayakawa K, Makino T, Iwamoto T. 2008. Polybrominated diphenyl ethers in human serum and sperm quality. *Bull Environ Contam Toxicol* 80:345-350

47 Meeker JD, Johnson PI, Camann D, Hauser R. 2009. Polybrominated diphenyl ether (PBDE) concentration in household dust are related to hormone levels in men. *Science of the Total Environment* doi: 10.1016/j.scitotenv.2009.01.030.

48 Lileenthal H, Hack A, Roth-Harer A, Grande SW, Talsness CE. 2006. Effects of developmental exposure to 2,2',4,4',5-pentabromodiphenyl ether (PBDE-99) on sex steroids, sexual development, and sexually dimorphic behavior in rats. *Environmental Health Perspectives* 114(2):194-201.

49 Schreder E. 2006. Pollution in People: A Study of Toxic Chemicals in Washingtonians. Available: www.pollutioninpeople.org

50 Chao HR, Wang SL, Lee WJ, Wang YF, Papke O. 2007. Levels of polybrominated diphenyl ethers (PBDEs) in breast milk from central Taiwan and their relation to infant birth outcome and maternal menstruation effects. *Environment International* 33(2): 239-245.

51 Canton RF, Scholten DE, Marsh G, de Jong PC, van den Berg M. 2008. Inhibition of human placental aromatase activity by hydroxylated polybrominated diphenyl ethers (OH-PBDEs). *Toxicology and Applied Pharmacology* 227(1): 68-75.

52 Hardell L, van Bavel B, Lindström G, Eriksson M, Carlberg M. 2006. In utero exposure to persistent organic pollutants in relation to testicular cancer risk. *International Journal of Andrology* 29:228-234.

53 Northwest Watch. 2004. Flame retardants in the bodies of Pacific Northwest residents: A study of toxic body burdens. Available: www.sightline.org

54 Norwegian Pollution Control Authority. 2008. Ban on Deca-BDE. Available: <http://www.sft.no/publikasjoner/2401/ta2401.pdf>

55 Government of Canada. 2008. Polybrominated diphenyl ether regulations. *Canada Gazette Part II* 142(14): 1663-1682.

56 US EPA. 2009. DecaBDE Phase-Out Initiative. Available: www.epa.gov/oppt/existingchemicals/pubs/actionplans/decadabe.html [NEED COPY]

57 State of Maine. 2007. Title 38, Chapter 16, Section 1609: Restrictions on sale and distribution of brominated flame retardants. Available: www.mainelegislature.org/legis/statutes/38/title38sec1609.html

58 State of Washington. 2008. Revised Code of Washington, chapter 70.76: Polybrominated diphenyl ethers -flame retardants. Available: <http://apps.leg.wa.gov/RCW/default.aspx?cite=70.76>



COMMONWEAL

Fact sheets on toxic chemicals for the *Mind, Disrupted* Biomonitoring Project provided by the Alaska Community Action on Toxics (www.akaction.net) and Commonweal (www.commonweal.org). For more information, please visit the *Mind, Disrupted* website at www.minddisrupted.org, or contact Pam Miller at pkmiller@akaction.net or Sharyle Patton at spatton@igc.org.

www.minddisrupted.org