

**HB**

**356**

<TARGET><BILL>HB 356</BILL><SUBJECT>HB  
356</SUBJECT><COMM>HSTA28</COMM></TARGET>



**CS FOR HOUSE BILL NO. 356(STA)**

IN THE LEGISLATURE OF THE STATE OF ALASKA

TWENTY-EIGHTH LEGISLATURE - SECOND SESSION

BY THE HOUSE STATE AFFAIRS COMMITTEE

Offered:  
Referred:

Sponsor(s): REPRESENTATIVE SEATON

**A BILL**

**FOR AN ACT ENTITLED**

1 **"An Act establishing the Advisory Committee on Wellness; and relating to the**  
2 **administration of state group health insurance policies."**

3 **BE IT ENACTED BY THE LEGISLATURE OF THE STATE OF ALASKA:**

4 \* **Section 1.** The uncodified law of the State of Alaska is amended by adding a new section  
5 to read:

6 LEGISLATIVE FINDINGS AND INTENT. (a) The legislature finds that

7 (1) individual Alaskans and the state share the burden of increasing health care  
8 and insurance costs resulting from avoidable disease in humans;

9 (2) state policy should focus more on avoiding disease and maintaining good  
10 health in the state's population;

11 (3) the state relies partly on federal one-size-fits-all human health policy  
12 recommendations, creating federal overreach into the state's administration of health policy;

13 (4) the state's human health policies should incorporate peer-reviewed  
14 scientific studies that illustrate health-related differences between the population of the state

1 and the populations of other states;

2 (5) the health policies of the state can affect the frequency and severity of  
3 disease present in the state's population;

4 (6) by establishing wellness initiatives to decrease the frequency and severity  
5 of disease present in the state's population, the state can slow the rate of annual growth in the  
6 state's cost of providing health care benefits to persons who are covered by a policy of group  
7 health insurance obtained under AS 39.30.090(a) or 39.30.091(a).

8 (b) It is the intent of the legislature that, by establishing a wellness committee and  
9 wellness initiatives that create incentives and methods to decrease the frequency and severity  
10 of disease in the state's population, the state will slow the rate of growth in the state's cost of  
11 providing health care benefits to persons who are covered by a policy of group health  
12 insurance obtained under AS 39.30.090(a) or 39.30.091(a) to a rate that does not exceed two  
13 percent annually.

14 \* **Sec. 2.** AS 39.30.090(a) is amended by adding a new paragraph to read:

15 (13) To the greatest extent legally and reasonably practicable, the  
16 Department of Administration shall work to hold the escalation of health care costs to  
17 less than two percent annually by administering policies of group health insurance  
18 obtained under this subsection in a manner that is likely to reduce the incidence of  
19 disease in the state's population and that facilitates implementation of the  
20 recommendations of the Advisory Committee on Wellness established under  
21 AS 39.30.093.

22 \* **Sec. 3.** AS 39.30.091 is amended by adding a new subsection to read:

23 (b) To the greatest extent legally and reasonably practicable, the Department  
24 of Administration shall work to hold the escalation of health care costs to less than two  
25 percent annually by administering policies of group health insurance obtained under  
26 this section in a manner that is likely to reduce the incidence of disease in the state's  
27 population and that facilitates implementation of the recommendations of the  
28 Advisory Committee on Wellness established under AS 39.30.093.

29 \* **Sec. 4.** AS 39.30 is amended by adding a new section to read:

30 **Sec. 39.30.093. Advisory Committee on Wellness.** (a) The Advisory  
31 Committee on Wellness is established in the Department of Administration. The

1 committee consists of at least seven volunteer members appointed by the  
2 commissioner of administration who serve at the pleasure of the commissioner.

3 (b) The committee shall select one of its members as chair.

4 (c) The commissioner shall appoint members who represent each region of the  
5 state to give the committee a reasonable geographic balance among regions of the state  
6 and shall appoint at least one member who is a professional provider of medical  
7 services.

8 (d) The committee shall meet at the call of the chair or at the request of a  
9 majority of its members. A member may attend a meeting by telephone or other means  
10 of telecommunication.

11 (e) At intervals of six months, the committee shall make recommendations to  
12 the commissioner of administration regarding the adoption of wellness initiatives to  
13 improve the physical and mental health of persons who are covered by a policy of  
14 group health insurance obtained under AS 39.30.090(a) or 39.30.091(a).

15 (f) Not later than six months after receiving a recommendation made under (d)  
16 of this section, the commissioner of administration shall provide to the committee a  
17 report on the commissioner's progress in advancing the recommendation and the  
18 recommendation's effect on

19 (1) the health of the population covered by a policy of group health  
20 insurance obtained under AS 39.30.090(a) or 39.30.091(a);

21 (2) the state's cost of providing health care to persons who are covered  
22 by a policy of group health insurance obtained under AS 39.30.090(a) or 39.30.091(a).

23 (g) Members of the committee serve without compensation.

## Nancy Manly

---

**From:** Nancy Manly  
**Sent:** Tuesday, April 15, 2014 9:57 AM  
**To:** LAA Legal  
**Subject:** FINAL CS TO HB356 v.C

04/15/2014

The House State Affairs Committee moved HB356 Version C this morning amended. Please draft a final CS incorporating the following amendment.

### **Conceptual Amendment#1 (Keller)**

Page 1 Line 9 after the first word "the" *delete* "state has not adequately focused" and *insert* after the word "state" **should focus more** so the sentence would read: (2) the state policy should focus more on avoiding disease...

Page 1 Line 11 after the word "relies" *insert* **in part** so the sentence would read: (3) the state relies in part on federal one-size-fits-all...

Page 1 Line 13 *delete* "have not incorporated" and *insert* **should incorporate** so the sentence would read: (4) the state's human health policies should incorporate peer-reviewed...

Nancy Manly, Chief of Staff and  
House State Affairs Committee Aide *for*  
**Representative Bob Lynn**  
**House District 23**  
907-465-2794 Fax: 907-465-4316

# Alaska State Legislature

State Capitol Room 102  
Juneau, Alaska 99801-1182  
(907) 465-2689  
Fax: (907) 465-3472  
1-800-665-2689



270 W Pioneer Ave. Suite B  
Homer, Alaska 99603  
(907) 235-2921  
Fax: (907) 235-4008  
1-800-665-2689

## REPRESENTATIVE PAUL SEATON HOUSE DISTRICT 30

### Sponsor Statement

#### HB 356

HB 356 requires the Commissioner of the Department of Administration to implement procedures for decreasing the incidence of disease in Alaska in order to hold the inflation of healthcare costs of active and retired Alaska state employees to 2% per year.

According to the Institute of Social and Economic Research, total health care spending in Alaska topped \$7.5 billion in Alaska in 2010, with state government employers paying over \$400 million. A major component of our \$12 billion unfunded pension liability is retiree healthcare costs. HB 356 requires the Commissioner of Administration to put in place programs that will decrease the incidence of disease in State of Alaska employees, both current and retired, in order to hold the inflation of costs to 2% per year.

This bill focuses on preventing the incidence of disease as opposed to treatment of disease. Prevention of disease is the policy approach unanimously requested of the Governor by the legislature through HCR 5 in 2011. This is an area of healthcare where the most economic impact can be achieved. For instance, recent studies show that an action as simple as taking a daily supplement of 5,000 IU of vitamin D can dramatically reduce the risk of heart disease, diabetes, cancer, autism, gingivitis, and many other conditions.

HB 354 creates an Advisory Committee on Wellness which is charged with making recommendations to the Commissioner of Administration on ways to decrease the incidence of disease in Alaska. HB 365 will enforce a paradigm shift for the Department of Administration and the Department of Health and Social Services. It will require the agencies to implement policies to keep Alaska state employees healthy by preventing disease, rather than the common, reactive policy of waiting until people get sick and then treating them.

# Fiscal Note

State of Alaska  
2014 Legislative Session

Bill Version: HB 356  
Fiscal Note Number: \_\_\_\_\_  
( ) Publish Date: \_\_\_\_\_

Identifier: HB356-DOA-DRB-03-14-14  
Title: ADVISORY COMMITTEE ON WELLNESS  
Sponsor: SEATON  
Requester: House Health and Social Services

Department: Department of Administration  
Appropriation: Centralized Administrative Services  
Allocation: Retirement and Benefits  
OMB Component Number: 64

**Expenditures/Revenues**

Note: Amounts do not include inflation unless otherwise noted below. (Thousands of Dollars)

	FY2015	Included in	Out-Year Cost Estimates				
	Appropriation Requested	Governor's FY2015 Request	FY 2016	FY 2017	FY 2018	FY 2019	FY 2020
<b>OPERATING EXPENDITURES</b>	<b>FY 2015</b>	<b>FY 2015</b>	<b>FY 2016</b>	<b>FY 2017</b>	<b>FY 2018</b>	<b>FY 2019</b>	<b>FY 2020</b>
Personal Services							
Travel							
Services							
Commodities							
Capital Outlay							
Grants & Benefits							
Miscellaneous							
<b>Total Operating</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>

**Fund Source (Operating Only)**

None							
<b>Total</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>

**Positions**

Full-time							
Part-time							
Temporary							

**Change in Revenues**

--	--	--	--	--	--	--	--

**Estimated SUPPLEMENTAL (FY2014) cost:** 0.0 *(separate supplemental appropriation required)*  
*(discuss reasons and fund source(s) in analysis section)*

**Estimated CAPITAL (FY2015) cost:** 0.0 *(separate capital appropriation required)*  
*(discuss reasons and fund source(s) in analysis section)*

**ASSOCIATED REGULATIONS**

Does the bill direct, or will the bill result in, regulation changes adopted by your agency? No  
If yes, by what date are the regulations to be adopted, amended or repealed? N/A

**Why this fiscal note differs from previous version:**

Not applicable, initial version.

Prepared By: Mike Barnhill, Deputy Commissioner Phone: (907)465-2200  
Division: Department of Administration Date: 03/14/2014 08:48 AM  
Approved By: Curtis Thayer, Commissioner Date: 03/14/14  
Agency: Department of Administration

## FISCAL NOTE ANALYSIS

STATE OF ALASKA  
2014 LEGISLATIVE SESSION

BILL NO. HB 356

### Analysis

HB 356 establishes an advisory committee on wellness in the Department of Administration and requires the Department to focus on wellness in an effort to limit the annual increase in health care costs for State of Alaska Employees to less than 2%.

The wellness committee would consist of at least seven volunteer members appointed by the Commissioner of the Department of Administration. The committee would make recommendations to the Commissioner of Administration regarding to adoption of wellness initiatives to improve the health of State of Alaska employees.

HB 356 also requires, to the extent legally and reasonably practicable, the Department of Administration shall work to hold the escalation of health care cost growth to less than 2% for health insurance plans administered by the Department.

The costs associated with the creation of the committee and coordinating committee meetings would be minimal and absorbed in the cost of normal business for the Department. The costs of specific initiatives recommended by the advisory committee on wellness, some of which could be substantial, cannot be estimated at this time. Therefore, the Department submits a zero fiscal note.

# ALASKA HEALTHCARE COSTS


# Alaska's Health-Care Bill: \$7.5 Billion and Climbing

By Mark A. Foster and Scott Goldsmith

UA Research Summary No. 18 • August 2011

Institute of Social and Economic Research • University of Alaska Anchorage

**Figure 1. Who Pays for Health Care in Alaska?**  
(2010 Spending: \$7.5 Billion)



<sup>a</sup>Insufficient data to break out categories

<sup>b</sup>The federal and state governments share the cost of Medicaid.

Source: Authors' estimates

Health-care spending for Alaskans reached about \$7.5 billion in 2010. For comparison, that's close to half the wellhead value of all the oil produced in Alaska that year. It's also roughly equal to half the wages Alaskans collected in 2010.

The state's health-care spending has been rising fast, tripling since 1990 and jumping 40% just between 2005 and 2010—and at current trends it could double by 2020, reaching more than \$14 billion.

Here we report on who's paying the bills, what we're buying, what's contributing to the growth, and other aspects of health-care spending. We conclude with a discussion of how Alaska could get better value for its health-care dollars.

• **Who pays the bills?** Individual Alaskans directly pay about 20%, state and federal programs around 40%, and private and government employers another 40% (Figure 1 and page 2).

• **What's the biggest cost?** Medicaid is the largest single expense, making up nearly 18% of all Alaska health-care spending. But that's down from 20% of total spending in 2005. Why? Because spending for Medicaid didn't grow as fast as other kinds of spending (page 3).

• **Are costs shifting?** Every category of spending increased since 2005—but because spending by individuals and private employers increased faster, their shares of total spending increased (page 4).

• **What are we buying?** Hospitals and doctors account for nearly 60% of total spending—but the next largest cost is the 10% that goes for administering private and government health insurance (page 4).

• **What's driving spending?** Over the past 50 years, technology, income growth, medical-price inflation, changing insurance coverage, and a growing, aging population have driven health-care spending (page 5).

• **How many Alaskans are uninsured?** The answer varies depending on how "uninsured" is measured and when. But recent estimates say about 18% of adults and 9% of children are uninsured. Based on 2010 census figures, that would be about 17,000 children and 94,000 adults (page 6).

• **How many Alaska businesses offer health insurance?** More than 90% of large firms offer insurance, compared with just 30% of small businesses—and that's down from 35% in 2003 (page 7).

• **Are prices higher in Alaska?** Yes. But Alaska's isolation, small markets, and other factors contribute to those higher prices—a day in the hospital costs on average 50% more than in the U.S. as a whole, and costs for common procedures are roughly 35% higher (page 8).

• **How is spending distributed?** Just 10% of Americans are responsible for two-thirds of all health-care spending in an average year (page 9).

• **What about the future?** Expanded insurance coverage; an aging population; and continued growth in technology, incomes, and medical prices will keep driving growth in health-care spending in the coming years. Controlling that growth will be an ongoing challenge (page 11).

### HOW HAVE PATTERNS OF SPENDING CHANGED?

Every category of health-care spending increased between 2005 and 2010, but the shares of spending shifted slightly among the various payers. We don't have enough information to say exactly what caused this shift—but several things likely contributed, as we describe below.

- *Individuals paid 20% of Alaska's health-care bills in 2010, up from 19% in 2005. As costs of health-care benefits increased rapidly, employers shifted more of those costs to employees (see page 7). Also, prices for policies individuals buy directly increased significantly.*

- *Private employers' share of spending increased from 17% to 18%. That increase was in part because private industry added nearly four times more jobs than governments did since 2005—and at least some of that bigger base of employees had health-care coverage.*

- *Government employers' share of spending was about the same, at 22%.*

- *Government health programs accounted for a somewhat smaller share of spending, down from about 41% to 39%.*

The federal and state governments have attempted to hold down growth in costs of health programs—but federal programs alone continue to make up nearly a third of all Alaska's health-care spending. Local government spending for health programs remains small, relative to that of the state and federal governments, and the increase in local spending was smaller as well.

### WHAT DO HEALTH-CARE DOLLARS BUY?


Alaska's \$7.5 billion health-care bill includes everything from visits to doctors and dentists to prescriptions and nursing-home care.<sup>5</sup> Figure 7 summarizes what Alaska's health-care dollars bought in 2010.

- *Hospital care was the largest expense, followed closely by payments for doctors and related clinical services—together they accounted for about 60% of Alaska health-care spending in 2010.*

- *Administering private and public insurance plans cost one of every ten dollars spent for Alaska health care in 2010. That's more than spending for prescriptions and medical equipment, and nearly twice the spending for dentists.*

- *Spending for nursing homes and home-health care made up only about 3% of total spending, even though spending for home health care has increased rapidly in the past decade. Much of this care is paid for under Medicaid.*

**Figure 6. Changes in Who Pays for Alaska Health-Care, 2005-2010**




### How About Health-Care Jobs?

This summary looks at health care from the perspective of spending for care—but it's important to remember that the spending also supports jobs for Alaskans. As the Alaska Department of Labor and Workforce Development reports in its August 2011 *Alaska Economic Trends*:

- Health-care spending directly supports 31,800 jobs in Alaska. That's one in ten of all wage and salary jobs—in hospitals, offices of doctors and other providers, nursing homes, and many other places.
- Many additional jobs related to health care—in government agencies, and among the self-employed—aren't included in that total.
- Alaska employment in health care has been increasing at an annual rate of 4.3% for the past decade.

**Figure 7. What Do Alaska's Health-Care Dollars Buy?**  
(2010 Spending: \$7.5 Billion)



\*Other personal and professional care and public health activities.

Source: Mark A. Foster and Associates estimates, based on Centers for Medicare and Medicaid Services, National Health Expenditure accounts

## WHO PAYS THE BILLS?

Individuals, private employers, and governments share the direct costs of health care in Alaska (Figure 1 and Table 1).

Individual Alaskans spent about \$1.5 billion for health care in 2010—20% of total spending.

- *Alaskans with employer-based insurance—both private and government—paid about \$640 million for premiums, and those with individual policies spent \$350 million.*

- *Out-of-pocket costs for Alaskans totaled about \$545 million in 2010. That includes deductibles and co-pays—the part of medical bills insurance doesn't pay. It also includes costs for services not covered by insurance, and money that uninsured Alaskans spent for medical bills.*

Private employers spent about \$1.4 billion—18% of total spending.

- *Alaska businesses spent around \$835 million to self-insure in 2010. They set aside money to pay medical bills themselves, rather than pay insurance premiums. They're betting that the medical bills will be less than the premiums they would have paid—and that their reserves will be enough to cover annual variation in claims. Many self-insured firms carry "stop loss" insurance, to protect them against very large claims. At first only large firms self-insured, but as insurance costs climbed, smaller businesses have also begun self-insuring.*

- *Businesses spent about \$400 million for insurance premiums in 2010. That's only about half what businesses spent to self-insure, showing how widespread the practice of self-insuring is.*

- *Medical bills of employees injured at work cost businesses about \$150 million in 2010. State law requires employers to pay for such injuries.*

Government employers spent \$1.6 billion for health benefits in 2010.

- *Local government employers—including school districts—spent about \$630 million, the federal government nearly \$590 million, and the state \$410 million.*

- *Like businesses, many public employers self-insure, rather than pay insurance premiums—but we don't have enough data to separate out those costs. The federal government also pays medical costs for active-duty and retired military personnel and veterans.*

Governments spent nearly \$3 billion for health programs in 2010.

- *Medicaid spending was nearly \$1.3 billion in 2010—\$871 million in federal money and \$409 million in state money. Medicaid is a federal program, but the state administers it and shares the costs (see page 3).*

- *Medicare spending was \$733 million in 2010, accounting for nearly 10% of all health-care spending. Medicare is a federal program for people 65 and older and those with certain disabilities. Medicare spending is expected to grow rapidly in the next decade, as older Alaskans make up an ever-growing share of the population (see page 5).*

- *The federal government spent close to \$650 million for other health programs in 2010, including the Indian Health Service, which provides medical care for Alaska Natives, and the Veterans Administration, which provides care for military veterans. Spending for these programs depends somewhat on enrollment, but it's also constrained by Congressional appropriations.*

- *Besides its share of Medicaid, the state government spent about \$260 million for a variety of other programs in 2010, including grants to local governments, the state-operated Pioneer Homes for older Alaskans, and the Alaska Psychiatric Institute.*

**Table 1. Health-Care Spending in Alaska, 2010**  
(Total Spending: \$7.5 Billion)

<b>Individuals</b>	<b>\$1,529 million</b>
Employee premiums	\$637
Out-of-pocket costs	\$544
Individual policies	\$348
<b>Private Employers*</b>	<b>\$1,384 million</b>
Insurance premiums	\$395
Self-insurance costs	\$836
Workers' compensation medical	\$153
<b>Government Employers*</b>	<b>\$1,625 million</b>
Federal	\$586
State	\$408
Local	\$631
<b>Federal Health Programs</b>	<b>\$2,250 million</b>
Medicare	\$733
Medicaid	\$871
IHS, VA, Community Health Centers, public health, K-12 health	\$646
<b>State Health Programs</b>	<b>\$670 million</b>
Medicaid	\$409
Local grants, API, Pioneer Homes, K-12 health, WAMI, Department of Corrections	\$261
<b>Local Health Programs</b>	<b>\$45 million</b>
Hospital and health program support	\$40
Other local	\$5

\*Includes coverage for current and retired employees.

Source: Authors' estimates. See page 12 for a description of what's included in health-care costs.

- *Local health programs are much smaller, at around \$45 million in 2010, largely support for hospitals and health programs.*

And finally, keep in mind that even though governments and businesses pay most of the *direct* costs of health care, individual Alaskans and other Americans *indirectly* pay all the costs of health care—because they buy goods and services, own businesses, and pay taxes.

ALASKA ACTIVE AND RETIREE HEALTH PLAN DATA

4th Quarter Report 2013

ACTIVE PLAN

**Aggregate Risk Profile**

Member Information			
Member Count	17338	Avg Forecasted Cost	\$6.670
Avg Age	35	Avg Total Cost	\$6.774
Percent Female	51%	Avg Forecasted Risk Index	1.09
Avg Months Enrolled	11	%/w Acute Impact Score >= 95	1.06%
		%/w Chronic Impact Score >= 95	5.38%
		%/w Motivation Rank >= 95	4.83%

Aggregate Risk Summary					
Risk Drivers	# Members	Avg Risk Contribution		Contribution to Forecast	Risk Contribution
Demographics	17338	SENIORS SKIN, FRACTURES, FALLS	\$345	\$5,987,784	5.18% X
Acute Respiratory Disorders	2880	TUBERCULOSIS	\$1,129	\$3,251,578	2.81% X
Arrhythmia Disorders	220		\$3,382	\$744,142	0.64%
CHF Conditions	673	CHRONIC HEART FAILURE	\$3,181	\$2,140,732	1.85% X
Cerebral Vascular Disorder	247		\$4,199	\$1,037,177	0.90%
Chronic Respiratory Disorders	1728	UPPER RESPIRATORY TRACT	\$2,318	\$4,005,542	3.46% X
Coronary Artery Related Conditions	1655	CORONARY HEART DISEASE	\$2,958	\$4,895,538	4.23% X
Dermatological Disorder	3075		\$1,272	\$3,911,805	3.38%
Diabetic Disorders	882	TYPE 1 AND TYPE 2	\$5,932	\$5,231,998	4.52% X
Female Reproductive Conditions	341	PRETERM BIRTHS	\$2,510	\$856,045	0.74% K
Gastrointestinal Disorders	2351	COLORECTAL CANCER	\$2,011	\$4,728,854	4.09% X
Heart Related Conditions	180		\$5,628	\$1,013,046	0.88%
Hypertension	1527	BLOOD PRESSURE	\$1,983	\$3,028,315	2.62% X
Hypotensive Drugs	1784		\$2,207	\$3,937,924	3.41%
Major Infection Related Conditions	2950	MRSA	\$2,023	\$5,968,497	5.16% X
Metabolic Conditions	3077	FIBROMYALGIA	\$2,680	\$8,247,421	7.13% X
Minor Infection Related Conditions	3704		\$1,340	\$4,965,037	4.29%
Miscellaneous Conditions	4750		\$2,303	\$10,940,696	9.46%
Musculo-skeletal Disorders	5173	INFANT MUSCLE, SENIOR FALLS	\$2,206	\$11,409,047	9.87% X
Myocardial Infarction Related Conditions	271		\$5,315	\$1,440,328	1.25%
Neonatal Issues	255	AUTISM, HEART PROGRAMMING	\$935	\$238,299	0.21% X
Neoplastic Related Conditions	638		\$4,332	\$2,763,900	2.39%
Neurological Disorder	3770	ALZHEIMER'S	\$1,435	\$5,409,047	4.68% X
Non-specific condition	5561		\$140	\$780,877	0.68%
Pneumonia	243	UPPER RESPIRATORY TRACT	\$3,822	\$928,744	0.80% X
Psychological Disorder	2688	S.A.D. AND DEPRESSION	\$2,771	\$7,447,883	6.44% X
Renal Disorders	309		\$15,145	\$4,679,794	4.05%
Trauma Related Condition	1822	TRAUMATIC BRAIN INJURY	\$1,455	\$2,651,019	2.29% X
Urinary Disorders	1381		\$2,170	\$2,996,360	2.59%

% total diseases directly related to Vitamin D status = 66.08 %

# RETIREE PLAN



4th Quarter Report 2013

## Aggregate Risk Profile

Member Information			
Member Count	65376	Avg Forecasted Cost	\$15,666
Avg Age	63	Avg Total Cost	\$17,726
Percent Female	54%	Avg Forecasted Risk Index	2.56
Avg Months Enrolled	12	%/w Acute Impact Score >= 95	5.47%
		%/w Chronic Impact Score >= 95	16.63%
		%/w Motivation Rank >= 95	9.70%

Aggregate Risk Summary				
Risk Drivers	# Members	Avg Risk Contribution	Contribution to Forecast	Risk Contribution
Demographics	65376	SENIORS SKIN, FRACTURES, FALLS \$454	\$29,702,943	2.90% X
Acute Respiratory Disorders	9520	TUBERCULOSIS \$1,669	\$15,893,121	1.55% X
Arrhythmia Disorders	5170	\$2,860	\$14,786,038	1.44%
CHF Conditions	10658	CHRONIC HEART FAILURE \$2,758	\$29,389,999	2.87% X
Cerebral Vascular Disorder	5021	\$3,726	\$18,710,595	1.83%
Chronic Respiratory Disorders	11241	UPPER RESPIRATORY TRACT \$3,093	\$34,763,411	3.39% X
Coronary Artery Related Conditions	24057	CORONARY HEART DISEASE \$2,900	\$69,776,210	6.81% X
Dermatological Disorder	15979	\$1,958	\$31,281,265	3.05%
Diabetic Disorders	10689	TYPE 1 AND TYPE 2 \$5,966	\$63,771,119	6.23% X
Female Reproductive Conditions	103	PRETERM BIRTHS \$1,489	\$153,413	0.01% X
Gastrointestinal Disorders	18753	COLORECTAL CANCER \$2,146	\$40,246,314	3.93% X
Heart Related Conditions	4346	\$4,374	\$19,007,254	1.86%
Hypertension	21394	BLOOD PRESSURE \$1,623	\$34,713,887	3.39% X
Hypotensive Drugs	21282	\$1,869	\$39,772,355	3.88%
Major Infection Related Conditions	13879	MRSA \$3,007	\$41,731,745	4.07% X
Metabolic Conditions	28763	FIBROMYALGIA \$2,545	\$73,213,862	7.15% X
Minor Infection Related Conditions	14339	\$1,726	\$24,754,410	2.42%
Miscellaneous Conditions	32888	\$2,762	\$90,833,634	8.87%
Musculo-skeletal Disorders	32886	INFANT MUSCLE, SENIOR FALLS \$2,498	\$82,154,255	8.02% X
Myocardial Infarction Related Conditions	5796	\$3,424	\$19,844,208	1.94%
Neonatal Issues	63	AUTISM, HEART PROGRAMMING \$1,504	\$94,751	0.01% X
Neoplastic Related Conditions	7447	\$3,928	\$29,253,071	2.86%
Neurological Disorder	27775	ALZHEIMER'S \$1,561	\$43,366,164	4.23% X
Non-specific condition	12687	\$113	\$1,429,752	0.14%
Pneumonia	1619	UPPER RESPIRATORY TRACT \$5,098	\$8,254,358	0.81% X
Psychological Disorder	16031	S.A.D. AND DEPRESSION \$2,659	\$42,623,390	4.16% X
Renal Disorders	4750	\$16,668	\$79,174,484	7.73%
Trauma Related Condition	8749	TRAUMATIC BRAIN INJURY \$1,904	\$16,660,268	1.63% X
Urinary Disorders	11515	\$2,504	\$28,836,425	2.82%

% total diseases directly related to Vitamin D status = 61.16%

AlaskaCare Retiree Plan - 4th Quarter Report 2013

Notes in red by office of Rep. Seaton

# DIABETES COST



# LEGISLATIVE RESEARCH SERVICES

Alaska State Legislature  
Division of Legal and Research Services  
State Capitol, Juneau, AK 99801

(907) 465-3991 phone  
(907) 465-3908 fax  
research@legis.state.ak.us

## Research Brief

TO: Representative Paul Seaton  
FROM: Tim Spengler, Legislative Analyst  
DATE: February 28, 2014  
RE: Average Annual Cost Estimates Related to Diabetes and Cancer  
*LRS Report 14.251*

---

*You asked for estimates on the average annual medical costs for individuals with diabetes and cancer. You requested estimates that consider a number of factors including doctor visits, equipment, and expected procedures, if available.<sup>1</sup>*

---

### *Diabetes Cost Estimates*

According to a major research study released in March 2013 commissioned by the American Diabetes Association (ADA), the estimated total costs of diagnosed diabetes nationwide have risen by 41 percent from 2007 to 2012.<sup>2</sup> Nationwide, costs associated with diabetes have increased from \$174 to \$245 billion during this time period. Most of these costs pertain to medical expenses although a portion relates to reduced productivity of diagnosed individuals. This in-depth study addresses the increased financial burden, health resources used, and lost productivity associated with diabetes.<sup>3</sup>

As for annual costs, the study concludes that people with diagnosed diabetes incur average medical expenditures directly attributable to the condition of around **\$7,900**. The largest medical expenditures are

- hospital inpatient care (43 percent of total medical cost);
- prescription medications (18 percent);
- anti-diabetic agents and diabetic supplies (12 percent);
- physician office visits (9 percent); and
- nursing/residential facility stays (8 percent).

Indirect costs of the disease pertain to the reduced productivity of those with diabetes. Such costs include increased absenteeism, reduced productivity in the workplace, inability to work as a result of disease-related disability, and lost productive capacity due to early mortality. Such costs exist, but were not calculated on a per person average.

The ADA-commissioned study also relates that people with diagnosed diabetes, on the average, have medical expenditures approximately 2.3 times higher than what expenditures would be in the absence of diabetes. Additionally, more than one in ten health care dollars spent in the United States is spent directly on diabetes and its related complications.

---

<sup>1</sup> You were also interested in the prevalence of diabetes and cancer among active and retired State of Alaska employees. We looked, but found no such data.

<sup>2</sup> You were particularly interested in costs for type 2 diabetes. While the study does not disaggregate by type, around 95 percent of diabetes cases are of the type 2 variety.

<sup>3</sup> The American Diabetes Association-commissioned study, "Economic Costs of Diabetes in the U.S. in 2012," can be accessed at <http://care.diabetesjournals.org/content/36/4/1033.full>.

According to the ADA, in 2012 there were around 22.3 million people—about seven percent of the U.S. population—with diagnosed diabetes. As many as seven million more people, by some estimates, likely have the disease but are, as of yet, unaware of it. Should current trends continue, by 2050, up to one in three American may have diabetes.

### *Cancer Cost Estimates*

We identified the cancer cost estimates in this section from a 2013 study funded by the Centers for Disease Control (CDC) and published by American Cancer Society. The study's findings are presented in a June 2013 original article entitled "State-Level Cancer Treatment Costs," which we include as Attachment A.<sup>4</sup> According to the authors it is the first time state-level estimates of cancer treatment costs have been published.

The study, which looked at cancer care costs across the nation during 2004 to 2008, concludes that expenditures for cancer treatment were substantial in all states and accounted for a sizable fraction of medical expenditures for all payers: Medicare, Medicaid, and private insurance. The high financial costs that cancer imposes on society underscore the importance of preventing and controlling cancer as one approach to managing state-level costs, according to the article. This is in addition to, of course, the terrible human costs that the disease causes.

The estimated average annual cancer cost per person in Alaska during 2004 to 2008 was right around \$10,000 a year.<sup>5</sup> This is slightly less than the \$11,100 average for all states. Treatment costs were highest in Michigan at around \$12,600 per year, while Arizona and California were the least expensive at around \$9,600. The study did not disaggregate costs by particular types of cancer.<sup>6</sup> The article includes a great deal of additional information that you may find of interest. For example, Table 1 estimates the average annual cancer prevalence rates for each state. Alaska's rate for all residents was 3.3 percent compared to the median national average of 4.2 percent.

Another document that you may find illuminating is the American Cancer Society's "Cancer Facts and Figures, 2014." The document estimates that in 2014 about 1,665,540 new cancer cases are expected to be diagnosed across the country. Of these diagnoses, it is estimated that 3,750 will occur in Alaska.<sup>7</sup> It also disaggregates the estimated cancers by type; in Alaska, the most commonly diagnosed cancers are predicted to be prostate, breast, lung, and colon in that order.

Finally, you may wish to peruse the CDC's "Cancer Rates by States" (<http://www.cdc.gov/cancer/dcpc/data/state.htm>). The site includes both incident and death rates for cancer disaggregated by state and type of cancer. The prevalence rates in Alaska appear to be generally on the middle or lower end of the nationwide spectrum.

We hope this is helpful. If you have questions or need additional information, please let us know.

---

<sup>4</sup> An "original" research article is a detailed account of research activity written by the scientists who did the research—not by someone else who is reporting on the research; it is considered a primary resource.

<sup>5</sup> Considering inflation, \$10,000 in 2008 would be equivalent to around \$11,000 in 2014.

<sup>6</sup> Another resource is the CDC's "cost calculator" for various chronic diseases, including cancer. The calculator must be downloaded but worked well for us. The calculator estimates the cost per person to treat cancer in Alaska to be nearly \$10,000, the same cost as "State-Level Cancer Treatment Costs," which the CDC was also involved with.

<sup>7</sup> This document can be accessed at <http://www.cancer.org/acs/groups/content/@research/documents/document/acspc-041770.pdf>. Information on rates by state can be found on pages five through eight.

**Diabetes Incidence:  
Comparing NHANES and D\*action (18+ years)**

In a comparison of data from the National Health and Nutrition Examination Survey (NHANES), 2005-2006, and GrassrootsHealth D\*action participants in the United States, we found the following:


**Incidence of Diabetes:**

NHANES: 8.5/1,000 person-years  
D\*action: 0.9/1,000 person-years

A full 90% reduction in incidence - before adjusting for co-factors.

NHANES blood level 21 ng/ml

D\*Action blood level 48 ng/ml



(Both groups had a similar average BMI, within 3 points.)

Rate Ratio = 9.7 (P=0.0002)

Chart Date: 8/6/13

© 2013 GrassrootsHealth. Preliminary data, not yet published.



**GrassrootsHealth**

A Public Health Promotion Organization

[www.grassrootshealth.net](http://www.grassrootshealth.net)

# Blood 25-Hydroxy Vitamin D Levels and Incident Type 2 Diabetes

## A meta-analysis of prospective studies

1. Yiqing Song, MD, SCD<sup>1</sup>†,
2. Lu Wang, MD, PHD<sup>1</sup>,
3. Anastassios G. Pittas, MD, MS<sup>2</sup>,
4. Liana C. Del Gobbo, PHD<sup>3</sup>,
5. Cullin Zhang, MD, PHD<sup>4</sup>,
6. JoAnn E. Manson, MD, DRPH<sup>1,5</sup> and
7. Frank B. Hu, MD, PHD<sup>1,5,6</sup>

### ± Author Affiliations

1. <sup>1</sup>Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts
2. <sup>2</sup>Division of Endocrinology, Diabetes and Metabolism, Tufts Medical Center, Boston, Massachusetts
3. <sup>3</sup>Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts
4. <sup>4</sup>Epidemiology Branch, Division of Epidemiology, Statistics, and Prevention Research, Eunice Kennedy Shriver National Institute of Child Health & Human Development, National Institutes of Health, Bethesda, Maryland
5. <sup>5</sup>Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts
6. <sup>6</sup>Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts

1. Corresponding author: Yiqing Song, [ysong3@rics.bwh.harvard.edu](mailto:ysong3@rics.bwh.harvard.edu).

### Next Section

## Abstract

**OBJECTIVE** To quantitatively assess the strength and shape of the association between blood 25-hydroxy vitamin D [25(OH)D] levels and incident risk of type 2 diabetes.

**RESEARCH DESIGN AND METHODS** A systematic search of the MEDLINE and Embase databases and a hand search of references from original reports were conducted up to 31 October 2012. Prospective observational studies that assessed the association between blood levels of 25(OH)D and risk of incident type 2 diabetes were included for meta-analysis. DerSimonian and Laird's random-effects model was used. A quadratic spline regression analysis was used to examine the shape of the association with a generalized least-squares trend test performed for the dose-response relation.

**RESULTS** A total of 21 prospective studies involving 76,220 participants and 4,996 incident type 2 diabetes cases were included for meta-analysis. Comparing the highest to the lowest category of 25(OH)D levels, the summary relative risk for type 2 diabetes was 0.62 (95% CI 0.54–0.70). A spline regression model showed that higher 25(OH)D

38% Lower Risk of Type 2 with higher vitamin D

Active State Of Alaska employees, Retirees and dependents – **83,000**

Employees, Retirees and dependents minus those with Diabetes already – **71,143**

New incidences of diabetes per year – **8.5 per 1,000 per year (.0085)**

Average cost of annual medical expenditures directly attributable to diabetes – **\$7,900**

**=**

Current Diabetes Cost per year= **\$4,777,252**

Per year Savings at 90% reduction = **\$4,299,527**

**(GrassrootsHealth D\*Action study)**

Per year Savings at 38% reduction = **\$1,815,356**

**(Meta-analysis of prospective studies - Song et.al.)**

**COLORECTAL CANCER**

## Association Between Vitamin D and Risk of Colorectal Cancer: A Systematic Review of Prospective Studies

Yanlei Ma, Peng Zhang, Feng Wang, Jianjun Yang, Zhihua Liu, and Huanlong Qin

### ABSTRACT

#### Purpose

To conduct a systematic review of prospective studies assessing the association of vitamin D intake or blood levels of 25-hydroxyvitamin D [25(OH)D] with the risk of colorectal cancer using meta-analysis.

#### Methods

Relevant studies were identified by a search of MEDLINE and EMBASE databases before October 2010 with no restrictions. We included prospective studies that reported relative risk (RR) estimates with 95% CIs for the association between vitamin D intake or blood 25(OH)D levels and the risk of colorectal, colon, or rectal cancer. Approximately 1,000,000 participants from several countries were included in this analysis.

#### Results

Nine studies on vitamin D intake and nine studies on blood 25(OH)D levels were included in the meta-analysis. The pooled RRs of colorectal cancer for the highest versus lowest categories of vitamin D intake and blood 25(OH)D levels were 0.88 (95% CI, 0.80 to 0.96) and 0.67 (95% CI, 0.54 to 0.80), respectively. There was no heterogeneity among studies of vitamin D intake ( $P = .19$ ) or among studies of blood 25(OH)D levels ( $P = .96$ ). A 10 ng/mL increment in blood 25(OH)D level conferred an RR of 0.74 (95% CI, 0.63 to 0.89).

#### Conclusion

Vitamin D intake and blood 25(OH)D levels were inversely associated with the risk of colorectal cancer in this meta-analysis.

33% Lower Risk

*J Clin Oncol* 29:3775-3782. © 2011 by American Society of Clinical Oncology

All authors: The Sixth People's Hospital affiliated with Shanghai Jiao Tong University, Shanghai, People's Republic of China.

Submitted March 4, 2011; accepted June 8, 2011; published online ahead of print at www.jco.org on August 29, 2011.

Supported by Grants No. 11QA1404800 from the Shanghai Rising-Star Program, 81001089 from the National Natural Science Foundation of China, and 2009AA02Z118 from the National 863 High Technology Foundation.

Y.M. and H.Q. contributed equally to this work.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Corresponding author: Huanlong Qin, Department of Surgery, The Sixth People's Hospital affiliated with Shanghai Jiao Tong University, 600 Yishan Rd, Shanghai 200233, People's Republic of China; e-mail: hl-qin@hotmail.com.

© 2011 by American Society of Clinical Oncology

0732-183X/11/2928-3775/\$20.00

DOI: 10.1200/JCO.2011.35.7566

### INTRODUCTION

25-hydroxyvitamin D [25(OH)D] is the precursor of the physiologically active form of vitamin D. The serum level of 25(OH)D is a result of exposure of the skin to sunlight, total vitamin D intake, and other factors such as age and skin pigmentation.<sup>1-2</sup> Vitamin D has the ability to inhibit cell proliferation and increase apoptosis *in vitro*, and several tissues can locally produce the physiologically active form of vitamin D, which has anticarcinogenic properties.<sup>3-6</sup> In addition, many cell types, including colorectal epithelial cells, contain vitamin D receptors. These cells are able to convert the circulating 25(OH)D into active 1 to 25(OH)D metabolites, which in turn bind to the cells' own vitamin D receptors to produce an autocrine effect by inducing cell differentiation and inhibiting proliferation, invasiveness, angiogenesis, and metastatic potential.<sup>7</sup> Therefore, low vitamin D levels may increase the risk of colorectal cancer through the above potential mechanism. Currently, vitamin D deficiency is an impor-

tant health problem in the industrial world<sup>8-9</sup>; in the United States, 25% to 58% of adolescents and adults are deficient in vitamin D.<sup>10</sup>


The results from prospective studies that have examined the association between vitamin D intake or 25(OH)D levels in the blood and the risk of colorectal cancer have been inconsistent. The aim of this review was to evaluate the evidence from prospective studies on vitamin D intake or blood levels of 25(OH)D and the risk of colorectal cancer by summarizing it quantitatively with a meta-analysis approach.

### METHODS

#### Search Strategy

The literature search was conducted before October 2010 in the MEDLINE and EMBASE databases without restrictions and included articles ahead of publication. The following keywords were used in searching: "vitamin D or 25(OH)D" and "colorectal cancer or colon cancer or rectal cancer." Moreover, we searched

# Meta-analysis



**Figure 1.** Dose-response gradient for colorectal cancer according to serum 25(OH)D concentration, all five studies combined.<sup>1,4-7</sup> The five points are the odds ratios for each quintile of 25(OH)D based on combined data from the five studies.

Active State Of Alaska employees, Retirees and dependents – **83,000**

Incidence of Colorectal Cancer per year in AK - **43 per 100,000 (.0043)**

Average cost of annual medical expenditures directly attributable to Colon Cancer  
– **\$11,000**

=

AK State Cost for Colorectal Cancer per year **\$ 3,925,900**


**50% per year savings with vitamin D**

**\$1,962,950**

**(meta-analysis Gorham et. al.)**

# BREAST CANCER

# Meta-analysis of breast cancer risk



Dose-response gradient of risk of breast cancer according to serum 25-hydroxyvitamin D concentration, pooled analysis.

Active State Of Alaska employees, Retirees and dependents – **83,000**

**Female percentage of AK employees and retirees: 53% = 43,990**

Incidence of Breast Cancer per year in AK - **125 per 100,000 (.0125)**

Average cost of annual medical expenditures directly attributable to Breast  
Cancer – **\$11,000**

=

Per year AK State Cost for Breast Cancer: **\$ 6,048,625**

**50% reduction with vitamin D**

**Per Year Savings with vitamin D:**

**\$3,024,312**

# PRETERM BIRTHS

Select a summary



search

go

news: Fetal and

## Alaska

Find maternal and infant health data on a state level, or by county or city. Narrow your results or compare with another region.

Location: Alaska [edit](#)

Topic: Preterm by race/ethnicity [edit](#)

Format: Bar Graph [edit](#)


search

reset

view tray

slides (0)

### Preterm by race/ethnicity: Alaska, 2009-2011 Average





## GRASSROOTSHEALTH

A Public Health Promotion Organization

### Protect Our Children *NOW!*

*A community outreach program to reduce the incidence of preterm births quickly, easily, and safely by attending to solving the vitamin D deficiency epidemic through the engagement of pregnant women in a value changing program of Good Health vs 'Treating Illness'*

Approximately 1300 infants will be born prematurely in Alaska in 2014 per the March of Dimes. Fully 25-50% of these in the state, 325-650 babies and their families, could possibly have this trauma prevented with vitamin D supplementation to the pregnant mother. Premature births are closely associated with cerebral palsy, mental impairment and permanent hearing loss among other deficiencies.

The March of Dimes estimates that the cost of each premature infant is \$55,000, adding up to a total annual cost of \$72MM of which \$18-36MM could likely be saved (on an annual basis). Data from randomized trials and others works from Dr. Wagner et al. showed a potential reduction of 50% in preterm births and significant reductions in preeclampsia and gestational diabetes as well as other complications of pregnancy. The problem now is getting the results into practice quickly vs waiting the standard 15-25 years.

Solving this problem requires nothing less than *Changing Cultural Values*, from 'Early Detection' to 'Primary Prevention'; from 'Affordable Care' to 'Good Health'; from individuals 'Taking Advice' of physicians to 'Consultation' with them. In order to accomplish this, the timing is perfect to link a new, highly accepted technology (internet application) to the new value of HEALTH through the environment of 'MyOWNHealth™' which captures all the essential ingredients of change: the science, the proven recommendations for pregnancy, clear methods for setting priorities for the individual, personal feedback and rewards for performance, a process for total engagement from learning through personal reward systems. This is a personal portal for the patient.

Behind the scenes, information is tracked by the system to provide information to the providers, the insurers, the scientists about what's working, what needs changing, i.e., a complete feedback loop to perfect the process. This will improve processes as well as strengthen public health.

A full demonstration of this process, to serve as a 'seed' for an entire community, has been developed by an international non-profit public health promotion organization, GrassrootsHealth, in conjunction with the leading researcher, Dr. Carol L. Wagner of the Medical University of South Carolina as the Principal Designer/Leader. They have in place not only a vitamin D testing program for the mothers and infants, but all the pieces of the MyOWNHealth™ system: simple, interactive educational programs for participants and physicians, engaging games and reward systems, programs to track the progress of healthy behavioral changes, management feedback to provide ongoing enhancements to the process.

This community project will involve the active participation of about 500 pregnant women. With the 500 women participating, there could be 25 children saved this problem with a potential cost savings of \$1,375,000 for this group alone in the first implementation.

Next steps to explore this program would include a meeting with Carole Baggerly, director of GrassrootsHealth and Dr. Wagner to highlight the details of a project plan for the community site.

- Approximate number of births per year in Alaska = 11,000
- Assumption: 500 births per year to State of Alaska Employees, Retirees and Dependents
- For each 500 pregnancies in the Alaska insured and dependent category with vitamin D sufficiency 25 preterm births avoided
- Savings to the state by avoiding 25 preterm births = \$1,375,000




# UPPER RESPIRATORY TRACT INFECTIONS



## Upper Respiratory Tract Infections

Recently, a study was conducted with seven hundred forty-three children ages 3-15 in a Canadian Hutterite Community. **The findings of the study show that children with higher vitamin D blood levels had a 50% lower relative risk of contracting an Upper Respiratory Tract infection.** Those children at the United States national average of 21 ng/ml vitamin D levels were at a 70% greater risk of contracting respiratory infections. Illnesses such as RTI's are commonly a factor in children's absences from school. Making sure your child has sufficient vitamin D will not only increase their health, but will lead to less school absences due to illness.



*Low Serum 25 Hydroxyvitamin D level and Risk of Upper Respiratory tract infection in Children and Adolescents Science et. al. Journal of Clinical Infectious Diseases, August 2013 volume 57.*

Prepared by the office of Representative Paul Seaton

# Vitamin D<sub>3</sub> supplementation in patients with frequent respiratory tract infections: a randomised and double-blind intervention study

Peter Bergman,<sup>1,2,3</sup> Anna-Carin Norlin,<sup>2,4</sup> Susanne Hansen,<sup>2</sup> Rokeya Sultana Rekha,<sup>5</sup> Birgitta Agerberth,<sup>5</sup> Linda Björkhem-Bergman,<sup>6</sup> Lena Ekström,<sup>6</sup> Jonatan D Lindh,<sup>6</sup> Jan Andersson<sup>3</sup>

To cite: Bergman P, Norlin A-C, Hansen S, *et al*. Vitamin D<sub>3</sub> supplementation in patients with frequent respiratory tract infections: a randomised and double-blind intervention study. *BMJ Open* 2012;2:e001663. doi:10.1136/bmjopen-2012-001663

► Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-001663>).

Received 13 June 2012  
Revised 15 November 2012  
Accepted 15 November 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

PB and ACN contributed equally to this study.

For numbered affiliations see end of article.

Correspondence to Dr Peter Bergman; [peter.bergman@ki.se](mailto:peter.bergman@ki.se)

## ABSTRACT

**Background:** Low serum levels of 25-hydroxyvitamin D<sub>3</sub> are associated with an increased risk of respiratory tract infections (RTIs). Clinical trials with vitamin D<sub>3</sub> against various infections have been carried out but data are so far not conclusive. Thus, there is a need for additional randomised controlled trials of effects of vitamin D<sub>3</sub> on infections.

**Objective:** To investigate if supplementation with vitamin D<sub>3</sub> could reduce infectious symptoms and antibiotic consumption among patients with antibody deficiency or frequent RTIs.

**Design:** A double-blind randomised controlled trial.

**Setting:** Karolinska University Hospital, Huddinge.

**Participants:** 140 patients with antibody deficiency (selective IgA subclass deficiency, IgG subclass deficiency, common variable immune disorder) and patients with increased susceptibility to RTIs (>4 bacterial RTIs/year) but without immunological diagnosis.

**Intervention:** Vitamin D<sub>3</sub> (4000 IU) or placebo was given daily for 1 year.

**Primary and secondary outcome measures:** The primary endpoint was an infectious score based on five parameters: symptoms from respiratory tract, ears and sinuses, malaise and antibiotic consumption.

Secondary endpoints were serum levels of 25-hydroxyvitamin D<sub>3</sub>, microbiological findings and levels of antimicrobial peptides (LL-37, HNP1–3) in nasal fluid.

**Results:** The overall infectious score was significantly reduced for patients allocated to the vitamin D group (202 points) compared with the placebo group (249 points; adjusted relative score 0.771, 95% CI 0.604 to 0.985, p=0.04).

**Limitations:** A single study centre, small sample size and a selected group of patients. The sample size calculation was performed using p=0.02 as the significance level whereas the primary and secondary endpoints were analysed using the conventional p=0.05 as the significance level.

**Conclusions:** Supplementation with vitamin D<sub>3</sub> may reduce disease burden in patients with frequent RTIs.

## ARTICLE SUMMARY

### Article focus

- Recent evidence suggests that vitamin D<sub>3</sub> has potent extraskeletal effects, such as suppression of inflammation and strengthening of mucosal immunity by induction of antimicrobial peptides.
- Data from observational studies suggest that low levels of 25-hydroxyvitamin D<sub>3</sub> are associated with an increased risk of respiratory tract infections.
- Results from a limited number of randomised controlled trials on the protective role of vitamin D<sub>3</sub> against respiratory tract infections are inconclusive and thus additional studies are warranted.

### Key messages

- Therefore we designed and carried out a randomised controlled trial where a large dose (4000 IU) of vitamin D<sub>3</sub> was given to patients with an increased susceptibility to infections for 1 year.
- The main conclusion is that vitamin D<sub>3</sub> supplementation reduces symptoms and antibiotic consumption among patients with an increased frequency of respiratory tract infections. Thus, vitamin D<sub>3</sub> supplementation may be an alternative strategy to reduce antibiotic use among patients with recurrent respiratory tract infections.

### Strengths and limitations of this study


- A high daily dose of vitamin D<sub>3</sub> was used, the study time was a full year covering all seasons and patients with an increased frequency of respiratory tract infections were studied.
- A single study centre, small sample size (n=140) and a selected group of patients.

## INTRODUCTION

Vitamin D was discovered when it was noted that rachitic children were improved by exposure to sunlight.<sup>1</sup> It was later shown by Holick *et al*<sup>2</sup> that vitamin D<sub>3</sub> is synthesised in the skin under the influence of ultraviolet light. Vitamin D<sub>3</sub> is further hydroxylated in the liver

# INFLAMMATION

## Inflammatory gum disease



Hiremath VP, Rao CB, Naik V, Prasad KVV. Anti-inflammatory Effect of Vitamin D on Gingivitis: A Dose-Response Randomised Control Trial. Oral Health Prev. Dent. 2013; 11(1):61-69. (notes by Seaton)

# MUSCLE FUNCTION



JAMDA

journal homepage: [www.jamda.com](http://www.jamda.com)

## Original Study

## Vitamin D and Muscle Function: Is There a Threshold in the Relation?

Hennie C.J.P. Janssen MD<sup>a,\*</sup>, Marielle H. Emmelot-Vonk MD, PhD<sup>a</sup>, Harald J.J. Verhaar MD, PhD<sup>a</sup>, Yvonne T. van der Schouw Prof<sup>b</sup><sup>a</sup> Department of Geriatrics, University Medical Center Utrecht, Utrecht, The Netherlands<sup>b</sup> Julius Center for Health Sciences and Primary Care, Utrecht University, Utrecht, The Netherlands

## A B S T R A C T

**Keywords:**  
Vitamin D  
muscle function  
performance  
elderly

**Objectives:** First, to determine the association between serum 25 hydroxyvitamin D (25OHD) concentration and muscle mass, strength, and performance. Second, to explore if there is a threshold in the association.

**Design:** Cross-sectional, single-center study.

**Setting:** The central part of the Netherlands (52° Northern latitude).

**Participants:** A total of 802 independently living men and postmenopausal women 40 to 80 years of age.

**Measurements:** Health-related and lifestyle factors, including physical activity, 25OHD concentration, lean mass, handgrip strength, knee extension strength, and physical performance were determined.

**Results:** Overall, higher 25OHD level was significantly associated with higher lean mass (22.6 g per nmol/L, 95% CI 7.3–37.9), handgrip strength (0.020 kg per nmol/L, 95% CI 0.001–0.038), and physical performance (0.006 points per nmol/L, 95% CI 0.001–0.012), after adjustment for various confounders. This association was most pronounced below a 25OHD level of 60 nmol/L, with lean mass increase 79.6 g per nmol/L (95% CI 40.8–118.4,  $P < .01$ ), handgrip strength 0.09 kg per nmol/L (95% CI 0.045–0.141,  $P < .01$ ), and physical performance 0.02 points per nmol/L (95% CI 0.005–0.032,  $P < .01$ ), and these significant associations attenuated to null above this threshold.

**Conclusion:** In middle aged men and (postmenopausal) women, a higher 25OHD level was significantly associated with higher lean mass, muscle strength, and performance. These associations were most pronounced below 60 nmol/L and absent above 60 nmol/L, indicating a ceiling effect.

Copyright © 2013 - American Medical Directors Association, Inc.

Evidence on the diverse actions of vitamin D has been growing exponentially over the past decades. In addition to its well-known role in bone metabolism, vitamin D involvement has been reported in autoimmune disease, reproductive function, malignancy, mood disorder, the metabolic syndrome, and, recently, even in sleep disorders.<sup>1,2</sup> One of the major fields of investigation regarding vitamin D has been in the prevention of falls and fractures in the elderly.<sup>3–12</sup>

Annually, at least 30% of independently living older people experience a fall,<sup>13</sup> with a quarter of those who fall having serious

injury requiring medical attention and about 6% experiencing a fracture.<sup>14</sup> This has profound implications on quality of life,<sup>15</sup> and in a US population-based survey, no fewer than 50% of independently living patients with a fall-related injury admitted to hospital were discharged to a nursing home.<sup>16</sup>

Several mechanisms have been postulated for a causal role of vitamin D deficiency in falls and fractures. First, vitamin D deficiency may impair bone metabolism and thereby increase proneness to fracture, should a fall occur.<sup>17</sup> Second, vitamin D deficiency may cause muscle weakness,<sup>5</sup> and, finally, it may exert a negative effect on postural stability and body sway.<sup>18,19</sup>

In severe vitamin D deficiency, vitamin D supplementation, with or without calcium, improved muscle function and balance.<sup>18,20–22</sup> However, evidence from meta-analyses on falls and fractures is still inconclusive.<sup>23–26</sup> This may be partly because it is unclear what serum hydroxyvitamin D concentration constitutes adequate vitamin D status with regard to bone health and extraskelatal vitamin D actions. International guidelines advise a serum hydroxyvitamin D concentration of 50 nmol/L<sup>27</sup> and 75 nmol/L as adequate.<sup>28</sup> However, in a meta-analysis on fall prevention in older people, a minimum serum hydroxyvitamin D concentration of 60 nmol/L was necessary

This study was funded by ZON-MW (grant no. 2100.0011) and the International Health Foundation, Geneva, Switzerland. ZON-MW and The International Health Foundation neither controlled nor influenced the contents of the research or of this article, nor played any part in the decision to submit this manuscript for publication. The costs of the vitamin D measurements were partly sponsored by Roche Diagnostics Nederland b.v. Roche Diagnostics neither controlled nor influenced the contents of this research.

The authors declare no conflicts of interest.

\* Address correspondence to Hennie C.J.P. Janssen, MD, Department of Geriatrics, 805.256, University Medical Center Utrecht, PO Box 85500, 3508 GA Utrecht, The Netherlands.

E-mail address: [hcjp.janssen@gmail.com](mailto:hcjp.janssen@gmail.com) (H.C.J.P. Janssen).

## A randomized study on the effect of vitamin D<sub>3</sub> supplementation on skeletal muscle morphology and vitamin D receptor concentration in older women

Lisa Ceglia<sup>1,2</sup>, Sathit Niramitmahapanya<sup>2,3</sup>, Mauricio da S. Morais<sup>4</sup>, Donato A. Rivas<sup>4</sup>, Susan S. Harris<sup>2</sup>, Heike Bischoff-Ferrari<sup>2,5,6</sup>, Roger A. Fielding<sup>4</sup>, Bess Dawson-Hughes<sup>2</sup>

1Division of Endocrinology, Diabetes and Metabolism, Tufts Medical Center, Boston, MA; <sup>2</sup>Bone Metabolism Laboratory and <sup>4</sup>Nutrition, Exercise Physiology and Sarcopenia Laboratory, Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA; <sup>3</sup>Department of Medicine, Rajavithi Hospital, College of Medicine, Rangsit University, Bangkok, Thailand; <sup>5</sup>Centre on Aging and Mobility, University of Zurich and <sup>6</sup>City Hospital Waid; Department of Geriatrics and Aging Research, University Hospital Zurich

**Context:** Studies examining whether vitamin D supplementation increases muscle mass or muscle-specific vitamin D receptor (VDR) concentration are lacking.

**Objective:** To determine whether vitamin D<sub>3</sub> 4000 IU/d alters muscle fiber cross-sectional area (FCSA) and intramyonuclear VDR concentration over 4 months.

**Design and Setting:** Randomized, double-blind, placebo-controlled study in a single center.

**Participants:** 21 mobility-limited women (aged  $\geq 65$  years) with serum 25-hydroxyvitamin D (25OHD) levels 22.5–60 nmol/L.

**Main Outcome Measures:** Baseline and 4-month FCSA and intramyonuclear VDR were measured from *vastus lateralis* muscle cross-sections probed for muscle fiber type (I/IIa/IIx) and VDR using immunofluorescence.

**Results:** At baseline, mean ( $\pm$ SD) age was  $78 \pm 5$  years; body mass index (BMI) was  $27 \pm 5$  kg/m<sup>2</sup>; 25OHD was  $46.3 \pm 9.5$  nmol/L; and a short physical performance battery score was  $7.95 \pm 1.57$  out of 12. At 4 months, 25OHD level was  $52.5 \pm 17.1$  (placebo) vs.  $80.0 \pm 11.5$  nmol/L (VD;  $P < 0.01$ ) and change in 25OHD level was strongly associated with percent change in intramyonuclear VDR concentration independent of group ( $r = 0.87$ ,  $P < 0.001$ ). By treatment group, percent change in intramyonuclear VDR concentration was  $7.8 \pm 18.2\%$  (placebo) vs.  $29.7 \pm 11.7\%$  (VD;  $P = 0.03$ ) with a more pronounced group difference in type II vs. I fibers. Percent change in total (type I/II) FCSA was  $-7.4 \pm 18.9\%$  (placebo) vs.  $10.6 \pm 20.0\%$  (VD;  $P = 0.048$ ).

**Conclusion:** Vitamin D<sub>3</sub> supplementation increased intramyonuclear VDR concentration by 30% and increased muscle fiber size by 10% in older, mobility-limited, vitamin D-insufficient women. Further work is needed to determine whether the observed effect of vitamin D on fiber size is mediated by the VDR and to identify which signaling pathways are involved.



Low vitamin D status has been associated with reduced muscle mass, strength, and performance in older adults (1–5). Several intervention studies have reported

that vitamin D supplementation increases appendicular muscle strength and improves physical function particularly in older women with low vitamin D status (6–9).

ISSN Print 0021-972X ISSN Online 1945-7197

Printed in U.S.A.

Copyright © 2013 by The Endocrine Society

Received July 11, 2013. Accepted October 7, 2013.

Abbreviations:

## Maternal Antenatal Vitamin D Status and Offspring Muscle Development: Findings From the Southampton Women's Survey

Nicholas C. Harvey,\* Rebecca J. Moon,\* Avan Aihie Sayer, Georgia Ntani, Justin H. Davies, M. Kassim Javaid, Sian M. Robinson, Keith M. Godfrey, Hazel M. Inskip, Cyrus Cooper, and The Southampton Women's Survey Study Group

Medical Research Council Lifecourse Epidemiology Unit (N.C.H., R.J.M., A.A.S., G.N., S.M.R., K.M.G., H.M.I., C.C.), University of Southampton, Southampton SO16 6YD, United Kingdom; National Institute for Health Research Southampton Biomedical Research Centre (N.C.H., K.M.G., C.C.), University of Southampton and University Hospital Southampton National Health Service Foundation Trust, Southampton SO16 6YD, United Kingdom; Paediatric Endocrinology (R.J.M., J.H.D.), University Hospital Southampton National Health Service Foundation Trust, Southampton SO16 6YD, United Kingdom; and National Institute for Health Research Musculoskeletal Biomedical Research Unit (M.K.J., C.C.), University of Oxford, Nuffield Orthopedic Centre, Headington, Oxford OX3 7HE, United Kingdom

**Context:** Maternal 25-hydroxyvitamin D [25(OH)D] status in pregnancy has been associated with offspring bone development and adiposity. Vitamin D has also been implicated in postnatal muscle function, but little is known about a role for antenatal 25(OH)D exposure in programming muscle development.

**Objective:** We investigated the associations between maternal plasma 25(OH)D status at 34 weeks of gestation and offspring lean mass and muscle strength at 4 years of age.

**Design and Setting:** We studied a prospective UK population-based mother-offspring cohort: the Southampton Women's Survey (SWS).

**Participants:** Initially, 12 583 nonpregnant women were recruited into the SWS, of whom 3159 had singleton pregnancies; 678 mother-child pairs were included in this analysis.

**Main Outcomes Measured:** At 4 years of age, offspring assessments included hand grip strength and whole-body dual-energy x-ray absorptiometry, yielding lean mass and percent lean mass. Physical activity was assessed by 7-day accelerometry in a subset of children (n = 326).

**Results:** The maternal serum 25(OH)D concentration in pregnancy was positively associated with offspring height-adjusted hand grip strength ( $\beta = 0.10$  SD/SD,  $P = .013$ ), which persisted after adjustment for maternal confounding factors, duration of breastfeeding, and child's physical activity at 4 years ( $\beta = 0.13$  SD/SD,  $P = .014$ ). Maternal 25(OH)D was also positively associated with offspring percent lean mass ( $\beta = 0.11$  SD/SD,  $P = .006$ ), but not total lean mass ( $\beta = 0.06$  SD/SD,  $P = .15$ ). However, this association did not persist after adjustment for confounding factors ( $\beta = 0.09$  SD/SD,  $P = .11$ ).

**Conclusions:** This observational study suggests that intrauterine exposure to 25(OH)D during late pregnancy might influence offspring muscle development through an effect primarily on muscle strength rather than on muscle mass. (*J Clin Endocrinol Metab* 99: 330–337, 2014)

\*

ISSN Print 0021-972X ISSN Online 1945-7197  
Printed in U.S.A.

Copyright © 2014 by The Endocrine Society

Received August 22, 2013 Accepted October 23, 2013

First Published Online October 31, 2013

\* N.C.H. and R.J.M. contributed equally to the study.

Abbreviations: DXA, dual-energy x-ray absorptiometry; MVPA, moderate to vigorous physical activity; 25(OH)D, 25-hydroxyvitamin D; SWS, Southampton Women's Survey; VDR, vitamin D receptor.



# SENIOR FALLS AND FRACTURES



## Vitamin D project helps prevent falls and saves health costs

Published By [Live News](#) / August 8, 2013 / [No Comments](#)

Source: New Zealand Government – Press Release/Statement:

Headline: Vitamin D project helps prevent falls and saves health costs

Associate Minister of Health Jo Goodhew says MidCentral DHB's vitamin D project is a good example of how a simple intervention can improve lives and save health dollars.

In 2010 the DHB, in partnership with ACC, began encouraging health professionals to prescribe vitamin D to residents in aged care facilities. Between March 2010 and June 2012 the uptake of vitamin D by aged care residents increased from 15 to 74 per cent.

“Comparisons from before and after the start of the project show a 32 per cent reduction in aged residential care residents going to the emergency department with falls-related fractures, and a 41 per cent reduction in their hospital admissions due to these fractures.” Mrs Goodhew said.

“The benefits of preventing falls in older people cannot be overstated. Preventing falls enables older people to maintain their independence and confidence.

“Of older people who suffer a hip fracture, nearly 20 per cent will die within a year. Almost half will require long-term care and half will require help at home. Half of those who walked without help before fracturing a hip will be unable to walk without assistance in the year following the fracture.”

The vitamin D project is also estimated to have saved MidCentral DHB more than \$540,000 because of fewer people coming to the emergency department and reduced admissions to hospital. Further savings are also likely because of reduced need for clinical support, hospital pharmacy services, and physiotherapy and rehabilitation services. \*

International evidence shows that taking vitamin D significantly reduces older adults' risk of falling.

“We know older people are less likely to fall and injure themselves if they keep their muscles and bones in good condition.

Vitamin D has been shown to increase the number and size of type II muscle fibres, which play an important role in balance and mobility. Vitamin D also helps maintain bone strength,” MidCentral DHB pharmacy advisor Andrew Orange says.

The Health Quality & Safety Commission's national patient safety campaign *Open for better care* is currently focusing on falls prevention. For more information about the *Open* campaign, go to [www.open.hqsc.govt.nz](http://www.open.hqsc.govt.nz).



**CLINICAL PROTOCOL: Vitamin D Protocol in Residential Care Facilities**

**AUTHORIZATION:**  
**RESIDENTIAL CARE &  
ASSISTED LIVING PROGRAM,  
OLDER ADULT PROGRAM &  
THE CLINICAL POLICY OFFICE**

**Date Released:**  
November 8,  
2011

**PURPOSE:**

To standardize and optimize vitamin D supplementation for people living in Fraser Health (FH) residential care facilities.

**BACKGROUND**

Vitamin D supplementation has been extensively studied as a treatment to prevent both falls and fractures<sup>1-3</sup>. Vitamin D is an important nutrient involved in calcium metabolism, bone health, and muscle function, hence its direct beneficial effect on falls and fracture prevention. In addition, prospective epidemiologic studies have suggested that vitamin D may reduce the risk of cardiovascular disease and some forms of cancer, and may have positive effects on immune responses and anti-inflammatory benefits<sup>4,5</sup>.

Vitamin D deficiency can arise from limited sun exposure, impaired ability of the liver or kidneys to activate vitamin D, limited dietary intake, or poor absorption from the intestine<sup>6,7</sup>. Vitamin D deficiency can cause osteomalacia (characterized by weakness of the bone and muscle), contribute to the development of osteoporosis, immune system dysfunction, and bone pain and is also associated with an increased risk of falls and higher risk of fractures in older adults<sup>6,7</sup>.

Despite the evidence that vitamin D is an important treatment for falls and fracture prevention, as well as potential cancer and cardiovascular benefits, vitamin D supplementation is not standard in most residential facilities and the prevalence of vitamin D deficiency is high in institutionalized people<sup>8,9</sup>.

To address this gap, FH (with funding from the Canadian Institutes of Health Research), brought together a Specialist Advisory Group (see Appendix A) comprised of health professionals, researchers, decision makers, policy makers, and other relevant stakeholders for a series of meetings with the purpose of developing an evidence based, practical and sustainable vitamin D protocol for residents of residential care facilities. The main objective of the group was to develop recommendations to indicate who should and should not be placed on the vitamin D protocol, how much vitamin D is safe and effective, and what the optimum dosing frequency is with minimal impact on staff workload and cost.

According to the recommendation from the Specialist Advisory Group, the adequate safe and effective dosage for older adult residents living in residential care facilities is 20,000 IU weekly of vitamin D<sub>3</sub> (2X 10,000 IU in tablet form). Only residents with hypercalcaemia<sup>Δ</sup> and/or severe renal

\* The upper level intakes of Vitamin D set by the Institute of Medicine (4,000 IU/day) and the Endocrine Society Clinical Practice Guideline (10,000 IU/day) represent the safe boundary at the high end of the scale and should not be misunderstood as amounts people need or should strive to consume<sup>6,10</sup>.

<sup>Δ</sup> Diagnosing hypercalcaemia should not utilize serum calcium level testing to implement protocol unless physicians have significant evidence to test.

failure (GFR <20 mL/ min) or those who refuse supplementation, should be excluded from the protocol.

**DEFINITIONS**

**Vitamin D:** is a fat soluble steroid hormone derived from cholesterol. Vitamin D<sub>3</sub> (cholecalciferol) is one form that is synthesized in our skin when exposed to sunlight. Another form of vitamin D, vitamin D<sub>2</sub> (ergocalciferol) can be obtained from certain foods, supplements or by prescription.

**Vitamin D deficiency:** blood serum levels are less than 25 nmol/L

**Vitamin D insufficiency:** blood serum levels are between 25-75 nmol/L

**Vitamin D sufficiency:** blood serum levels are between 75-250 nmol/L

\*

**Vitamin D toxicity:** blood serum levels are above 375 nmol/L

**Tolerable Upper Limit:** is the highest dose a person can chronically consume without risk of adverse effects

**RELATED RESOURCES**

- I. Vitamin D: A proven D-fence against falls. 2008. Available at:  
[http://www.acc.co.nz/PRD\\_EXT\\_CSMP/groups/external\\_ip/documents/publications\\_promotion/prd\\_ctrb095324.pdf](http://www.acc.co.nz/PRD_EXT_CSMP/groups/external_ip/documents/publications_promotion/prd_ctrb095324.pdf)
- II. Stay on your feet and stay active - with a little help from Vitamin D. 2008. Available at:  
[http://www.acc.co.nz/PRD\\_EXT\\_CSMP/groups/external\\_ip/documents/publications\\_promotion/prd\\_ctrb095323.pdf](http://www.acc.co.nz/PRD_EXT_CSMP/groups/external_ip/documents/publications_promotion/prd_ctrb095323.pdf)
- III. Standing up to falls. 2009. Available at:  
[http://www.acc.co.nz/PRD\\_EXT\\_CSMP/groups/external\\_ip/documents/guide/wim2\\_059285.pdf](http://www.acc.co.nz/PRD_EXT_CSMP/groups/external_ip/documents/guide/wim2_059285.pdf)
- IV. Vitamin D program - Information for pharmacists. 2011. Available at:  
[http://www.acc.co.nz/PRD\\_EXT\\_CSMP/groups/external\\_ip/documents/publications\\_promotion/wpc087748.pdf](http://www.acc.co.nz/PRD_EXT_CSMP/groups/external_ip/documents/publications_promotion/wpc087748.pdf)
- V. Vitamin D Prescribing Criteria. 2008. Available at: <http://www.acc.co.nz/search-results/index.htm?ssUserText=vitamin+D+prescribing+criteria+>
- VI. Identification of fall risks and interventions for falls and injury reduction - Clinical Practice Guideline (CPG) Residential Complex Care. 2009. Available at:  
[http://fhpulse/clinical\\_programs/residential\\_care\\_and\\_assisted\\_living/resources/CPG/Fraser%20Health/Falls%20CPG%20-%20Residential.pdf](http://fhpulse/clinical_programs/residential_care_and_assisted_living/resources/CPG/Fraser%20Health/Falls%20CPG%20-%20Residential.pdf)
- VII. Hanley DA, Cranney A, Jones G, Whiting SJ, Leslie WD. Vitamin D in adult health and disease: a review and guideline statement from Osteoporosis Canada (summary). *CMAJ*. 2010;182(12):1315-1319



REVIEW OF RECENT EVIDENCE ON VITAMIN D AND  
HEALTH





# Autoimmunity Reviews

journal homepage: [www.elsevier.com/locate/autrev](http://www.elsevier.com/locate/autrev)

## Review

### Vitamin D effects on musculoskeletal health, immunity, autoimmunity, cardiovascular disease, cancer, fertility, pregnancy, dementia and mortality—A review of recent evidence

Pawel Pludowski<sup>a,\*</sup>, Michael F. Holick<sup>b</sup>, Stefan Pilz<sup>c,d</sup>, Carol L. Wagner<sup>e</sup>, Bruce W. Hollis<sup>e</sup>, William B. Grant<sup>f</sup>, Yehuda Shoenfeld<sup>g</sup>, Elisabeth Lerchbaum<sup>c</sup>, David J. Llewellyn<sup>h</sup>, Katharina Kienreich<sup>c</sup>, Maya Soni<sup>h</sup>

<sup>a</sup> Department of Biochemistry, Radioimmunology and Experimental Medicine, The Children's Memorial Health Institute, Warsaw, Poland

<sup>b</sup> Department of Medicine, Section of Endocrinology, Nutrition, and Diabetes, Vitamin D, Skin and Bone Research Laboratory, Boston University Medical Center, Boston, MA, USA

<sup>c</sup> Department of Internal Medicine, Division of Endocrinology and Metabolism, Medical University of Graz, Austria

<sup>d</sup> Department of Epidemiology and Biostatistics, EMGO Institute for Health and Care Research, VU University Medical Centre, Amsterdam, The Netherlands

<sup>e</sup> Division of Neonatology, Department of Pediatrics, Children's Research Institute, Medical University of South Carolina, Charleston, SC, USA

<sup>f</sup> Sunlight, Nutrition, and Health Research Center, San Francisco, CA, USA

<sup>g</sup> Zabłudowicz Center for Autoimmune Diseases, Chaim Sheba Medical Center, Incumbent of the Laura Schwarz-Kipp Chair for Research of Autoimmune Diseases, Sackler Faculty of Medicine, Tel-Aviv University, Israel

<sup>h</sup> University of Exeter Medical School, Exeter, United Kingdom

## ARTICLE INFO

### Article history:

Received 12 February 2013

Accepted 28 February 2013

Available online 28 March 2013

### Keywords:

Vitamin D  
25(OH)D level  
Falls  
Fractures  
Cardiovascular  
Cancer  
Immunity  
Autoimmunity  
Fertility  
Pregnancy  
Infancy  
Infections  
Mortality  
Dementia

## ABSTRACT

**Background:** Optimal vitamin D intake and its status are important not only for bone and calcium-phosphate metabolism, but also for overall health and well-being. Vitamin D deficiency and insufficiency as a global health problem are likely to be a risk for wide spectrum of acute and chronic illnesses.

**Methods:** A review of randomized controlled trials, meta-analyses, and other evidence of vitamin D action on various health outcomes.

**Results:** Adequate vitamin D status seems to be protective against musculoskeletal disorders (muscle weakness, falls, fractures), infectious diseases, autoimmune diseases, cardiovascular disease, type 1 and type 2 diabetes mellitus, several types of cancer, neurocognitive dysfunction and mental illness, and other diseases, as well as infertility and adverse pregnancy and birth outcomes. Vitamin D deficiency/insufficiency is associated with all-cause mortality.

**Conclusions:** Adequate vitamin D supplementation and sensible sunlight exposure to reach optimal vitamin D status are among the front line factors of prophylaxis for the spectrum of disorders. Supplementation guidance and population strategies for the eradication of vitamin D deficiency must be included in the priorities of physicians, medical professionals and healthcare policy-makers.

© 2013 Elsevier B.V. All rights reserved.

## Contents

1. Introduction . . . . .	977
2. Defining vitamin D deficiency and its consequences on skeletal health . . . . .	977
3. Vitamin D and musculoskeletal system . . . . .	978

<sup>\*</sup> Sources of support: This work is supported in part by the Grant of MNISW 5412/B/P01/2010/39, EU Structural Grant # POIG.02.01.00-14-059/09, NIH CTSI Grant # UL1-RR025771, The Thrasher Research Fund, NIH RR01070, UL1 RR029882, The Children's Memorial Health Institute Internal Grants S109/2009 and 181/2007, The Alzheimer's Association Grant # NIRC-11-200737, and the Department of Biochemistry, Radioimmunology and Medicine, The Children's Memorial Health Institute, Warsaw, Poland as well as the Division of Neonatology, Medical University of South Carolina, Charleston, SC, USA.

\* Corresponding author at: The Children's Memorial Health Institute, Department of Biochemistry, Radioimmunology and Experimental Medicine, Aleja Dzieci Polskich 20, 04-730 Warsaw, Poland.

E-mail address: [p.pludowski@czd.pi](mailto:p.pludowski@czd.pi) (P. Pludowski).

Controversy remains regarding the appropriate adjustment for covariates in observational studies of vitamin D in relation to cognitive decline or dementia. For example, the sources of vitamin D itself (sunlight exposure, dietary intake, fortification and supplementation) are not likely to be confounders, and adjustment for these variables or proxy measures such as time spent outdoors or latitude is likely to represent over adjustment. Even adjustment for age is not without controversy, as human skin is known to become less efficient at vitamin D production with age. Age is therefore related to the synthesis of vitamin D and is not just a proxy measure for possible unmeasured confounding.

Ultimately randomized controlled trials are needed to establish whether vitamin D supplementation has clinical relevance in this context and can be used to prevent, delay or treat dementia. At this point no large well-designed randomized controlled trials have been conducted, and the causal relationship between vitamin D and dementia remains uncertain and caution should be exercised. Existing trials on vitamin D and cognitive decline have produced inconclusive results and had a number of important drawbacks including small sample sizes (<100) [195,196] and the use of low doses of vitamin D (<520 IU/day) with a combination of other nutrients [195,197], making interpretation difficult. However, several large trials are currently underway which will provide important new information. The DOHealth trial is being conducted in around 2000 participants aged 70 years and older across eight European cities. Vitamin D3 supplements (2000 IU/day) are one of the three interventions incorporated and cognitive outcomes will be measured over 3 years. Another key trial is the VITAL study in the US that aims to recruit around 20,000 middle aged and older adults. Again one of the interventions investigated will be vitamin D3 supplements (2000 IU/day), although cognitive outcomes over 4.5 years will only be assessed in a subsample of around 10% of participants. Neither trial targets older adults who are known to have low levels of vitamin D and early cognitive changes indicating that they are at high risk for dementia. If these trials do not produce promising results we may be left wondering if a more targeted approach or a different dose of vitamin D supplementation might be more effective.

## 11. Conclusion

It is now recognized that vitamin D deficiency and insufficiency are a global health problem [1,5,198–201]. A multitude of studies have suggested that vitamin D deficiency and insufficiency not only have negative consequences on bone health but are also likely to be a risk for many acute and chronic illnesses including infectious diseases, autoimmune diseases, cardiovascular disease, type 1 and type 2 diabetes mellitus, several types of cancer, neurocognitive dysfunction and mental illness, and other diseases, as well as infertility and adverse pregnancy and birth outcomes [24,26,37,49,55,75–79,85,90–94,100–105,109,117,118,136,141,146,186,187,202,203].

It is interesting that healthy black children in South Africa have blood levels of 25(OH)D of  $49 \pm 4$  ng/mL [204] similar to adult Maasai herders of  $47 \pm 10$  ng/mL [205]. It is well documented that blood levels of 25(OH)D are maximum at the end of the summer and are at their nadir at the end of the winter even in Denmark [206]. Physiologically it makes no sense to have wide swings in the circulating levels of 25(OH)D. This is the reason why a three-part strategy to maintain circulating levels of 25(OH)D of at least 30 ng/mL should be encouraged. Sensible sun exposure, which remains the major source of vitamin D for most children and adults [1,207], along with including foods that naturally contain or are fortified with vitamin D [1], and taking a daily supplement of vitamin D should be able to sustain blood levels of 25(OH)D in a range similar to our hunter-gatherer forefathers, i.e. 25(OH)D  $\sim 40$ – $50$  ng/mL. Since there is no downside to increasing children's and adults' vitamin D status (with the exception of patients with granulomatous disorders) it is reasonable to attain and maintain a circulating level of 25(OH)D of 40–60 ng/mL as recommended by the Endocrine Society Experts or even slightly lower (30–50 ng/mL)

as recommended in "Practical guidelines for supplementation of vitamin D and treatment of deficits in Central Europe: Recommended vitamin D intakes in general population and groups being at risk of vitamin D deficiency" [208], not only for optimal bone health but also for overall health and well-being.

## Take-home messages

- Vitamin D deficiency is a global health problem for children and adults. Vitamin D deficiency is associated with rickets and growth retardation in children and osteoporosis and osteomalacia in adults. Vitamin D deficiency has also been linked to many acute and chronic illnesses including some cancers, autoimmune diseases, cardiovascular disease, type 1 and type 2 diabetes mellitus, infectious diseases and neurocognitive dysfunction and other diseases, as well as infertility and adverse pregnancy and birth outcomes.
- A three-part strategy should be implemented to combat the vitamin D deficiency pandemic which includes:
  - Eating foods that naturally contain vitamin D,
  - Encouraging food fortification with vitamin D in countries that do not practice this fortification and,
  - Providing guidelines for both vitamin D supplementation of general population and for sensible sun exposure as a reliable source of vitamin D.
- Anti-fall and anti-fracture action of vitamin D administration of at least 800 IU/day with at least 24 ng/mL (60 nmol/L) of 25(OH)D serum levels appeared effective and beneficial for musculoskeletal machinery.
- Vitamin D may be instrumental in the immune system homeostasis, and in preventing autoimmune diseases and lowering risk of infections.
- Vitamin D deficient individuals are at increased cardiovascular risk even after adjustments for common cardiovascular risk factors.
- Risk for breast and colorectal cancer decreases as serum 25(OH)D level increases to 30–40 ng/mL (75–100 nmol/L).
- All-cause mortality risk in general population seems to be the lowest at 25(OH)D levels ranging from 30 to 45 ng/mL (75 to 112.5 nmol/L).
- Vitamin D supplementation up to 4000 IU/day in pregnant woman is safe and effective in achieving sufficiency and improving health not only in the mother but also in the developing fetus, every 10 ng/mL increase in maternal 25(OH)D at delivery reduces the risk of four main comorbidities of pregnancy by 16%.
- It is reasonable to attain and maintain a circulating level of 25(OH)D of 30–60 ng/mL as recommended by the Endocrine Society or even slightly lower (30–50 ng/mL) as recommended in "Practical guidelines for supplementation of vitamin D and treatment of deficits in Central Europe", not only for optimal bone health but also for overall health and well-being.

## References

- [1] Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266–81.
- [2] IOM (Institute of Medicine). Dietary reference intakes for calcium and vitamin D. Committee to review dietary reference intakes for calcium and vitamin D. Washington DC: The National Academies Press. Institute of Medicine; 2011.
- [3] Rosen CJ, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, et al. IOM committee members respond to Endocrine Society vitamin D guideline. *J Clin Endocrinol Metab* 2012;97(4):1146–52.
- [4] Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96(7):1911–30.
- [5] Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Controversy in clinical endocrinology: guidelines for preventing and treating vitamin D deficiency and insufficiency revisited. *J Clin Endocrinol Metab* 2012;97:1153–8.
- [6] Holick MF. Resurrection of vitamin D deficiency and rickets. *J Clin Invest* 2006;116(8):2062–72.

Total Cost per year Breast Cancer: \$6,048,625

**Vitamin D savings: \$3,024,312**

Colorectal Cancer Cost: \$3,925,900

**Vitamin D savings: \$1,962,950**

Diabetes Cost: \$4,777,952

**Vitamin D savings at 90%: \$4,299,527**

Preterm births

**Savings by avoiding 25 preterm births: \$1,375,000**

**Annual Cost Savings with vitamin D: \$10,661,789**

# Vitamin D for Health: A Global Perspective

Arash Hossein-nezhad, MD, PhD, and Michael F. Holick, PhD, MD

## Abstract

It is now generally accepted that vitamin D deficiency is a worldwide health problem that affects not only musculoskeletal health but also a wide range of acute and chronic diseases. However, there remains cynicism about the lack of randomized controlled trials to support the association studies regarding the nonskeletal health benefits of vitamin D. This review was obtained by searching English-language studies published up to April 1, 2013, in PubMed, MEDLINE, and the Cochrane Central Register of Controlled Trials (search terms: *vitamin D* and *supplementation*) and focuses on recent challenges regarding the definition of vitamin D deficiency and how to achieve optimal serum 25-hydroxyvitamin D concentrations from dietary sources, supplements, and sun exposure. The effect of vitamin D on fetal programming epigenetics and gene regulation could potentially explain why vitamin D has been reported to have such wide-ranging health benefits throughout life. There is potentially a great upside to increasing the vitamin D status of children and adults worldwide for improving musculoskeletal health and reducing the risk of chronic illnesses, including some cancers, autoimmune diseases, infectious diseases, type 2 diabetes mellitus, neurocognitive disorders, and mortality.

© 2013 Mayo Foundation for Medical Education and Research ■ Mayo Clin Proc. 2013;88(7):720-755



From the Department of Medicine, Section of Endocrinology, Nutrition, and Diabetes, Vitamin D, Skin, and Bone Research Laboratory, Boston University Medical Center, Boston, MA.

Vitamin D deficiency has been recognized as a pandemic with a myriad of health consequences.<sup>1,2</sup> Low vitamin D status has been associated with an increased risk of type 1 diabetes mellitus, cardiovascular disease, certain cancers, cognitive decline, depression, pregnancy complications, autoimmunity, allergy, and even frailty.<sup>1-4</sup> Low prenatal and neonatal vitamin D status may also increase susceptibility to schizophrenia, type 1 diabetes, and multiple sclerosis (MS) in later life via specific target organ effects, including the immune system, or through epigenetic modification.<sup>5</sup>

Despite the many important health benefits of vitamin D, there is controversy regarding the definition of vitamin D deficiency and what the vitamin D requirement should be.<sup>2,6,7</sup> In addition, critical windows of exposure to adequate vitamin D levels during fetal maturation remain to be defined<sup>5,6</sup> owing, in part, to the lack of well-designed controlled clinical trials with long-term follow-up.<sup>5-7</sup>

This review, obtained, in part, from searching English-language studies published up to April 1, 2013, in PubMed, MEDLINE, and the Cochrane Central Register of Controlled Trials (search terms: *vitamin D* and *supplementation*), focuses on recent challenges about how to achieve an optimal serum level of 25-hydroxyvitamin D [25(OH)D] from dietary sources, supplements,

and sun exposure and evidence-based benefits for skeletal and nonskeletal health. Also, we explore fetal programming and epigenomic mechanisms that could potentially explain why vitamin D has been reported to have such wide-ranging health benefits throughout life.

## VITAMIN D METABOLISM AND BIOLOGICAL FUNCTIONS

Vitamin D (D represents D<sub>2</sub>, D<sub>3</sub>, or both) is a secosteroid produced endogenously in the skin from sun exposure or obtained from foods that naturally contain vitamin D, including cod liver oil and fatty fish (eg, salmon, mackerel, and tuna); UV-irradiated mushrooms; foods fortified with vitamin D; and supplements.<sup>2,7</sup>

During exposure to sunlight, 7-dehydrocholesterol (7-DHC) in the skin is converted to previtamin D<sub>3</sub>. The 7-DHC is present in all the layers of human skin.<sup>7-9</sup> Approximately 65% of 7-DHC is found in the epidermis, and greater than 95% of the previtamin D<sub>3</sub> that is produced is in the viable epidermis and, therefore, cannot be removed from the skin when it is washed.<sup>9</sup> Once previtamin D<sub>3</sub> is synthesized in the skin, it can undergo either a photoconversion to lumisterol, tachysterol, and 7-DHC or a heat-induced membrane-enhanced isomerization to vitamin D<sub>3</sub> (Figure 1).<sup>7,8</sup> The cutaneous production of previtamin D<sub>3</sub> is regulated. Solar photoproducts (tachysterol and

lumisterol) inactive on calcium metabolism are produced at times of prolonged exposure to solar UV-B radiation, thus preventing sun-induced vitamin D intoxication.<sup>7,8</sup> Vitamin D<sub>3</sub> is also sensitive to solar irradiation and is, thereby, inactivated to suprasterol 1 and 2 and to 5,6-trans-vitamin D<sub>3</sub>.<sup>7</sup> Cutaneous vitamin D<sub>3</sub> production is influenced by skin pigmentation, sunscreen use, time of day, season, latitude, altitude, and air pollution.<sup>1,2,7,8</sup> An increase in the zenith angle of the sun during winter and early morning and late afternoon results in a longer path for the solar UV-B photons to travel through the ozone layer, which efficiently absorbs them. This is the explanation for why above and below approximately 33° latitude little if any vitamin D<sub>3</sub> is made in the skin during winter.<sup>10,11</sup> This is also the explanation for why—whether being at the equator and in the far northern and southern regions of the world in summer, where the sun shines almost 24 hours a day—vitamin D<sub>3</sub> synthesis occurs only between approximately 10 AM and 3 PM.<sup>1,11</sup> Similarly, in urban areas, such as Los Angeles, California, and Mexico City, Mexico, where nitrogen dioxide and ozone levels are high, few vitamin D<sub>3</sub>-producing UV-B photons reach the people living in these cities.<sup>7,11</sup> Similarly, because glass absorbs all UV-B radiation, no vitamin D<sub>3</sub> is produced in the skin when the skin is exposed to sunlight that passes through glass.


Once formed, vitamin D<sub>3</sub> is ejected out of the keratinocyte plasma membrane and is drawn into the dermal capillary bed by the vitamin D binding protein (DBP).<sup>7,8</sup> Vitamin D that is ingested is incorporated into chylomicrons, which are released into the lymphatic system, and enters the venous blood,<sup>2,7</sup> where it binds to DBP and lipoproteins transported to the liver.<sup>1,2,7</sup> Vitamin D<sub>2</sub> and vitamin D<sub>3</sub> are 25-hydroxylated by the liver vitamin D-25-hydroxylase (CYP2R1) to produce the major circulating vitamin D metabolite, 25(OH)D, which is used to determine a patient's vitamin D status.<sup>1,2,7</sup> This metabolite undergoes further hydroxylation by the 25(OH)D-1 $\alpha$ -hydroxylase (CYP27B1) in the kidneys to form the secosteroid hormone 1 $\alpha$ ,25-dihydroxyvitamin D [1,25(OH)<sub>2</sub>D] (Figure 1).<sup>2,7,12</sup> The 25(OH)D bound to DBP is filtered in the kidneys and is reabsorbed in the proximal renal tubules by megalin cubilin receptors.<sup>6,12</sup> The renal

#### ARTICLE HIGHLIGHTS

- Vitamin D deficiency is a common underdiagnosed condition.
- Recent evidence from hundreds of studies suggests that vitamin D is important for reducing the risk of type 1 diabetes mellitus, cardiovascular disease, certain cancers, cognitive decline, depression, pregnancy complications, autoimmunity, allergy, and even frailty.
- The blood level of 25(OH)D is the best method to determine vitamin D status.
- Vitamin D deficiency during pregnancy may influence fetal “imprinting” that may affect chronic disease susceptibility soon after birth as well as later in life.
- An effective strategy to prevent vitamin D deficiency and insufficiency is to obtain some sensible sun exposure, ingest foods that contain vitamin D, and take a vitamin D supplement.

1 $\alpha$ -hydroxylation is closely regulated, being enhanced by parathyroid hormone (PTH), hypocalcemia, and hypophosphatemia and inhibited by hyperphosphatemia, fibroblast growth factor-23, and 1,25(OH)<sub>2</sub>D itself.<sup>7,13,14</sup>

The 1,25(OH)<sub>2</sub>D performs many of its biologic functions by regulating gene transcription through a nuclear high-affinity vitamin D receptor (VDR).<sup>15,16</sup> This active metabolite of vitamin D binds to the nuclear VDR, which binds retinoic acid X receptor to form a heterodimeric complex that binds to specific nucleotide sequences in the DNA known as vitamin D response elements. Once bound, a variety of transcription factors attach to this complex, resulting in either up-regulation or down-regulation of the gene's activity.<sup>2,7,17</sup> There are estimated to be 200 to 2000 genes that have vitamin D response elements or that are influenced indirectly, possibly by epigenetics, to control a multitude of genes across the genome.<sup>2,16</sup> A recent microarray study on the influence of vitamin D status and vitamin D<sub>3</sub> supplementation on genome-wide expression in white blood cells before and after vitamin D<sub>3</sub> supplementation found that an improved serum 25(OH)D concentration was associated with at least a 1.5-fold alteration in the expression of 291 genes.<sup>17</sup> This study suggested that any improvement in vitamin D status will significantly affect the expression of genes that have a variety of biologic functions of



**FIGURE 1.** Schematic representation of the synthesis and metabolism of vitamin D for skeletal and nonskeletal function. 1-OHase = 25-hydroxyvitamin D-1 $\alpha$ -hydroxylase; 24-OHase = 25-hydroxyvitamin D-24-hydroxylase; 25(OH)D = 25-hydroxyvitamin D; 1,25(OH)<sub>2</sub>D = 1,25-dihydroxyvitamin D; CaBP = calcium-binding protein; CYP27B1, Cytochrome P450-27B1; DBP = vitamin D-binding protein; ECaC = epithelial calcium channel; FGF-23 = fibroblast growth factor-23; PTH = parathyroid hormone; RANK = receptor activator of the NF- $\kappa$ B; RANKL = receptor activator of the NF- $\kappa$ B ligand; RXR = retinoic acid receptor; TLR2/1 = Toll-like receptor 2/1; VDR = vitamin D receptor; vitamin D = vitamin D<sub>2</sub> or vitamin D<sub>3</sub>. Copyright Holick 2013, reproduced with permission.

more than 80 pathways linked to cancer, autoimmune disorders, and cardiovascular disease, which have been associated with vitamin D deficiency.<sup>17</sup>

One of the major physiologic functions of vitamin D is to maintain serum calcium and phosphorus levels in a healthy physiologic range to maintain a variety of metabolic functions, transcription regulation, and bone metabolism (Figure 1).<sup>2,7</sup> The 1,25(OH)<sub>2</sub>D interacts with its VDR in the small intestine to increase the efficiency of intestinal calcium absorption from approximately 10% to 15% up to 30% to 40% and intestinal phosphorus absorption from approximately 60% to 80%.<sup>7</sup> It also interacts with VDR in osteoblasts to stimulate a receptor activator of nuclear factor  $\kappa$ B ligand, which, in turn, interacts with receptor activator of nuclear factor  $\kappa$ B on immature preosteoclasts, stimulating them to become mature bone-resorbing osteoclasts (Figure 1).<sup>7,18</sup> The mature osteoclast removes calcium and phosphorus from the bone to maintain blood calcium and phosphorus levels. In the kidneys, 1,25(OH)<sub>2</sub>D stimulates calcium reabsorption from the glomerular filtrate.<sup>2,7</sup>

The VDR is present in most tissues and cells in the body.<sup>1,2,7,19-25</sup> Many of these organs and cells, including the brain, vascular smooth muscle, prostate, breast, and macrophages, not only have a VDR but also have the capacity to produce 1,25(OH)<sub>2</sub>D.<sup>1,2,7,19-25</sup> This production probably depends on the availability of circulating 25(OH)D, indicating the biological importance of sufficient blood levels of this vitamin D metabolite.<sup>2,15,26</sup>


The estimated 2000 genes that are directly or indirectly regulated by 1,25(OH)<sub>2</sub>D<sup>2,7,17-24,26</sup> have a wide range of proven biological actions, including inhibiting cellular proliferation and inducing terminal differentiation, inhibiting angiogenesis, stimulating insulin production, inducing apoptosis, inhibiting renin production, and stimulating macrophage cathelicidin production.<sup>1,2,7,16,17,26</sup> In addition, 1,25(OH)<sub>2</sub>D stimulates its own destruction in the kidneys and in cells that have a VDR and responds to 1,25(OH)<sub>2</sub>D by enhancing expression of the 25(OH)D-24-hydroxylase (CYP24A1) to metabolize 25(OH)D and 1,25(OH)<sub>2</sub>D into water-soluble inactive forms that are excreted in the bile (Figure 1).<sup>1,7,19,27</sup>

## VITAMIN D METABOLISM DURING PREGNANCY

Vitamin D metabolism is enhanced during pregnancy and lactation. The placenta is formed at 4 weeks of gestation.<sup>2,25</sup> From this time to term, 25(OH)D is transferred across the placenta, and the fetal cord blood concentration of 25(OH)D is correlated with the mother's concentration.<sup>27</sup> However, the active metabolite 1,25(OH)<sub>2</sub>D does not readily cross the placenta.<sup>25,27</sup> The fetal kidneys and the placenta provide the fetal circulation with 1,25(OH)<sub>2</sub>D by expressing CYP27B1 (Figure 2).<sup>28</sup>

The maternal (decidual) and fetal placental (trophoblastic) components of the placenta have CYP27B1 activity; cultured human syncytiotrophoblasts and decidual cells synthesize 1,25(OH)<sub>2</sub>D.<sup>27</sup> The spatiotemporal organization of placental CYP27B1 and the VDR across gestation has also been characterized, confirming that the enzyme and receptor are localized to the maternal and fetal parts of the placenta.<sup>29</sup> Serum levels of DBP increase 46% to 103% during pregnancy, suggesting that DBP may play a role in directing vitamin D metabolism and function during pregnancy.<sup>2,27,30</sup> The DBP has a much higher binding affinity for 25(OH)D than for 1,25(OH)<sub>2</sub>D, and in kidney epithelial cells, DBP plays a pivotal role in conserving 25(OH)D by facilitating the recovery of 25(OH)D from the glomerular filtrate.<sup>31,32</sup>

Transplacental transfer of calcium to the fetus is also facilitated by expression of all the key mediators of vitamin D metabolism in the placenta. Hormones involved in fetal growth and that influence CYP27B1 activity include insulin-like growth factor 1 and human placental lactogen, PTH-related protein (PTHrP), estradiol, and prolactin.<sup>2,31,33,34</sup> The PTHrP acts as a calciotropic hormone during fetal life and in lactation.<sup>35-37</sup> The exact role of circulating PTHrP in pregnancy is unknown, but its rise may stimulate renal CYP27B1 and contribute to the increase in 1,25(OH)<sub>2</sub>D concentration and, indirectly, the suppression of PTH levels.<sup>27,35,37</sup> The PTHrP arises from several sources, including the breast, myometrium, decidua, amnion, and fetal parathyroids.<sup>36</sup> Several roles of PTHrP are postulated from animal studies, including fetal chondrocyte maturation, fetal calcium transfer, and stimulation of CYP27B1 activity.<sup>33,34,36</sup> Furthermore, the carboxy terminal of PTHrP (osteostatin) may suppress osteoclastic activity



**FIGURE 2.** Vitamin D metabolism during pregnancy and lactation. Maternal 25(OH)D is thought to freely cross the human placenta. The placenta expresses vitamin D receptors (VDR) and also produces 1-OHase to convert 25(OH)D to 1,25(OH)<sub>2</sub>D. 1,25-dihydroxyvitamin D does not readily cross the placenta, and fetal 1,25(OH)<sub>2</sub>D levels are normally lower than maternal serum levels. The low fetal concentrations of 1,25(OH)<sub>2</sub>D reflect the low fetal PTH and high phosphorus concentrations, which suppress renal 1-OHase. Although PTHrP is elevated in the fetal circulation, it appears to be less able to stimulate the renal 1-OHase than PTH. Total (free and bound) 1,25(OH)<sub>2</sub>D concentrations double or triple in the maternal circulation starting in the first trimester, but studies have only shown increased free concentrations during the third trimester. This increase is due to maternal synthesis by the renal 1-OHase. Vitamin D passes readily into breast milk, 25(OH)D passes very poorly, and 1,25(OH)<sub>2</sub>D does not appear to pass at all.<sup>2</sup> 1,25(OH)<sub>2</sub>D levels fall rapidly after pregnancy and are normal during lactation.<sup>7</sup> Near-exclusive breastfeeding for 6 months leads, on average, to maternal calcium loss 4 times higher than that in pregnancy. Phosphorus can rise above the normal range, probably because of accelerated resorption from the skeleton. Parathyroid hormone-related protein levels are higher than PTH concentrations in nonpregnant women and show some pulsatility in response to suckling. Parathyroid hormone-related protein (produced by the lactating breast) in combination with low estradiol concentrations appears to drive the main physiologic adaptation to meet the calcium demands of lactation. Suckling and prolactin both inhibit ovarian function and stimulate PTHrP. Together, PTHrP and low estradiol concentrations stimulate skeletal resorption. Renal calcium reabsorption rates increase, presumably due to PTHrP, which mimics the actions of PTH on the renal tubules. *For definitions of abbreviations, see Figure 1.* Copyright Holick 2013, reproduced with permission.

and may have a possible bone protection role in the mother during pregnancy.<sup>32,35-37</sup>


Calcitonin, an important component of calcium homeostasis during pregnancy,<sup>38,39</sup> is known to promote transcription of CYP27B1<sup>40</sup> and may, therefore, be a key determinant of placental vitamin D metabolism.<sup>41</sup> Thus, PTHrP and calcitonin, as well as other factors, cause 1,25(OH)<sub>2</sub>D levels to increase, being 2-fold higher in serum of women in the third trimester of pregnancy than in nonpregnant or postpartum women.<sup>2,27</sup> Normally, 1,25(OH)<sub>2</sub>D regulates its own metabolism via a feedback loop such that elevated concentrations induce the expression of CYP24A1, with concomitant down-regulation of CYP27B1.<sup>7,25,42</sup> This process results in a reduction of 25(OH)D and 1,25(OH)<sub>2</sub>D levels.<sup>9,29,30</sup> However, during pregnancy, this process becomes uncoupled, resulting in elevated maternal concentrations of circulating 1,25(OH)<sub>2</sub>D.<sup>27,43</sup> The placental methylation of the CYP24A1 promoter reduces the capacity for CYP24A1 induction and down-regulates basal promoter activity and abolishes vitamin D-mediated feedback activation. This epigenetic decoupling of vitamin D feedback catabolism also plays an important role in enhancing 1,25(OH)<sub>2</sub>D bioavailability at the fetomaternal interface.<sup>44</sup>

#### VITAMIN D, PLACENTA DEVELOPMENT, FETAL PROGRAMMING, AND EPIGENETIC MODIFICATION

Epidemiologic evidence has suggested a link between fetal life events and susceptibility to disease in adult life.<sup>45-47</sup> This paradigm, referred to as *fetal programming* or *developmental origins of health and disease*, may have a profound effect on public health strategies for the prevention of major illnesses.<sup>2,48</sup> The role of vitamin D in implantation tolerance and placental development has been studied. The 1,25(OH)<sub>2</sub>D<sub>3</sub> regulates key target genes associated with implantation, such as Homeobox A10 (*HOXA10*), whereas the potent immunosuppressive effects of 1,25(OH)<sub>2</sub>D<sub>3</sub> suggest a role in placental development.<sup>49</sup> Increasing expression of CYP27B1 and VDR in first-trimester human trophoblasts and deciduas<sup>50</sup> may be related to the immunosuppressive effects of 1,25(OH)<sub>2</sub>D<sub>3</sub> and may help improve implantation tolerance. Placental development plays a critical role in pregnancy health, and its link to maternal vitamin D deficiency may

explain related adverse outcomes.<sup>5,45</sup> In neonatal rats exposed prenatally to low maternal serum 25(OH)D levels, there was a general slowing of cardiac development, with significantly lower heart weights, decreased citrate synthase and 3-hydroxyacyl CoA dehydrogenase activity, and a 15% lower myofibrillar protein content.<sup>46</sup> A 2-month-old human infant with dilated cardiomyopathy and severe vitamin D deficiency had dramatic improvement of her ejection fraction (17%-66%) after vitamin D supplementation.<sup>47</sup> In addition, maternal vitamin D deficiency in rats stimulated nephrogenesis in offspring, with a 20% increase in nephron number but a decrease in renal corpuscle size observed between replete and deficient rats, despite there being no difference in body weight or kidney weight and volume.<sup>5,51</sup> These findings support the role of vitamin D influencing fetal programming and placental development.

Epigenetic modification refers to heritable changes in gene expression that are not mediated by alterations in DNA sequence.<sup>52</sup> This hypothesis, first articulated by Barker et al,<sup>53</sup> postulated that in utero epigenetic fetal programming (as a result of environmental events during pregnancy) induced specific genes and genomic pathways that controlled fetal development and subsequent disease risk. The role of vitamin D in epigenetic modification and fetal programming could potentially explain why vitamin D has been reported to have such wide-ranging health benefits. Recent studies have suggested that epigenetic decoupling of vitamin D feedback catabolism plays an important role in maximizing 1,25(OH)<sub>2</sub>D bioavailability at the fetomaternal interface.<sup>25,44</sup> Modified expression of the genes encoding placental calcium transporters, by epigenetic regulation by 1,25(OH)<sub>2</sub>D, might represent the means whereby maternal vitamin D status could influence bone mineral accrual in the neonate.<sup>54,55</sup> Vitamin D deficiency during pregnancy may, therefore, not only impair maternal skeletal preservation and fetal skeletal formation but also influence fetal "imprinting" that may affect chronic disease susceptibility soon after birth as well as later in life (Figure 3).<sup>15,56</sup> Transgenerational hormonal imprinting caused by vitamin D treatment of newborn rats has been previously reported.<sup>57</sup> A recent study reported that VDR binds to the  $\epsilon$  germline gene promoter and exhibits transrepressive activity.<sup>58</sup> Inhibition



**FIGURE 3.** Vitamin D, placenta development, fetal programming, and epigenetic modification. A= adenosine; CH<sub>3</sub> = methyl group; HIV = human immunodeficiency virus; IVF = in vitro fertilization; miRNA, micro RNA; SAH = S-adenosylhomocysteine; SAM= single carbon from adenosylmethionine. Copyright Holick 2013, reproduced with permission.


of IgE production by 1,25(OH)<sub>2</sub>D was mediated by its transrepressive activity through the VDR-corepressor complex, affecting chromatin compacting around the I $\epsilon$  region.<sup>58</sup> Also, the associations of early-life sun exposure and germline variation in VDR and CYP24A1 with non-Hodgkin lymphoma risk was reported in a clinic-based case-control study.<sup>59</sup>

#### DEFINITION OF VITAMIN D DEFICIENCY

The blood level of 25(OH)D is the best method to determine vitamin D status. Although 1,25(OH)<sub>2</sub>D is the biologically active form, it provides no information about vitamin D status because it is often normal or even elevated in children and adults who are vitamin D deficient.<sup>7,15,60-63</sup> Recently, the Institute of Medicine (IOM) and the Endocrine Society released separate guidelines for vitamin D requirements.<sup>60,62</sup> The recommended dietary allowances (RDAs)

of the IOM and the Endocrine Society guidelines for vitamin D intake are summarized in Figure 4.

The revised guidelines by the IOM stress that the daily requirements for vitamin D are generally met by most of the population and are appropriate to reach the "sufficient" level of 20 ng/mL (to convert to nmol/L, multiply by 2.496).<sup>62</sup> The IOM guidelines used a population model to prevent vitamin D deficiency in 97.5% of the general population. Also, note that the IOM report focuses only on bone health (calcium absorption, bone mineral density, and osteomalacia/rickets) and found no evidence that a serum 25(OH)D concentration greater than 20 ng/mL had beneficial effects at a population level. However, considering the available evidence on skeletal and extraskeletal effects of vitamin D, the few negative studies, and the lack of toxicity potential of vitamin D supplementation at



recommended doses, the US Endocrine Society, which used a medical model, recommended that serum 25(OH)D levels of 30 ng/mL should be attained to avoid other risks connected with an inadequate vitamin D status.<sup>7,60</sup> Therefore, the Endocrine Society recommended that vitamin D deficiency be defined as a 25(OH)D level of 20 ng/mL or less, vitamin D insufficiency as 21 to 29 ng/mL, and vitamin D sufficiency as 30 ng/mL or greater for children and adults.<sup>60</sup> It suggested that maintenance of a 25(OH)D level of 40 to 60 ng/mL is ideal (this takes into account assay variability) and that up to 100 ng/mL is safe.<sup>60</sup>

**MUSCULOSKELETAL CONSEQUENCES OF VITAMIN D DEFICIENCY**

According to current evidence from biochemical testing, observational studies, and randomized controlled trials (RCTs), serum 25(OH)D levels of at least 20 ng/mL are required for normalization of PTH levels, to minimize the risk of osteomalacia, and for optimal bone and muscle function, with many experts regarding 30 ng/mL as the threshold for optimal bone health.<sup>7,16,61,64-66</sup> The skeletal consequences of 25(OH)D insufficiency include secondary hyperparathyroidism, increased bone turnover and bone loss, and increased risk of low-trauma fractures.<sup>7,15,61,64</sup>

The most common etiology of rickets, historically and presently, is vitamin D deficiency. Low maternal 25(OH)D levels were found to correlate with increased fetal distal femoral splaying, determined by ultrasonography measurements of femoral length and metaphyseal width.<sup>63,65</sup> Children begin to manifest classic clinical signs of rickets between 6 months and 1.5 years that include rachitic rosary, widened epiphyseal plates at the end of long bones, and bowing deformities of the legs.<sup>66</sup> A common early symptom in newborns is excessive sweating due to neuromuscular irritability,<sup>66</sup> and a 25(OH)D level less than 20 ng/mL is common in children presenting with vague limb or back pain.

From a skeletal perspective for adults, evidence from RCTs suggests that vitamin D may be considered a threshold nutrient, with little bone benefit observed at levels of 25(OH)D above which PTH is normalized.<sup>62,67</sup> A literature review of 70 studies generally found a threshold for a decline in serum PTH levels with increasing serum 25(OH)D levels, but there was no consistency in the threshold level of serum 25(OH)D, which varied from 20 to 50 ng/mL.<sup>68</sup> A study of 4100 older adults (>60 years old) from the Third National Health and Nutrition Examination Survey (NHANES III) found that higher 25(OH)D levels were associated with better lower extremity function.<sup>61</sup> Much of the improvement occurred at 25(OH)D levels ranging from 9 to 16 ng/mL but continued to be seen at levels up to 40 ng/mL.<sup>69</sup> A systematic review revealed that supplemental vitamin D at daily doses of 800 to 1000 IU consistently had beneficial effects on muscle strength and balance.<sup>70</sup> Several RCTs have reported positive effects of vitamin D supplementation on muscle function and fall prevention.<sup>71-73</sup> Adequate calcium intake is imperative to gain optimal benefit from improving the vitamin D status in those with insufficient 25(OH)D levels.<sup>67</sup> In contrast, a study of 173 young Asian Indian females revealed that after supplementation with vitamin D<sub>3</sub> (60,000 IU/wk for 8 weeks followed by 60,000 IU every 2 weeks) and calcium (500 mg twice per day for 6 months), and despite significant improvement in serum 25(OH)D levels, there was no significant change in their skeletal muscle strength.<sup>74</sup> Thus, age, baseline and final 25(OH)D concentrations, and whether and how much calcium supplementation was included

in the clinical trial could affect outcome measures related to muscle performance and vitamin D status.

Proximal muscle weakness is a prominent clinical feature of vitamin D deficiency.<sup>7,60</sup> The relative contributions of vitamin D and calcium for reducing fracture risk remain unclear<sup>75</sup> because improving calcium intake is also associated with suppression of PTH levels independent of vitamin D status.<sup>67,76,77</sup> A meta-analysis of data from RCTs found a dose-response relationship between a higher vitamin D dose and higher achieved serum 25(OH)D levels, with prevention of falls and fractures.<sup>73</sup> The greatest benefit was observed at 700 to 1000 IU/d or a mean serum 25(OH)D concentration of 30 to 44 ng/mL.<sup>71,73</sup> Similar results were reported in a more recent meta-analysis of pooled participant-level data from 11 double-blind RCTs of oral vitamin D supplementation, with or without calcium, compared with placebo or calcium alone in persons 65 years or older.<sup>78</sup> Reduction in the risk of fracture occurred only at the highest vitamin D intake level (median, 800 IU/d; range, 792-2000 IU/d), with a 30% reduction in the risk of hip fracture and a 14% reduction in the risk of any nonvertebral fracture.<sup>78</sup> This reduction was independent of the assigned treatment dose of vitamin D, age group, sex, type of dwelling, and study.<sup>78</sup> Several previous meta-analyses have suggested that the dose of vitamin D is irrelevant when vitamin D is combined with calcium.<sup>79-82</sup> In contrast, a pooled subgroup analysis of the 8 double-blind RCTs that used vitamin D combined with calcium indicated that with combined supplementation, the risk of fracture was reduced only at the highest actual intake level of vitamin D. These findings support that a 25(OH)D level of more than 24 ng/mL may be most beneficial for reducing the risk of fractures.<sup>78</sup>

With a similar tone and theme, a report from the US Preventive Services Task Force concluded that current evidence is insufficient to assess the balance of benefits and harms of combined vitamin D and calcium supplementation for the primary prevention of fractures in premenopausal women or in men.<sup>83</sup> Furthermore, they concluded that there was insufficient evidence to assess the balance of benefits and harms of daily supplementation with greater than 400 IU of vitamin D<sub>3</sub> and greater than 1000 mg of calcium for primary prevention of fractures in noninstitutionalized postmenopausal women.

They recommended against daily supplementation with 400 IU or less of vitamin D<sub>3</sub> and 1000 mg or less of calcium for the primary prevention of fractures in noninstitutionalized postmenopausal women. They also stated that it was unclear whether higher doses of vitamin D and calcium are effective in preventing fractures in postmenopausal women, younger women, or men.<sup>83</sup> The Task Force, however, concluded that vitamin D supplementation is effective in preventing falls in community-dwelling adults 65 years or older, which, in turn, reduces the risk of fracture. This could help explain the observation by the Women's Health Initiative (WHI) that, in the subgroup of long-adherent women who took their calcium and vitamin D, there was a reduced risk of hip but not total fractures.<sup>84</sup> Therefore, what is still unknown is whether adequate intake of calcium, especially from dietary sources, and maintenance of serum 25(OH)D levels of at least 20 ng/mL as recommended by the IOM<sup>62</sup> or at least 30 ng/mL as recommended by the Endocrine Society<sup>60</sup> throughout life will reduce the risk of fracture. Most evidence suggests that adequate calcium and vitamin D intake along with exercise during childhood will maximize bone mineral content that can be sustained in young and middle-aged adults as long as they also have a healthy lifestyle, adequate calcium intake, and a healthy vitamin D status.<sup>60-62,84-86</sup> Accruing maximum bone mineral content during childhood, and maintaining peak bone mineral density in young and middle-aged adults, will likely reduce the risk of fracture later in life, when there is a disruption in bone remodeling due to menopause and aging.

Recent recommendations of the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO)<sup>87</sup> for the optimal management of elderly and postmenopausal women regarding vitamin D supplementation have also indicated that patients with serum 25(OH)D levels less than 20 ng/mL have increased bone turnover, bone loss, and, possibly, mineralization defects compared with patients with serum 25(OH)D levels of 20 ng/mL or greater. Similar relationships have been reported for frailty, nonvertebral and hip fracture, and all-cause mortality, with poorer outcomes at less than 20 ng/mL.<sup>87</sup> Thus, ESCEO recommended that 20 ng/mL be the minimal serum 25(OH)D concentration

at the population level and in patients with osteoporosis to ensure optimal bone health. Also, in fragile elderly individuals who are at elevated risk for falls and fractures, ESCEO recommended a minimal serum 25(OH)D level of 30 ng/mL for the greatest effect on fracture.<sup>87</sup> This coincides with the recommendation from the Endocrine Society<sup>60</sup> and with the observation of Priemel et al,<sup>79</sup> who reported that of 675 presumed healthy adults (aged 20-90+ years) who died in an accident, 36% had evidence of osteomalacia. However, Priemel et al<sup>79</sup> observed no osteomalacia in those who had a 25(OH)D level greater than 30 ng/mL.

#### EVIDENCE-BASED SKELETAL AND NONSKELETAL HEALTH BENEFITS OF VITAMIN D

Observational studies have found a decreased risk of many disorders, including certain types of cancer, mental disorders, infectious disease, cardiovascular disease, type 2 diabetes mellitus, and autoimmune disorders, associated with serum 25(OH)D levels greater than 28 to 32 ng/mL.<sup>7,60,67</sup> It has, therefore, been argued that 25(OH)D levels should be in the range of 28 to 40 ng/mL to maximize these nonskeletal benefits.<sup>1,2,7,19,60,61</sup>

The results of some clinical trials provide evidence confirming the results of observational and association studies, whereas others do not. The Table summarizes the meta-analyses on vitamin D supplementation, comparing the beneficial and nonbeneficial effects of vitamin D supplementation in randomized trials for musculoskeletal and nonskeletal outcomes. The Table provides the foundation for clinical decision making for recommending vitamin D supplementation and identifies gaps in our knowledge that require additional RCTs to provide insights as to whether vitamin D supplementation has nonskeletal health benefits.

#### VITAMIN D AND NONSKELETAL HEALTH ASSOCIATIONS AND MECHANISMS

##### Cancers

Association studies have related higher serum levels of 25(OH)D to reduced incidence of many types of cancers. It has been hypothesized that the local conversion of 25(OH)D to 1,25(OH)<sub>2</sub>D in healthy cells in the colon, breast, and prostate can help prevent malignancy by

TABLE. Summary of Meta-analyses of Vitamin D Supplementation

Reference, year	Included studies	Sample size (No.)	Participants	Dose/duration	Outcomes	Effects
Thome-Lyman and Fawzi, <sup>88</sup> 2012	5 randomized trials 2 observational studies	28,943	Pregnant women	Vitamin D <sub>2</sub> and vitamin D <sub>3</sub> Various doses and patterns of intake during pregnancy	Perinatal and infant health	Positive effect on low birth weight No effect on small-for-gestational-age (2 trials) No effect on preterm delivery
De-Regil et al, <sup>89</sup> 2012	6 trials	1023	Women during pregnancy	Vitamin D (1200 IU/d) alone or combined with 375 mg of elemental calcium	Safely improve maternal and neonatal outcomes	No effect on preeclampsia Positive effect on concentrations of 25(OH)D in serum Positive effect on birth weight No effect on adverse effects
Bischoff-Ferrari et al, <sup>78</sup> 2012	11 double-blind RCTs	31,022	Persons aged $\geq 65$ y	Oral vitamin D supplementation with or without calcium	Fracture reduction	No effect on risk of hip fracture until 800 IU/d Positive effect on hip and any nonvertebral fracture by highest intake level according to quartiles
Lai et al, <sup>90</sup> 2010	7 eligible RCTs and 17 identified case-control studies	801	Persons aged 74.8-85 y	Vitamin D <sub>2</sub> and vitamin D <sub>3</sub> (400-1100 IU)	Hip fracture risk	No effect on hip fracture risk
Bergman et al, <sup>80</sup> 2010	8 controlled trials	12,658	Postmenopausal women	Vitamin D <sub>3</sub> supplementation (800 IU/d) with or without calcium	Increasing BMD Preventing fractures	Positive effect on nonvertebral and hip fractures
Bischoff-Ferrari et al, <sup>81</sup> 2009	12 double-blind RCTs	83,165	Older individuals ( $\geq 65$ y)	>400 IU/d	Preventing nonvertebral and hip fractures	Positive effect on nonvertebral fracture prevention with vitamin D is dose dependent (only high dose)
Avenell et al, <sup>82</sup> 2009	45 trials	83,741	Older people	Vitamin D or related compounds	Preventing fractures	Positive effect by vitamin D with calcium on hip fracture No effect by vitamin D alone on hip fracture
Abrahamsen et al, <sup>75</sup> 2010	7 major randomized trials	68,517	Persons aged 47-107 y	Vitamin D <sub>2</sub> and vitamin D <sub>3</sub> (10 $\mu$ g/d to 300,000 IU/12 mo)	Antifracture efficacy	Positive effect of vitamin D with calcium on fracture No effect of vitamin D alone
Izaks, <sup>91</sup> 2007	11 trials	NA	General population	Vitamin D <sub>2</sub> and vitamin D <sub>3</sub> follow-up >1 y	Fracture risk	High-dose vitamin D may be effective in institutionalized persons but probably is not effective in the general population
Jackson et al, <sup>92</sup> 2007	9 studies	2410	Postmenopausal women	Vitamin D <sub>3</sub> (excluding the potential effect of calcium supplementation)	Risk of fall and fracture	Positive effect on risk of fall in patients treated with vitamin D <sub>3</sub>
Boonen et al, <sup>93</sup> 2007	10 RCTs	54,592	Postmenopausal women and/or older men ( $\geq 50$ y)	Oral vitamin D with or without calcium vs placebo/no treatment	Prevention of hip fractures	Positive effect of oral vitamin D on reducing the risk of hip fractures only with calcium supplementation
Avenell et al, <sup>94</sup> 2005	57 trials	82,986	Older people	Vitamin D or an analogue, alone or with calcium, vs placebo	Fracture	Positive effect of vitamin D with calcium supplements on hip and other nonvertebral fractures

Bischoff-Ferrari et al, <sup>84</sup> 2005	5 RCTs for hip fracture 7 RCTs for nonvertebral fracture risk	19,114	Older people	Oral vitamin D supplementation (cholecalciferol, ergocalciferol) with or without calcium supplementation vs calcium supplementation	Preventing hip and nonvertebral fractures	Positive effect (700-800 IU/d) on hip and any nonvertebral fractures in ambulatory or institutionalized elderly persons No effect (400 IU/d) on fracture prevention
Winzenberg et al, <sup>85</sup> 2011	6 studies	884	Healthy children and adolescents (aged 1 mo to <20 y)	Vitamin D supplementation vs placebo for $\geq 3$ mo	Improving BMD (effects vary with factors such as vitamin D dose and vitamin D status)	No effect on total body BMC or on hip or forearm BMD Positive small effect on lumbar spine BMD Positive effect with low serum vitamin D on total body BMC and lumbar spine bone
Huncharek et al, <sup>86</sup> 2008	21 RCTs	NA	Children	Dietary calcium/dairy supplementation	BMC	Positive effect of dietary calcium/dairy products, with and without vitamin D, on total body and lumbar spine BMC in children (with low baseline intakes)
Bischoff-Ferrari et al, <sup>95</sup> 2009	8 RCTs	2426	Older individuals	Vitamin D <sub>2</sub> and vitamin D <sub>3</sub> (200-1000 IU) with or without calcium	Preventing falls	Positive effect of supplemental vitamin D (700-1000 IU/d) on the risk of falling No effect at a dose <700 IU
Kalyani et al, <sup>96</sup> 2010	10 articles	2932	Older adults (aged $\geq 60$ y)	200-1000 IU/d of vitamin D for 1-36 mo	Fall prevention	Positive effect on fall prevention
Chung et al, <sup>97</sup> 2011	19 RCTs (3 for cancer and 16 for fracture outcomes) 28 observational studies (for cancer outcomes)	NA	Adults	Vitamin D with or without calcium (limited data from RCTs assessed high-dose vitamin D [1000 IU/d])	Benefits and harms of vitamin D with or without calcium supplementation on clinical outcomes of cancer and fractures	Positive effect of high-dose vitamin D on reduced risk of total cancer Positive effect on fracture Negative effect on renal and urinary tract stones
Buttiglierio et al, <sup>98</sup> 2011	25 studies (3 randomized trials involving patients with advanced prostate cancer explored the prognostic role of vitamin D supplementation)	1273	Cancer patients	1 trial: doxercalciferol 2 trials: calcitriol Duration: 11.7-18.32 mo	Influence of hypovitaminosis D on prognosis of cancer Improvement outcome of vitamin D supplementation	No effect on survival
Bjelakovic et al, <sup>99</sup> 2011	50 randomized trials	94,148	Adults; Most trials included elderly women (>70 y)	Supplemental vitamin D (vitamin D <sub>3</sub> [cholecalciferol] or vitamin D <sub>2</sub> [ergocalciferol]) or an active form of vitamin D (1 $\alpha$ -hydroxyvitamin D [alfacalcidol] or 1,25-dihydroxyvitamin D [calcitriol]) at any dose, duration, and route of administration vs placebo or no intervention	Beneficial and harmful effects of vitamin D for prevention of mortality	Positive effect of vitamin D <sub>3</sub> on mortality Negative effect of vitamin D <sub>3</sub> combined with calcium on nephrolithiasis Negative effect on hypercalcemia

Continued on next page

TABLE. Continued

Reference, year	Included studies	Sample size (No.)	Participants	Dose/duration	Outcomes	Effects
Iram et al, <sup>100</sup> 2010	16 additional trials (only 1 trial was single supplements of vitamin D)	22,120 participants in the trials	Adults and children with HIV infection	NA	Reducing mortality and morbidity	No effect
Autier et al, <sup>101</sup> 2012	76 trials	6207	White persons aged >50 y	Doses of 5-250 µg/d (median, 20 µg/d)	Circulating 25(OH)D level	Positive effect of vitamin D <sub>3</sub> intake without calcium on serum 25(OH)D concentrations No effect of concomitant use of calcium supplementation and high 25(OH)D concentration at baseline
Bjorkman et al, <sup>102</sup> 2009	52 clinical trials	6290	Chronically immobile patients	Vitamin D supplementation	Responses of parathyroid hormone	Positive effect in chronically immobile patients on 25(OH)D levels but a slight effect on PTH decrease
Tripkovic et al, <sup>103</sup> 2012	17 studies	1016	Persons aged 18-97 y	Varying dosages and treatment periods	Compared the effects of vitamin D <sub>2</sub> and vitamin D <sub>3</sub> on serum 25(OH)D concentrations	Positive effect of vitamin D <sub>3</sub> compared with vitamin D <sub>2</sub> in the raising of serum 25(OH)D concentrations
Kandula et al, <sup>104</sup> 2011	22 studies	264	Patients with non-dialysis-dependent CKD, dialysis-dependent CKD, and renal transplant	Ergocalciferol or cholecalciferol	Benefits and harms of vitamin D supplementation	Positive effect on 25(OH)D and PTH levels
Song et al, <sup>105</sup> 2013	21 prospective studies	81,216	Healthy individuals and patients with type 2 diabetes	Circulating 25(OH)D	Association between blood levels of 25(OH)D and risk of incident type 2 diabetes	Inverse and significant association between circulating 25(OH)D levels and risk of type 2 diabetes
George et al, <sup>106</sup> 2012	15 trials	NA	Nondiabetes to diabetes	Vitamin D or analogues	Glycemia, insulin resistance, progression to diabetes, and complications of diabetes	No effect on fasting glucose, hemoglobin A <sub>1c</sub> , or insulin resistance Small positive effect on fasting glucose and insulin resistance in patients with diabetes or impaired glucose tolerance No effect on glycated hemoglobin in diabetic patients
Bath-Hextall et al, <sup>107</sup> 2012	11 studies	596	Atopic eczema/dermatitis	Vitamin D vs vitamin E	Treating established atopic eczema/dermatitis	Negative effect at high doses
Muir et al, <sup>70</sup> 2011	13 trials	NA	Older adults (≥60 y)	Vitamin D (800-1000 IU/d)	Muscle strength, gait, and balance	Positive effect on balance and muscle strength
Annweiler et al, <sup>108</sup> 2009	16 trials	24-33,067	Persons aged ≥80 y	NA	Muscle, balance, and gait performance	No significant effect on balance and gait Positive/no effect on muscle strength No effect on sit-to-stand test

Wang et al, <sup>109</sup> 2010	18 trials	NA	Adults	NA	Reduce the risk of cardiovascular events	No effect on cardiovascular disease risk
Pittas et al, <sup>110</sup> 2010	18 trials	37,162	Generally healthy adults	Vitamin D (400-8571 IU/d) with or without calcium	Cardiometabolic outcomes	No effect on glycemia or incident diabetes, blood pressure, and cardiovascular outcomes
Ferguson and Chang, <sup>111</sup> 2009	3 double-blind randomized crossover trials	41	Adults and children with cystic fibrosis	800 and 1600 IU of vitamin D alone with or without 1 g of calcium	Respiratory outcomes	No adequate evidence of benefit or harm
Abba et al, <sup>112</sup> 2008	12 trials	3393	Patients with tuberculosis	Several vitamins and minerals and diets	Promote the recovery of tuberculosis	No effect on number of deaths or number of participants with positive sputum test results
Auter and Gandini, <sup>113</sup> 2007	18 independent RCTs	57,311	At risk for dying of any cause	Vitamin D supplements varied from 300 to 2000 IU/d	Any health condition	Not enough evidence for effective decision

BMC = bone mineral content; BMD = bone mineral density; CKD = chronic kidney disease; HIV = human immunodeficiency virus; NA = not available; PTH = parathyroid hormone; RCT = randomized controlled trial; 25(OH)D = 25-hydroxyvitamin D.

inducing cellular maturation, inducing apoptosis, and inhibiting angiogenesis while enhancing the expression of genes including *P21* and *P27* to control cellular proliferation (Figure 1).<sup>1,2,7,16,26</sup> Another vitamin D–regulated gene is *CYP3A4*, whose protein product detoxifies the bile acid lithocholic acid.<sup>114</sup> Lithocholic acid is believed to damage the DNA of intestinal cells, and it may promote colon carcinogenesis. Stimulating the production of a detoxifying enzyme by 1,25(OH)<sub>2</sub>D could explain a protective role for improving vitamin D status against colon cancer.<sup>114</sup> Because vitamin D regulates a gamut of physiologic processes, including immune modulation, resistance to oxidative stress, and modulation of other hormones, it is not surprising that low vitamin D status has been associated with increased risk of several cancers and cancer mortality.<sup>7,61,115-126</sup> As the importance of non-coding RNAs has emerged, the ability of 1,25(OH)<sub>2</sub>D to regulate microRNAs (miRNAs) has been found in several cancer cell lines, patient tissues, and sera. In vitamin D<sub>3</sub> intervention trials, significant differences in miRNAs were observed between treatment groups or between baseline and follow-up.<sup>116</sup> In patient sera from population studies, specific miRNA differences were associated with serum levels of 25(OH)D. The findings thus far indicate that increasing vitamin D<sub>3</sub> intake in patients and 1,25(OH)<sub>2</sub>D<sub>3</sub> in vitro not only regulates specific miRNA(s) but also up-regulates global miRNA levels.<sup>116</sup>

Epidemiologic studies have suggested that adequate levels of 25(OH)D are critical for the prevention of various solid tumors, including prostate, breast, ovarian, and colon cancers.<sup>97,114,115,117-120</sup> A meta-analysis for the US Preventive Services Task Force regarding vitamin D supplementation concluded that each 4-ng/mL increase in blood 25(OH)D levels was associated with a 6% reduced risk of colorectal cancer but not with statistically significant dose-response relationships for prostate and breast cancer.<sup>97</sup> In a large prospective study of lethal prostate cancer (1260 cases vs 1331 controls), men with the highest quartile of plasma 25(OH)D levels had less than half the risk of lethal prostate cancer compared with men with the lowest quartile of plasma 25(OH)D levels.<sup>115</sup> A meta-analysis including 1822 colon and 868 rectal cancers reported an inverse association

between circulating 25(OH)D levels and colorectal cancer, with a stronger association for rectal cancer.<sup>97,121</sup> Participants in the WHI who had a baseline 25(OH)D level less than 12 ng/mL and who took 400 IU of vitamin D<sub>3</sub> and 1000 mg of calcium daily had a 253% increased risk of colorectal cancer compared with women who took the same amount of vitamin D<sub>3</sub> and calcium for 7 years and had baseline serum 25(OH)D levels greater than 24 ng/mL.<sup>84,122</sup>

Although cross-sectional data have many limitations, the findings are hypothesis generating and can be used to develop protocols for RCTs.<sup>84,123,124</sup> The findings from prospective case-control cohort studies in which blood collection occurred many years before diagnosis add another dimension to the evidence.<sup>118</sup> The results of these studies generally support vitamin D supplementation in those with "low" vitamin D status. However, some have argued for caution before increasing 25(OH)D levels and associated dosing regimens beyond quantities clearly supported by RCTs and meta-analyses.<sup>7,97,103</sup> There are now several observational studies reporting a U- or J-shaped association between some cancers and serum 25(OH)D and latitude or UV-B radiation levels, in which those in the highest percentiles have an inverse risk compared with those in the lowest.<sup>118,125-127</sup> Many RCTs that were evaluated for nonskeletal benefits of vitamin D had problems with a high incidence of nonadherence, misinterpretation of the original data, and use of doses of vitamin D below the 2010 IOM recommendations.<sup>62,123,124,128</sup> A good example is the WHI.<sup>129</sup> More than 50% of participants in the WHI admitted not taking their calcium and vitamin D daily, and blood concentrations of 25(OH)D were often not measured at baseline or at study end.<sup>124,129</sup> Furthermore, the authors acknowledged that the 400 IU of vitamin D was inadequate to raise the blood level of 25(OH)D above 30 ng/mL, which most studies have suggested is required to reduce cancer risk and other nonskeletal acute and chronic diseases.<sup>7,26,61,127,128</sup> A reanalysis of the WHI concluded that in 15,646 women (43%) who were not taking personal calcium or vitamin D supplements at randomization, the calcium and vitamin D intervention significantly decreased the risk of total, breast, and invasive breast cancers by 14% to 20% and

the risk of colorectal cancer by 17%.<sup>84</sup> In another RCT, a 60% reduction in all cancers was observed in postmenopausal women who ingested 1100 IU of vitamin D<sub>3</sub> and 1500 mg of elemental calcium daily for 4 years.<sup>130</sup> There is conflicting evidence about vitamin D's relationship with risk of pancreatic cancer. A study of more than 120,000 men and women from the Health Professionals Follow-up Study and the Nurses' Health Study found that participants with higher dietary intake of vitamin D had a progressively lower risk of pancreatic cancer compared with those who had the lowest intake.<sup>131</sup> In a study of men and women enrolled in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial,<sup>132</sup> no association between serum 25(OH)D levels and pancreatic cancer risk was observed. A pooled analysis of 5 nested case-control studies reported an inverse association between plasma levels of 25(OH)D and the subsequent risk of pancreatic cancer.<sup>133</sup> Compared with individuals with 25(OH)D levels less than 20 ng/mL, those with 25(OH)D levels of 20 ng/mL or greater experienced an approximately 30% lower risk of pancreatic cancer.

Grant<sup>120,134,135</sup> reported that more than 13 cancers were reduced by adequate exposure to solar UV-B radiation. He calculated that in a span of 24 years (1970-1994), 566,400 Americans died of cancer because of inadequate exposure to solar UV-B radiation.<sup>120,134-136</sup> He also estimated that 50,000 to 63,000 Americans and 19,000 to 25,000 British citizens in the United Kingdom die prematurely of cancer each year due to vitamin D deficiency.<sup>135,136</sup> A large collaborative effort analyzed data from 10 prospective cohort studies to examine whether serum 25(OH)D levels were associated with 7 rare cancers.<sup>137</sup> The National Cancer Institute Cohort Consortium Vitamin D Pooling Project of Rarer Cancers included information on serum 25(OH)D levels and the incidence of rare cancers in a subset of more than 12,000 men and women. The researchers matched participants on date and season of blood collection and used other statistical techniques to adjust for seasonal variation in serum 25(OH)D levels. When the data from the different studies were pooled, there was no overall association between vitamin D status and risk of non-Hodgkin lymphoma or cancers of the endometrium, esophagus, stomach,

kidney, or ovary.<sup>137</sup> In contrast, a recent review of ecological studies associating solar UV-B exposure—vitamin D and cancers found strong inverse correlations with solar UV-B irradiance for 15 types of neoplasms: bladder, breast, cervical, colon, endometrial, esophageal, gastric, lung, ovarian, pancreatic, rectal, renal, and vulvar cancers and Hodgkin and non-Hodgkin lymphoma.<sup>138</sup> Weaker evidence was observed for 9 other types of cancer: brain, gallbladder, laryngeal, oral/pharyngeal, prostate, and thyroid cancers; leukemia; melanoma; and myeloma.<sup>138</sup> Although there was compelling evidence for the association between vitamin D intake and cancer risk, a meta-analysis by Buttigliero et al<sup>98</sup> found no effect of vitamin D supplementation on survival in patients with cancer.

### Mortality

Vitamin D deficiency is associated with an increased risk of total mortality.<sup>139</sup> Most, but not all, studies documented increased mortality rates in patients with low 25(OH)D concentrations. In a study of 247,574 individuals from the primary care sector, a reverse J-shaped relation was reported between serum level of 25(OH)D and all-cause mortality, with the lowest mortality rate at 20 to 24 ng/mL.<sup>126</sup> This finding underscores the importance of not only including the very low (4 ng/mL) but also the higher (56 ng/mL) levels of 25(OH)D in the analysis.<sup>126</sup> It also raises several questions. How do patients who had a 25(OH)D level greater than 50 ng/mL attain such a high level, which is usually observed only in Africans living outdoors?<sup>128,140</sup> These people are likely taking megadoses of vitamin D and possibly other supplements or are being treated for vitamin D deficiency. In some instances, studies are misrepresented or misinterpreted. For example, the IOM reported in their overview that there was evidence of increased mortality for those with a 25(OH)D level greater than 30 ng/mL. However, one of the studies used to support this IOM conclusion actually stated that mortality rates were reduced until the blood level of 25(OH)D reached 50 ng/mL and that mortality rates possibly increased only in women who had 25(OH)D levels greater than 50 ng/mL.<sup>141</sup> When these J curves are plotted, we are not informed what percentage of study participants had a 25(OH)D level greater than 50 ng/mL. By some estimates, less

than 10% of patients have a 25(OH)D level greater than 50 ng/mL, and in a recent meta-analysis,<sup>126</sup> only 1.5% of participants had a 25(OH)D level greater than 50 ng/mL. This raises questions about the validity of the so-called J-U curve analyses.

A meta-analysis of prospective cohort studies including 5562 deaths of 62,548 participants suggested a nonlinear decrease in mortality risk as circulating 25(OH)D concentration increases, with optimal outcomes occurring at concentrations of approximately 30 to 35 ng/mL.<sup>142</sup> In a similar meta-analysis, vitamin D intake and blood 25(OH)D levels were inversely associated with risk of colorectal cancer, and a 10-ng/mL increase in blood 25(OH)D levels conferred a risk rate (RR) of 0.74.<sup>143</sup> A meta-analysis of prospective studies of 6853 patients with chronic kidney disease found that the mortality risk decreased by 14% per 10-ng/mL increase in 25(OH)D levels.<sup>144</sup> The major cause of mortality was cardiovascular disease.

In a recent meta-analysis with 70,528 randomized participants (86.8% females) with a median age of 70 years, vitamin D supplementation with or without calcium reduced mortality by 7%. However, vitamin D supplementation alone did not affect mortality, but risk of death was reduced if vitamin D was given with calcium.<sup>145</sup> The Ludwigshafen Risk and Cardiovascular Health Study is a cohort study of patients referred for coronary angiography between 1997 and 2000, from which 1801 with the metabolic syndrome were investigated. Mortality was tracked for a median of 7.7 years.<sup>146</sup> Multivariable survival analysis was used to estimate the association between serum 25(OH)D levels and mortality. After full adjustment, including the metabolic syndrome components, patients with optimal 25(OH)D levels had a substantial reduction in all-cause (hazard ratio [HR], 0.25; 95% CI, 0.13-0.46) and cardiovascular disease (HR, 0.33; 95% CI, 0.16-0.66) mortality rates compared with those with severe vitamin D deficiency. For specific cardiovascular disease mortality, there was a strong reduction in sudden death (HR, 0.15; 95% CI, 0.04-0.63) and congestive heart failure (HR, 0.24; 95% CI, 0.06-1.04) but not for myocardial infarction. The reduction in the mortality rate was dose dependent for each of these causes.<sup>146</sup> Consistent with the beneficial effect of vitamin D on risk of mortality, a meta-analysis of 50

randomized trials by Bjelakovic et al<sup>99</sup> found a positive effect of vitamin D<sub>3</sub> on mortality.

### Cardiovascular Disorders and Type 2 Diabetes Mellitus

Observational studies in humans found that 25(OH)D and 1,25(OH)<sub>2</sub>D levels are inversely related to coronary artery calcifications<sup>147,148</sup> and are lower in patients with myocardial infarction.<sup>149</sup> An in vitro study suggested that low 25(OH)D levels influence the activity/expression of macrophages and lymphocytes in atherosclerotic plaques, thus promoting chronic inflammation in the artery wall.<sup>150</sup> Additionally, 1,25(OH)<sub>2</sub>D<sub>3</sub> inhibited foam cell formation and promoted angiogenesis in endothelial colony-forming cells in vitro, possibly due to an increase in vascular endothelial growth factor expression and pro-matrix metalloproteinase-2 activity.<sup>4,151</sup> A short course of treatment with vitamin D (4000 IU for 5 days) effectively attenuated the increase in circulating levels of inflammatory cytokines after an acute coronary event.<sup>150</sup> These findings provide support for the anti-inflammatory effects of vitamin D on the vascular system and suggest mechanisms that mediate some of its cardioprotective properties.<sup>4,150</sup> In addition, low 25(OH)D concentrations result in elevations in PTH levels, which have been linked to insulin resistance and significant increases in the serum levels of many acute phase proteins.<sup>149</sup>

Wang et al<sup>152</sup> studied 1739 Framingham Offspring Study participants (mean age, 59 years; 55% women; all of white race) without previous cardiovascular disease. During mean follow-up of 5.4 years, 120 individuals experienced a first cardiovascular event. Individuals with 25(OH)D levels less than 15 ng/mL had a multivariable-adjusted HR of 1.62 for incident cardiovascular events compared with those with 25(OH)D levels of 15 ng/mL or greater. This effect was evident in participants with hypertension (HR, 2.13; 95% CI, 1.30-3.48) but not in those without hypertension.<sup>152</sup> Observational studies indicated that a serum 25(OH)D level less than 30 ng/mL was strongly associated with hypertension and metabolic syndrome.<sup>153</sup> This effect is thought to be partly mediated through regulation of the renin-angiotensin-aldosterone axis.<sup>154</sup> The Intermountain Heart Collaborative Study Group prospectively analyzed a large electronic medical records

database that contained 41,504 patient records. Serum 25(OH)D levels less than 30 ng/mL were associated with highly significant increases in the prevalence of diabetes, hypertension, hyperlipidemia, and peripheral vascular disease. Serum 25(OH)D levels were also highly associated with coronary artery disease, myocardial infarction, heart failure, and stroke and with incident death, heart failure, coronary artery disease/myocardial infarction, stroke, and their composite.<sup>153</sup> Black normotensive children who received 2000 IU/d of vitamin D<sub>3</sub> were compared with those who received 400 IU/d for 16 weeks in an RCT. Teenagers who received 400 IU/d of vitamin D<sub>3</sub> increased their mean  $\pm$  SD plasma levels of 25(OH)D from 13.6 $\pm$ 4.2 to 23.9 $\pm$ 7.2 ng/mL and had no reduction in arterial wall stiffness. In contrast, teenagers who received 2000 IU/d of vitamin D<sub>3</sub> increased their mean  $\pm$  SD plasma levels of 25(OH)D from 13.2 $\pm$ 3.4 to 34.2 $\pm$ 12.1 ng/mL and significantly lowered their arterial wall stiffness.<sup>155</sup> This finding is supported by the observation that serum 25(OH)D levels less than 30 ng/mL were strongly associated with hypertension, elevated blood glucose, and metabolic syndrome in adolescents.<sup>156</sup> Children with vitamin D deficiency or insufficiency had a 2.5-fold higher risk of an elevated blood glucose level, a 2.4-fold increased risk of elevated blood pressure, and a 4-fold increased risk of metabolic syndrome, a prelude to type 2 diabetes.<sup>156</sup>

A meta-analysis of 11 prospective studies involving 3612 cases and 55,713 noncase participants provided the largest and most comprehensive assessment thus far of the association between circulating 25(OH)D levels and type 2 diabetes. It suggested a strong inverse association between serum 25(OH)D concentration and incidence of type 2 diabetes. The combined RR of 0.59 suggested that the risk of future diabetes may be reduced by 41% (95% CI, 33%-48%) by having a serum 25(OH)D level greater than 32 ng/mL compared with a serum 25(OH)D level less than 19.5 ng/mL at baseline.<sup>157</sup> The MIDSPAN family study was a prospective study of 1040 men and 1298 women from the West of Scotland recruited in 1996 and followed up for a median of 14.4 years.<sup>158</sup> Plasma levels of 25(OH)D less than 15 ng/mL were not associated with a risk of cardiovascular disorders in this cohort with very low 25(OH)D levels. The median plasma 25(OH)D level was 18.6 ng/mL, and the median vitamin D intake was 3.2  $\mu$ g/d

(128 IU/d). However, there was some evidence that a 25(OH)D level less than 15 ng/mL was associated with all-cause mortality.<sup>158</sup> There was an association between 25(OH)D levels and incidence of type 2 diabetes, but there was no evidence in this study of a beneficial effect of vitamin D supplementation on type 2 diabetes outcomes.<sup>106</sup> A meta-analysis of 15 trials by George et al<sup>106</sup> did not find sufficient evidence to recommend vitamin D supplementation for improving glycemia or insulin resistance in patients with diabetes, normal fasting glucose levels, or impaired glucose tolerance. Similarly, Wang et al<sup>109</sup> and Pittas et al<sup>110</sup> concluded in their meta-analyses that evidence from limited data suggested that vitamin D supplements at moderate to high doses may reduce the risk of cardiovascular disease,<sup>109</sup> but most studies that used lower doses found no clinically meaningful effect.<sup>110</sup>

### Autoimmune Diseases

Vitamin D has been defined as a natural immune modulator. Epidemiologic, genetic, and basic science studies indicate a potential role of vitamin D in the pathogenesis of certain systemic and organ-specific autoimmune diseases, such as type 1 diabetes mellitus, MS, rheumatoid arthritis (RA), and Crohn disease (CD).<sup>159</sup> Vitamin D's effects on the innate immune system are predominantly through the toll-like receptors and on the adaptive immune system through T-cell differentiation, particularly the T helper cell (T<sub>H</sub>) type 17 response. Because T<sub>H</sub>17 cells are critical in the pathogenesis of RA, this has led to an interest in the effects of vitamin D deficiency in RA.<sup>160</sup> Vitamin D inhibits immune reactions in general, but it enhances the transcription of endogenous antibiotics, such as cathelicidin and defensins.<sup>26,161</sup> Vitamin D suppresses autoimmune disease pathology by regulating the differentiation and activity of CD4<sup>+</sup> T cells, resulting in a more balanced T<sub>H</sub>1/T<sub>H</sub>2 response that favors less development of self-reactive T cells and autoimmunity.<sup>162</sup> The T<sub>H</sub>1-dependent autoimmune diseases, including MS, type 1 diabetes, CD, and RA, are also inhibited by 1,25(OH)<sub>2</sub>D<sub>3</sub> owing to inhibition of antigen presentation, reduced polarization of T<sub>H</sub>0 cells to T<sub>H</sub>1 cells, and reduced production of cytokines from the latter cells.<sup>161</sup> The 1,25(OH)<sub>2</sub>D<sub>3</sub> down-regulated the proinflammatory cytokine (interleukin 1 $\beta$ , interleukin 6, and tumor necrosis factor) production in human activated macrophages by significantly

decreasing the aromatase activity, especially in the presence of an estrogenic milieu, such as in RA synovial tissue.<sup>163</sup> A prospective cohort study of 29,368 women aged 55 to 69 years without a history of RA found an inverse association between vitamin D intake and RA after 11 years of follow-up.<sup>164</sup> There was a 34% reduction in the development of RA with greater vitamin D intake. Women using a multivitamin with 400 IU of vitamin D reduced their risk of RA by 40%.<sup>164</sup> Use of a high-dose vitamin D<sub>3</sub> analogue resulted in improvement of symptoms of RA in 89% of patients, with 45% of patients entertaining a complete remission.<sup>165</sup> Recent evidence has suggested a significant inverse relationship between serum 25(OH)D levels and visual analog scale scores in patients with RA.<sup>166</sup> Very low serum 25(OH)D levels ( $\leq 6$  ng/mL) were characterized by patients being positive for rheumatoid factor, a high percentage of patients with very high disease activity, and a high percentage of patients requiring treatment with at least 3 disease-modifying antirheumatic drugs.<sup>167</sup>

There is a large body of evidence linking a lack of vitamin D early in life to the development of type 1 diabetes.<sup>168</sup> Vitamin D supplementation during infancy was reported to confer partial protection against  $\beta$ -cell autoimmunity.<sup>169</sup> There is consistent evidence from observational studies for potential long-term programming effects of vitamin D supplementation on immunologic diseases, such as type 1 diabetes, MS, asthma, and allergic diseases.<sup>5</sup> There was a 63% decreased risk of islet cell antibodies in offspring with a single standard deviation (156 IU) increase in recalled maternal dietary vitamin D intake during pregnancy.<sup>5</sup> Similarly, higher maternal cod liver oil (a source of vitamin D) intake during pregnancy was associated with a decreased risk of type 1 diabetes in offspring, and fetal exposure to vitamin D deficiency was linked to a higher metabolic and cardiovascular disease risk in adult life.<sup>170</sup>

A Finnish study (10,366 children) found that children who regularly took the recommended dose of vitamin D (2000 IU/d) had a rate ratio of 0.22 (95% CI, 0.05-0.89) compared with those who regularly received less than the recommended amount.<sup>171</sup> The 1,25(OH)<sub>2</sub>D<sub>3</sub> has been reported in animal models and in cultured cells to improve insulin production, modulate T- and  $\beta$ -cell activity, enhance phagocytic killing activity, improve vascular smooth muscle resistance, and

reduce the risk of autoimmune diseases.<sup>7,155,162</sup> In contrast, in healthy youth (aged 8-18 years), plasma 25(OH)D concentrations had no independent relationship with parameters of glucose homeostasis and in vivo insulin sensitivity and  $\beta$ -cell function relative to insulin sensitivity.<sup>172</sup> It remains to be determined whether in youth with dysglycemia the relationships are different and whether vitamin D optimization enhances insulin sensitivity and  $\beta$ -cell function.<sup>172</sup>

Evidence continues to accumulate supporting a protective role for vitamin D in MS risk and progression. Notable recent findings are that high 25(OH)D levels at the time of a first demyelinating event predicts a lower MS risk and a decreased risk of MS in offspring whose mothers had high 25(OH)D levels.<sup>173</sup> An American study of more than 187,000 women followed up for 10 to 20 years reported promising results with women taking at least 400 IU of supplemental vitamin D daily. The risk of MS was decreased by 41%.<sup>174</sup> An epigenetic study in lymphoblastoid cell lines reported relevant insights into how vitamin D may influence the immune system and the risk of MS through VDR interactions with the chromatin state inside MS-associated genomic regions.<sup>175</sup> Higher 25(OH)D levels were associated with decreased exacerbation risk in relapsing-remitting MS.<sup>175</sup> However, the literature is limited by small study sizes, heterogeneity of dosing, form of vitamin D tested, and clinical outcome measures.<sup>176</sup> Whether vitamin D<sub>3</sub> immunomodulatory effects can be translated into clinical benefits in patients with MS is still a matter of debate.<sup>176</sup> High doses of vitamin D<sub>3</sub> (up to 280,000 IU/wk for 6 weeks) have been used safely in patients with MS.<sup>177,178</sup> Blood levels of 25(OH)D rose to a mean of 154 ng/mL without causing hypercalcemia. The progression and activity of MS were not affected in this study, but the number of gadolinium-enhancing lesions per patient assessed by nuclear magnetic brain scan was significantly reduced.<sup>177</sup> A trial using high-dose vitamin D<sub>2</sub> to achieve 25(OH)D levels of 52 to 78 ng/mL did not reduce magnetic resonance imaging lesions in relapsing-remitting MS.<sup>178</sup> In a trial using escalating doses up to 40,000 IU/d of vitamin D<sub>3</sub> for 28 weeks followed by 10,000 IU/d for 12 weeks, there were no significant adverse events, and there seemed to be significantly less progression of disability in the treatment group.<sup>179</sup>

A chromatin immunoprecipitation sequencing—defined genome-wide map of VDR binding reported that there were 2776 “binding sites” on the human genome with at least 229 genes associated with type 1 diabetes and CD.<sup>180</sup> In a clinical trial in patients with CD in remission, 1200 IU of vitamin D<sub>3</sub> daily increased mean  $\pm$  SD serum 25(OH)D levels from 27.6 $\pm$ 12.4 to 38.4 $\pm$ 10.8 ng/mL after 3 months.<sup>181</sup> The relapse rate was numerically lower in patients treated with vitamin D<sub>3</sub> (6 of 46 or 13%) than in patients treated with placebo (14 of 48 or 29%), although this did not quite reach significance ( $P=.06$ ). Monocyte-derived dendritic cells (DCs) from 20 patients with CD were cultured with either 25(OH)D<sub>3</sub> or 1,25(OH)<sub>2</sub>D<sub>3</sub> and were matured with lipopolysaccharide (LPS).<sup>182</sup> After stimulation with 25(OH)D<sub>3</sub>, DCs from patients with CD displayed a reduced response to LPS with a diminished capability to activate T cells compared with DCs stimulated with LPS alone. Compared with LPS alone, both metabolites of vitamin D<sub>3</sub> reduced the ability of DCs to activate lymphocytes. These data indicate that intrinsic activation of 25(OH)D<sub>3</sub> to 1,25(OH)<sub>2</sub>D<sub>3</sub> occurs in DCs from patients with CD and provides evidence that higher serum 25(OH)D<sub>3</sub> levels can potentially modulate DC function in CD.<sup>182</sup> Although several studies reported the immunomodulatory effects of vitamin D on biological functions and developing processes of autoimmune diseases, there is no strong evidence for recommending vitamin D supplementation to prevent or manage the autoimmune diseases on the basis of the results of some short-term clinical trials.<sup>177,178,181</sup>

### Respiratory Tract Diseases and Wheezing Disorders

At the turn of the past century, children with rickets were at higher risk for upper respiratory tract infections and for dying of them.<sup>26,183</sup> Macrophages have a VDR, and when they ingest an infectious agent, such as tuberculosis bacillus, the toll-like receptors are activated, resulting in signal transduction to increase the expression of VDR and *CYP27B1*.<sup>7,26,28</sup> In turn, 25(OH)D is converted to 1,25(OH)<sub>2</sub>D, which signals the nucleus to increase the expression of cathelicidin, a defensin protein that kills infective agents, such as tuberculosis bacillus.<sup>7,26,28</sup>

Cord blood 25(OH)D levels were associated with tolerogenic immune regulation and fewer respiratory tract infections in newborns.<sup>184</sup> Also, high 25(OH)D levels during maternity were associated with a decrease in childhood wheezing by nearly 50% compared with low maternal 25(OH)D levels. Newborns with 25(OH)D levels less than 10 ng/mL were twice as likely to develop respiratory tract infections compared with those with levels of 30 ng/mL or greater, and every 4-ng/mL increase in the cord blood 25(OH)D level lowered the cumulative risk of wheezing by age 5 years.<sup>184</sup> Serum concentrations of 25(OH)D in 198 healthy adults revealed that a concentration of 38 ng/mL or higher reduced the risk of acute viral respiratory tract infections and number of days ill by 2-fold.<sup>185</sup> Japanese children who received 1200 IU/d of vitamin D from December through March compared with those who received placebo reduced their risk of influenza A by 42%.<sup>186</sup> It was also observed that children who took vitamin D daily had a relative risk reduction of 93% for having an asthma attack compared with children who did not take a vitamin D supplement.<sup>186</sup> Vitamin D has also been implicated in the reversal of corticosteroid resistance and in airway remodeling, which are the hallmarks of chronic obstructive pulmonary disease and severe asthma. Dietary vitamin D may regulate epigenetic events, in particular on genes that are responsible for chronic obstructive pulmonary disease susceptibility.<sup>187</sup>

The potential role of vitamin D in reducing the risk of allergies also may be related to epigenetic regulation.<sup>188,189</sup> Misdirected epigenetic programming offered an explanation for why vitamin D deficiency in pregnancy may be associated with increased allergy rates in the offspring. The cord blood level of 25(OH)D found a U-shaped association, with a 2.4-fold odds ratio (OR) of low and a 4-fold OR of high levels of 25(OH)D to develop allergen-specific IgE.<sup>188,190</sup> Eczema was significantly more likely in those with 25(OH)D levels less than 20 ng/mL compared with those with 25(OH)D levels of 30 ng/mL or greater (OR, 2.66; 95% CI, 1.24-5.72;  $P=.01$ ).<sup>189</sup> On a molecular level, maternal vitamin D intake during pregnancy increased the messenger RNA levels of immunoglobulin-like transcript (ILT) 3 and ILT4 in umbilical cord blood.<sup>191</sup> Because ILT3 and ILT4 are critical for the generation of T suppressor cells

and the induction of immunologic tolerance, this finding may point toward an early induction of tolerogenic immune responses by maternal vitamin D intake in the developing child. In addition, vitamin D stimulates natural killer cells that are known to play an immunoregulatory role in the prevention of autoimmune diseases.<sup>2</sup> Thus, although vitamin D can favorably influence several pathways associated with respiratory tract diseases, there are few clinical trials to support the beneficial effect of vitamin D supplementation for these patients. Meta-analyses on respiratory outcomes<sup>111</sup> and recovery from tuberculosis<sup>112</sup> did not report a beneficial effect of supplementation for patients with cystic fibrosis or tuberculosis, respectively.

### Neurologic Disorders

The brain has a VDR and has the ability to produce 1,25(OH)<sub>2</sub>D<sub>3</sub>. In vivo mouse studies found that in utero hypovitaminosis D impairs brain development and leads to persistent changes in the adult brain.<sup>192</sup> The 1,25(OH)<sub>2</sub>D<sub>3</sub> is rapidly incorporated into embryonic hippocampal cells, moves into the nucleus, and then returns to the cytoplasm.<sup>193</sup> These events delay cell proliferation and induce cell differentiation characterized by the expression of differentiation markers, modification of soma lengthening, and increase in neurite length and branching.<sup>193</sup> At birth, rats with prenatal vitamin D deficiency had heavier and longer brains, enlarged lateral ventricles, and decreased cortical thickness.<sup>5,192,194-196</sup> Evidence from human studies is scanty. One recent study found that higher maternal serum 25(OH)D levels in late pregnancy (<12 vs >30 ng/mL) were associated with larger head circumference of offspring at 9 years old but not with measures of cognition or psychological health.<sup>5,194</sup> In addition, there may be a critical window during late gestation in which vitamin D insufficiency precipitates an altered adult behavioral phenotype.<sup>195</sup> In rats, offspring of vitamin D-deficient mothers had significant impairment of latent inhibition (ability to ignore irrelevant stimuli), a feature often associated with schizophrenia, whereas those transiently depleted had subtle and discrete alterations in learning and memory.<sup>196</sup> In a Finnish birth cohort study, 9114 individuals were drawn from the northern Finland 1966 birth cohort.<sup>197</sup> In males, the use of at least 2000 IU of vitamin D during the first year of life was associated with a

reduced risk of schizophrenia (RR, 0.23; 95% CI, 0.06-0.95) compared with those taking lower doses.<sup>197</sup>

There is minimal evidence for an association of low maternal vitamin D status with risk of autism.<sup>5</sup> Children of dark-skinned mothers, particularly immigrants to locations with low ambient UV radiation, such as Minnesota, may be at increased risk, but this finding has been inconsistent.<sup>198</sup>

The 1,25(OH)<sub>2</sub>D<sub>3</sub> seems to have a neuroprotective role, inducing remyelination by endogenous progenitor cells and stimulation of amyloid- $\beta$  clearance by macrophages of patients with Alzheimer disease.<sup>199</sup> A vitamin D<sub>3</sub>-enriched diet correlated with a decrease in the number of amyloid plaques and inflammation in the brains of A $\beta$ PP mice.<sup>199</sup> These observations suggest that a vitamin D<sub>3</sub>-enriched diet may reduce the risk of Alzheimer disease as well as depression and neurocognitive disorders. An Australian study of 743 white pregnant women found that maternal vitamin D insufficiency during pregnancy is significantly associated with offspring language impairment.<sup>200</sup> Vitamin D deficiency was also associated with prominent changes in behavior and brain neurochemistry in the adult mouse.<sup>201</sup> In the follow-up of a British birth cohort (n=7401), current and subsequent risk of depression in middle adulthood was associated with low serum 25(OH)D levels.<sup>202</sup> This study provides support for a lower risk of depression with serum 25(OH)D levels between 20 and 34 ng/mL. A meta-analysis of cohort studies reported that there was a significantly increased HR of depression for the lowest vs highest vitamin D categories (HR, 2.21; 95% CI, 1.40-3.49).<sup>203</sup> In a community setting, depressed adults had significantly lower 25(OH)D levels than those without depression.<sup>204</sup> A variety of studies found an association between a low level of 25(OH)D and a high depression score.<sup>205,206</sup> Patients who received 400 to 800 IU of vitamin D with calcium for 6 to 12 months did not have an improvement in their mental health scores. However, patients who received 400 to 800 IU of vitamin D for 5 days with calcium or a single 100,000-IU dose of vitamin D had an improvement in the assessments of depression.<sup>207,208</sup> Although there is a strong association between risk of neurologic disorders and serum 25(OH)D concentrations, there are

only a few short-term clinical trials of vitamin D in patients with MS that have not reported benefit and no clinical trials evaluating other neurologic disorders.<sup>177,178</sup>

### Adverse Pregnancy Outcomes

A recent meta-analysis of data from 24 studies found that women with circulating 25(OH)D levels less than 20 ng/mL in pregnancy experienced an increased risk of preeclampsia (OR, 2.09; 95% CI, 1.50-2.90), gestational diabetes mellitus (OR, 1.38; 95% CI, 1.12-1.70), preterm birth (OR, 1.58; 95% CI, 1.08-2.31), and small-for-gestational-age (OR, 1.52; 95% CI, 1.08-2.15).<sup>209</sup> However, many of these outcomes are rare and require a large sample size to study, representing a challenge for cohorts with a limited number of preserved samples. Experimental studies have provided evidence of disrupted vitamin D metabolic homeostasis in the preeclamptic placenta and have suggested that increased oxidative stress could be a causative factor of altered vitamin D metabolism in preeclamptic placentas.<sup>50</sup> In normal placenta, DBP, *CYP24A1*, and VDR expressions were localized mainly in trophoblasts, whereas *CYP2R1* and *CYP27B1* expressions were localized mainly in villous core fetal vessel endothelium.<sup>50</sup> Protein expression of *CYP2R1* and VDR were reduced, but *CYP27B1* and *CYP24A1* expressions were elevated in preeclamptic compared with normotensive placentas.<sup>50</sup> A similar pattern was observed in an in vitro model that found that hypoxia induced down-regulation of DBP, *CYP2R1*, and VDR and up-regulation of *CYP27B1* and *CYP24A1*.<sup>50</sup> These data indicate that fetal (trophoblastic) autocrine synthesis of 1,25(OH)<sub>2</sub>D<sub>3</sub> may play a pivotal role in controlling placental inflammation and preeclampsia.

One of the main pathogenic features of preeclampsia is maternal endothelial dysfunction that results from impaired angiogenesis and reduced endothelial repair capacity. The 1,25(OH)<sub>2</sub>D<sub>3</sub> improves the angiogenic properties of endothelial progenitor cells. These findings could explain the positive influence of vitamin D<sub>3</sub> in reducing preeclampsia risk.<sup>151</sup>

There was an inverse association with having a cesarean delivery and serum 25(OH)D levels. In a case-control study, after adjustment for race, age, educational level, insurance status, and alcohol use, women with 25(OH)D levels less than 15 ng/mL were

almost 4 times as likely to have a cesarean delivery than were women with 25(OH)D levels of at least 15 ng/mL.<sup>210</sup>

A meta-analysis of 3 trials involving 463 women suggested that women who received vitamin D supplements during pregnancy less frequently had a baby with a birth weight less than 2500 g than did those who received no treatment or placebo; the statistical significance was borderline.<sup>89</sup> In terms of other conditions, there were no significant differences in adverse effects, including nephrotic syndrome, stillbirths, and neonatal deaths, between women who received vitamin D supplements and women who received no treatment or placebo.<sup>89</sup> A meta-analysis indicated a significant inverse relation between serum 25(OH)D level and the incidence of gestational diabetes mellitus. Overall, vitamin D deficiency (25(OH)D <20 ng/mL) in pregnancy was significantly related to the incidence of gestational diabetes mellitus, with an OR of 1.61.<sup>211</sup> However, it remains unclear whether this association is causal owing to the observational design of the studies. Recently, meta-analyses by Thorne-Lyman and Fawzi<sup>88</sup> and De-Regil et al<sup>89</sup> reported a similar beneficial effect of vitamin D supplementation on birth weight but no significant effect on other maternal and neonatal outcomes.

### ASSESSING VITAMIN D STATUS

Although the generally accepted measure of vitamin D status is circulating 25(OH)D concentration, there is little consensus on which assay method should be used. Commonly used assays include competitive protein-binding assay, radioimmunoassay, enzyme immunoassay, chemiluminescence immunoassay, high-performance liquid chromatography, and liquid chromatography–tandem mass spectrometry (LC-MS/MS), each with its own advantages and disadvantages.<sup>212,213</sup> Binkley et al<sup>214</sup> reported that 25(OH)D results differed widely depending on the laboratory and the method used, with the mean result (from 10 healthy adults) varying from 17.1 to 35.6 ng/mL. A study conducted by the Vitamin D External Quality Assessment Scheme found a 31% overestimation by one immunoassay method.<sup>215</sup> Its specificity needs to exclude significant interferences from the C-3 epimer of 25(OH)D, which is more prevalent in infants younger than 1 year.<sup>216</sup>

DiaSorin radioimmunoassays (DiaSorin Corp) used in the NHANES III had a mean bias of greater than 12% comparing the vitamin D status of the US population of 1988-1994 with that of 2000-2004. This difference was probably caused by changes in reagents and calibration lots performed by the manufacturer.<sup>217,218</sup> This makes diagnostic and therapeutic decisions on the basis of absolute cutoff values for vitamin D deficiency extremely problematic<sup>219,220</sup> and hinders the comparability of results from prospective and epidemiologic studies. Since November 2010, efforts have been made to recalibrate 25(OH)D measurements from all NHANES samples with LC-MS/MS. This effort is within the context of an international approach to standardization of 25(OH)D measurements in national surveys, the Vitamin D Standardization Program, and publication of the results is planned for the middle of 2013.<sup>221</sup> To ensure that laboratories are providing accurate testing results, it is important that their vitamin D testing method measures total 25(OH)D levels [25(OH)D<sub>2</sub> and 25(OH)D<sub>3</sub>] and has acceptable precision. The immunoassay remains the predominant mode of measurement for 25(OH)D. Most, if not all, of these assays have problems with equimolar recovery of the 25(OH)D<sub>2</sub> and 25(OH)D<sub>3</sub> levels.<sup>222</sup> The level of 25(OH)D<sub>2</sub> is underestimated by 20% to 80% and can vary for different patients who received vitamin D<sub>2</sub> in the same assay. Standardization of all assays has been improved but not resolved with the currently available reference materials, as evidenced by the Vitamin D External Quality Assessment Scheme.<sup>222</sup> The choice of method for each laboratory remains a balance mainly among turnaround time, convenience, cost, and the specificity and accuracy of the information obtained.<sup>208</sup> Recognizing the importance of a 25(OH)D<sub>2</sub> and 25(OH)D<sub>3</sub> reference material, the National Institute of Standards and Technology released a 4-level Standard Reference Material set, SRM972.<sup>221</sup> Treatment with vitamin D<sub>2</sub>, therefore, may not be accurately monitored using many of the commercial assays because these antibody assays often underestimate 25(OH)D<sub>2</sub> levels and, thus, total 25(OH)D levels, which is what they record in the report. This issue is most important in patients who experience no improvement with replacement therapy with vitamin D<sub>2</sub> (the most

commonly used Food and Drug Administration—approved pharmaceutical form of vitamin D); the absence of a rise in the total 25(OH)D level may indicate nonadherence or malabsorption.<sup>223</sup> For these patients, use of the gold standard LC-MS/MS would quantitatively report 25(OH)D<sub>2</sub>, 25(OH)D<sub>3</sub>, and total 25(OH)D levels.

#### VITAMIN D STATUS DURING PREGNANCY, BIRTH, AND CHILDHOOD

Maternal vitamin D deficiency predisposes to low vitamin D stores in the newborn and increases infantile rickets<sup>224</sup> because the mother is the only source of vitamin D during pregnancy. The prevalence of vitamin D deficiency and insufficiency during pregnancy is of special concern and ranges from 8% to 100%, depending on the country of residence and the definitions of vitamin D deficiency and insufficiency (Figure 5).<sup>2</sup> In the United States, vitamin D deficiency and insufficiency is estimated to be 27% to 91% in pregnant women.<sup>2</sup> As shown in Figure 5, this rate is estimated to be 36% to 65% in Canada, 45% to 98% in Asia, 70% to 100% in Europe, and 25% to 65% in Australia and New Zealand.<sup>124,128</sup>

The prevalence of vitamin D deficiency and insufficiency in children in China is high, especially in children aged 6 to 16 years.<sup>211</sup> In the United States, it is estimated that 50% of children aged 1 to 5 years and 70% of children aged 6 to 11 years are vitamin D deficient or insufficient.<sup>156</sup> Recent studies reported that adolescents and young adults are also at risk for vitamin D deficiency.<sup>16,128,217</sup> Also, a high prevalence of vitamin D deficiency was reported in a cross-sectional study conducted at a tertiary care center in western India.<sup>225</sup>

Evidence suggests that children and adults in the United States are becoming more vitamin D deficient and insufficient because of an increase in the incidence of obesity, a decrease in milk consumption, and an increase in sun protection.<sup>15,217</sup> This recent evidence emphasizes the high prevalence of vitamin D deficiency throughout the world, not only in at-risk groups (Figure 5).<sup>7,15,217,224-230</sup>

#### PREVALENCE OF VITAMIN D DEFICIENCY IN ADOLESCENTS AND ADULTS

It has been estimated that 20% to 80% of US, Canadian, and European men and women are vitamin D deficient.<sup>228,229</sup> The prevalence of


serum 25(OH)D levels less than 20 ng/mL was almost one-third of the US population (32%).<sup>217</sup> More than 70% of non-Hispanic black individuals and more than 40% of Hispanic/Mexican individuals were at risk for a 25(OH)D level less than 20 ng/mL.<sup>228</sup> In a national Canadian cohort, serum 25(OH)D levels less than 30 ng/mL were evident in 57.5% of men and in 60.7% of women, and they rose to 73.5% in spring (men) and 77.5% in winter (women).<sup>229</sup> In the Healthy Lifestyle in Europe by Nutrition in Adolescence study, 25(OH)D levels less than 30 ng/mL were reported to be approximately 80% in adolescents from the 9 European countries.<sup>230</sup> Levels of 25(OH)D were higher in northern Europe than in southern Europe and were higher in western Europe than in eastern Europe.<sup>230</sup> The higher levels in northern Europe were also observed in some multicenter studies in which a single laboratory facility was used.<sup>230</sup>

The higher serum 25(OH)D levels in Norway and Sweden are probably due to a high intake of fatty fish and cod liver oil. The lower serum 25(OH)D levels in Spain, Italy, and Greece may be due to more skin pigmentation, sunshine-avoiding behavior, and air pollution with ozone and nitrogen dioxide, which reduce sun-induced vitamin D production.<sup>226</sup>

In the Middle East and Asia, vitamin D deficiency in children and adults is highly prevalent.<sup>7,227</sup> Children and adults of color are especially at high risk owing to the inefficient cutaneous production of vitamin D<sub>3</sub>.<sup>1,2,7</sup> In a study on the vitamin D status of Australian adults, vitamin D deficiency (25[OH]D <20 ng/mL) was 31% (22% in men and 39% in women); 73% had 25(OH)D levels less than 30 ng/mL.<sup>226</sup> Women who practice purdah (ie, the use of clothing and other approaches to screen themselves from men and strangers) and children and adults who avoid all sun exposure or wear sunscreen protection are equally at high risk.<sup>7,231</sup>

#### CAUSES OF VITAMIN D DEFICIENCY AND RISK FACTORS

Traditional risk groups for vitamin D deficiency include pregnant women, children, older persons, the institutionalized, and non-Western immigrants.<sup>7,228</sup> The major source of vitamin D for children and adults is exposure to natural sunlight.<sup>7,61</sup> The Maasai and




Hadzabe tribes in Tanzania (East Africa) with traditional lifestyles, living in the presumed cradle of humankind, who are exposed daily to tropical sunlight had a mean circulating 25(OH)D level of 46 ng/mL.<sup>140</sup>

A variety of factors influence the cutaneous production of vitamin D. A sunscreen with a sun protection factor of 30 applied properly reduces the ability of the skin to produce vitamin D by as much as 95% to 99%. People of color who have natural sunscreen protection from their increased melanin pigment are less efficient by more than 90% in producing vitamin D in their skin compared with white individuals.<sup>232</sup> In addition, air pollution with increased ozone and nitrogen dioxide levels (both known to compromise several health outcomes) absorbs UV-B radiation and is an often-neglected risk factor for hypovitaminosis D.<sup>61,233</sup> Important risk factors for vitamin D deficiency are shown in Figure 6.

The prevalence of vitamin D deficiency and insufficiency is affected by seasonal variation and latitude. The prevalence increases in late

winter/spring and decreases in summer.<sup>234</sup> A study of the effect of education on vitamin D status found that low-educated women had lower 25(OH)D levels compared with high-educated women, and women in the lowest 25(OH)D quartile had a higher risk of small-for-gestational-age offspring.<sup>235</sup>

The elderly population is particularly at risk for clinical complications related to low 25(OH)D levels. With increasing age, solar exposure is usually limited because of changes in lifestyle factors, such as clothing and less outdoor activity. Diet may also become less varied, with a lower natural vitamin D content. Most important, however, the cutaneous production of vitamin D after exposure to solar UV-B radiation decreases with age because of atrophic skin changes, with a reduced amount of its precursor 7-DHC.<sup>236,237</sup> A comparison of the amount of previtamin D<sub>3</sub> produced in skin from 8- to 18-year-old individuals with the amount produced in skin from 77- to 82-year-old individuals revealed that aging can decrease by greater than 2-fold the capacity of the skin to produce previtamin D<sub>3</sub>.<sup>237</sup>



Although the heritability of vitamin D status seems considerable, the specific genetic determinants of 25(OH)D levels are only beginning to be identified. A recent examination of 141 single nucleotide polymorphisms (SNPs) in a discovery cohort of 1514 white participants from the community-based Cardiovascular Health Study found that lower serum 25(OH)D levels were associated with HRs for the risk of the composite outcome of 1.40 for those who had 1 minor allele at rs7968585 (in VDR) and 1.82 for those with 2 minor alleles.<sup>238,239</sup> This candidate gene study indicates that known associations of low serum 25(OH)D levels with clinical outcomes may vary according to genetic differences in the VDR. In black patients, there

were significant associations in 3 SNPs in vitamin D pathway genes (rs2282679, rs2298849, and rs10877012), all of which replicate earlier findings in populations of European ancestry.<sup>239</sup> Included among these was rs2282679, a highly significant result from 2 recent genome-wide association studies (GWASs),<sup>240,241</sup> one of which reported a 49% increased risk of vitamin D deficiency (25[OH]D <20 ng/mL) associated with the rs2282679 minor allele in white individuals.<sup>241</sup> Another study of genetic predictors of 25(OH)D in black individuals involved 513 participants from 42 families in Los Angeles, California, and evaluated 30 SNPs in DBP, VDR, and CYP27B1.<sup>242</sup> Recent epigenomic findings confirmed 3 genes (*DHCR7*,

*CYP2R1*, and *CYP24A1*) of the 4 genes in the GWAS findings, which reinforces the crucial roles played by those 3 genes in vitamin D metabolism.<sup>243</sup> *DHCR7* encodes the enzyme 7-DHC reductase, which converts 7-DHC to cholesterol, thereby removing the substrate from the synthetic pathway of vitamin D<sub>3</sub>.<sup>239</sup> *DHCR7* is a novel gene for association with 25(OH)D levels, as identified in 2 recent GWASs.<sup>240,241</sup> *CYP24A1*, which encodes 25(OH)D-24-hydroxylase, has been identified as a candidate gene for vitamin D insufficiency in one GWAS but not in the other.<sup>240-242</sup> This mitochondrial protein initiates the degradation of 1,25(OH)<sub>2</sub>D<sub>3</sub> and plays a role in calcium homeostasis and vitamin D metabolism. These epigenomic findings suggest that individuals with vitamin D deficiency are more likely to have reduced synthesis and increased catabolism of 25(OH)D and 1,25(OH)<sub>2</sub>D.<sup>243</sup>

The genetic contributions to circulating 25(OH)D represent a complex trait for which family studies have estimated heritability ranging from 43% to 80%.<sup>244</sup> Genomic and epigenomic data integration provided greater understanding of the physiology and etiology of the complex traits. Further elucidation of the genetic architecture of this complex trait beyond environmental determinants of 25(OH)D has the potential to identify those at risk for vitamin D insufficiency.<sup>244</sup> It may also provide a useful proxy for lifetime vitamin D exposure that may be applied in instrumental variable analyses investigating the association between vitamin D and common complex diseases. However, a recent GWAS of prospectively collected 25(OH)D data in 5 studies with 5575 individuals reported that known GWAS-associated SNPs explain only a fraction of the observed variance in circulating 25(OH)D levels (ie, approximately 5.2%).<sup>244</sup> On rare occasions, some patients who deny taking a vitamin D supplement have unexplained high normal 25(OH)D levels in the range of 40 to 80 ng/mL. It is believed that this is due to a genetic mutation of the *cyp24A1* that reduces the catabolism of 25(OH)D and 1,25(OH)<sub>2</sub>D and can be a cause of infantile hypercalcemia.<sup>245</sup> Therefore, these recent genomic and epigenetic data provide additional evidence of genetic-environmental interactions and their effects on circulating 25(OH)D levels.

#### TREATMENT AND PREVENTION OF VITAMIN D DEFICIENCY AND INSUFFICIENCY WITH SUN EXPOSURE AND UV-B IRRADIATION

Humans obtain a considerable amount of their vitamin D requirement from sun exposure.<sup>7,140</sup> Although excessive exposure to sunlight increases the risk of nonmelanoma skin cancer, which is easy to detect and easy to treat, there is no evidence that sensible sun exposure, as our hunter-gatherer forefathers likely experienced, increases risk.<sup>124,246</sup> More importantly, the most deadly form of melanoma skin cancer that occurs on the least sun-exposed areas is less likely to occur in adults who have outdoor occupations.<sup>1,124,246,247</sup> Therefore, it is not unreasonable to consider sensible sun exposure as a good source of vitamin D.<sup>7,124</sup> An adult in a bathing suit exposed to 1 minimal erythemal dose (slight pinkness to the skin 24 hours after exposure) is the equivalent to taking approximately 20,000 IU (500 µg) of vitamin D<sub>2</sub> orally.<sup>7,15</sup> Thus, exposure of arms and legs to 0.5 minimal erythemal dose is equivalent to ingesting approximately 3000 IU of vitamin D<sub>3</sub>.<sup>7,60</sup> Adults who frequented a tanning salon had robust levels of 25(OH)D, on average 46 ng/mL, and had higher bone mineral density in their hips compared with healthy adults who did not go to a tanning salon in Boston, Massachusetts, at the end of winter.<sup>248</sup> It was estimated that if all the people in the United States were to double their solar UV-B irradiance to raise their serum 25(OH)D levels to 45 ng/mL, the net result could be as many as 400,000 reduced deaths compared with only 11,000 increased deaths from melanoma and other skin cancer.<sup>249</sup> Time of day during sun exposure, season, latitude, and degree of skin pigmentation dictate how much vitamin D<sub>3</sub> is produced during sun exposure. Exposure of the arms and legs (abdomen and back when possible) to sunlight 2 to 3 times a week for approximately 25% to 50% of the time it would take to develop a mild sunburn (minimal erythemal dose) will cause the skin to produce enough vitamin D. For a white person, if 30 minutes of June noontime sun would cause a mild sunburn, then 10 to 15 minutes of exposure followed by good sun protection should be sufficient to produce adequate vitamin D.<sup>7</sup> There is no need to ever expose the face because although it is the most sun exposed of all the body areas, it provides little vitamin D<sub>3</sub>. A free app, [dminder.info](http://dminder.info), provides the user with

information about sensible sun exposure and vitamin D production. For patients with fat malabsorption syndromes that render oral consumption of supplemental vitamin D ineffective, exposure to a lamp that emits UV-B radiation can be effective in raising blood levels of 25(OH)D.<sup>250</sup>

### Food Sources

Very few foods naturally contain vitamin D; examples of foods with ample vitamin D stores include wild-caught salmon and UV-exposed mushrooms.<sup>7</sup> Foods fortified with vitamin D usually contain 100 IU per serving. An analysis of the vitamin D intake of children and adults in the United States revealed that they were unable to achieve the RDA for vitamin D from any dietary sources.<sup>251</sup>

Vitamin D intake can be increased by eating foods fortified with vitamin D. A recent systematic review found that food fortification with vitamin D (especially in milk) is effective in significantly increasing 25(OH)D levels in the population.<sup>7,252</sup> Other foods include some cereals, juices, other dairy products, and some margarines. A mean individual intake of approximately 11 µg/d (440 IU/d) from fortified foods (range, 120-1000 IU/d) increased 25(OH)D concentrations by 7.7 ng/mL, corresponding to a 0.48-ng/mL increase in 25(OH)D for each 40 IU (1 µg) ingested.<sup>252</sup>

### Vitamin D Supplementation

The RDA for vitamin D and tolerable upper-limit levels vary in different age groups and in certain circumstances.<sup>7,60,62</sup> Although it is recommended that RDAs of 600 to 800 IU daily should meet the requirements to optimize bone health<sup>62</sup> in most of the population, higher vitamin D intakes (1000-2000 IU) are needed to reach and maintain 25(OH)D levels greater than 30 ng/mL.<sup>7,60</sup> It is recognized that for every 100 IU of vitamin D ingested, the blood level of 25(OH)D increases by approximately 0.6 to 1 ng/mL.<sup>253</sup> When the serum 25(OH)D level is less than 15 ng/mL, 100 IU of vitamin D will increase the 25(OH)D level by as much as 2 to 3 ng/mL.<sup>7,71</sup> An effective strategy to treat vitamin D deficiency and insufficiency in children and adults is to give them 50,000 IU of vitamin D<sub>2</sub> once a week for 6 and 8 weeks, respectively.<sup>60,254</sup> To prevent recurrence of vitamin D deficiency in children, administration of 600 to 1000 IU/d is effective.<sup>60</sup> For adults, to prevent recurrence of

vitamin D deficiency, administration of 50,000 IU of vitamin D<sub>2</sub> every 2 weeks is effective.<sup>7,60,255</sup>

This strategy was shown to be effective in maintaining blood levels of 25(OH)D at approximately 40 to 60 ng/mL for up to 6 years without any evidence of toxic effects.<sup>255</sup>

Vitamin D can be administered daily, weekly, monthly, or every 4 months to sustain an adequate serum 25(OH)D concentration.<sup>7,256-258</sup> A bolus of high doses of vitamin D (up to 300,000 IU) can be initially used in persons with extreme vitamin D deficiency. Repeated boluses of high-dose vitamin D at 6- to 12-month intervals have been used in a nursing home setting, but a steady-state serum 25(OH)D concentration is likely to be maintained by more frequent, lower doses of vitamin D. One study has suggested that a 500,000-IU bolus dose of vitamin D<sub>3</sub> increases the risk of fracture within 3 months,<sup>256</sup> but other studies have reported reduced risk of fracture.<sup>257,258</sup>

### Vitamin D Supplementation During Pregnancy and Lactation

The 25(OH)D passes from the placenta into the bloodstream of the fetus. Because the half-life for 25(OH)D is approximately 2 to 3 weeks, the infant can remain vitamin D sufficient for several weeks after birth, as long as the mother is vitamin D sufficient.<sup>2</sup>

In a study of 40 mostly black pregnant women who were documented to be ingesting approximately 600 IU of vitamin D a day, at the time that they gave birth, 76% were vitamin D deficient as defined by the IOM cutoff value of a 25(OH)D level less than 20 ng/mL.<sup>259</sup> Eighty-one percent of their newborns were vitamin D deficient.<sup>259</sup> Maternal supplementation with 2000 and 4000 IU/d of vitamin D during pregnancy improved the maternal/neonatal vitamin D status.<sup>260</sup> None of the pregnant women developed significant changes in their serum calcium or 24-hour urinary calcium levels. Evidence of risk reduction in infection, preterm labor, and preterm birth was suggestive, requiring additional studies powered for these end points.<sup>260,261</sup>

Human breast milk and unfortified cow's milk have little vitamin D.<sup>7,260,261</sup> Only after lactating women were given 4000 to 6000 IU/d of vitamin D was enough vitamin D transferred in breast milk to satisfy the infant's requirement.<sup>260,261</sup>

### Vitamin D Supplementation in Special Conditions

Because body fat can sequester vitamin D, it is now recognized that children and adults who are obese require 2 to 5 times more vitamin D to treat and prevent vitamin D deficiency.<sup>7,60</sup> Patients taking antiepileptic medications, AIDS medications, and glucocorticoids often require more vitamin D to satisfy their requirements.<sup>7,60</sup> However, patients with granulomatous disorders, such as sarcoidosis and tuberculosis, are at risk for hypercalciuria and hypercalcemia when blood levels of 25(OH)D are greater than 30 ng/mL owing to the increased serum levels of 1,25(OH)<sub>2</sub>D produced in the macrophages in the granulomas.<sup>7</sup> Therefore, their vitamin D intake needs to be carefully monitored and controlled.<sup>7,60</sup> Hence, daily requirements of vitamin D to reach and maintain the desired serum 25(OH)D level can be estimated from the baseline 25(OH)D concentration. Supplemental vitamin D is preferentially administered orally or intramuscularly (not available in the United States), and the vitamin D-producing Sperti lamp can be used, where available, in patients with malabsorption syndromes.<sup>250,262</sup>

### Type of Vitamin D Supplementation

Either vitamin D<sub>2</sub> or vitamin D<sub>3</sub> can be used for vitamin D supplementation, although there is controversy regarding vitamin D<sub>3</sub> vs vitamin D<sub>2</sub> for achieving and maintaining higher serum 25(OH)D levels. Although a recent meta-analysis indicated that vitamin D<sub>3</sub> is more efficacious at raising serum 25(OH)D concentrations than is vitamin D<sub>2</sub>,<sup>2,102</sup> several prospective studies have found them to be equally effective in raising and maintaining serum 25(OH)D levels in children and adults.<sup>263,264</sup> Holick et al<sup>264</sup> found that an 11-week course of treatment with 1000 IU/d of vitamin D<sub>2</sub>, 1000 IU/d of vitamin D<sub>3</sub>, or a combination of 500 IU/d of vitamin D<sub>2</sub> and 500 IU of vitamin D<sub>3</sub> daily caused an equivalent increase in serum total 25(OH)D levels. Furthermore, the group that received vitamin D<sub>2</sub> did not experience a significant change in serum 25(OH)D<sub>3</sub> levels. Gordon et al<sup>265</sup> and Thacher et al<sup>266</sup> also found that in infants and toddlers treated for 6 weeks, 2000 IU of vitamin D<sub>2</sub> and 2000 IU of vitamin D<sub>3</sub> daily or a single dose of 50,000 IU of vitamin D<sub>2</sub> or vitamin D<sub>3</sub> were equally effective in increasing the serum total 25(OH)D level. The

bioavailability of vitamin D<sub>3</sub> is well established, and the bioavailability of vitamin D<sub>2</sub> from mushrooms in humans has been found to be comparable with that of a vitamin D<sub>2</sub> supplement.<sup>267</sup> Finally, adults treated with vitamin D<sub>2</sub> not only raised their total blood levels of 25(OH)D but also maintained total blood levels of 1,25(OH)<sub>2</sub>D to the same degree as adults who received the same dose of vitamin D<sub>3</sub>.<sup>263</sup>

### SAFETY AND INTOXICATION

Vitamin D intoxication is characterized by hypercalcemia, hypercalciuria, and hyperphosphatemia, which, in turn, are responsible for soft-tissue and vascular calcifications and nephrolithiasis in the long term. Serum 25(OH)D levels are usually markedly elevated (>150 ng/mL) in individuals with vitamin D intoxication.<sup>7,60,90</sup> Daily doses of vitamin D<sub>3</sub> up to 10,000 IU were safe in healthy males, and there was no evidence of hypercalcemia or hypercalciuria for 5 months.<sup>253,268</sup> This amount is far above the tolerable upper level indicated in the IOM guidelines (4000 IU). Higher doses of vitamin D (up to 40,000 IU/d) are still safe provided that a serum 25(OH)D concentration of 200 ng/mL is not exceeded. A recent report of an infant inadvertently receiving 12,000 IU of vitamin D<sub>3</sub> daily for 20 days and achieving a serum 25(OH)D level of 425 ng/mL had no signs of vitamin D intoxication. Once the vitamin D use was stopped, the serum 25(OH)D level was less than 100 ng/mL within 2 months.<sup>269</sup>

### CONCLUSION

Vitamin D deficiency is a common underdiagnosed condition that has received increasing attention in the world. The US Endocrine Society guidelines and the IOM recommend screening only in populations at risk, as no evidence currently exists to support screening at a population level. Candidates for vitamin D screening include those who are at specific risk for vitamin D deficiency and patients who are experiencing or are at risk for specific medical conditions associated with hypovitaminosis D.

Recent evidence from hundreds of studies has suggested that vitamin D is important for reducing the risk of a variety of chronic illnesses. The identification of a VDR in most tissues and cells and the observation that a multitude of genes may be directly or indirectly regulated by 1,25(OH)<sub>2</sub>D have provided a

rationale for the nonskeletal health benefits of vitamin D. A study in healthy adults who received either 400 or 2000 IU/d of vitamin D<sub>3</sub> for 3 months in winter reported that 291 genes were either up-regulated or down-regulated. That these genes affected as many as 80 different metabolic pathways (from immune modulation to enhanced antioxidant activity) emphasizes the importance of improving the world's vitamin D status.<sup>17</sup> The observation that 1,25(OH)<sub>2</sub>D may also influence epigenetics provides additional support for the concept that there is no downside to increasing the vitamin D status of children and adults. Vitamin D deficiency during pregnancy may adversely influence placental development and fetal programming. Vitamin D deficiency in both parents may influence adverse pregnancy outcomes and susceptibility to developing disease in adult life and even into the next generation.

There is potentially a great upside (in terms of improving overall health and well-being) to increasing serum 25(OH)D levels above 30 ng/mL. An effective strategy to prevent vitamin D deficiency and insufficiency is to obtain some sensible sun exposure, ingest foods that contain vitamin D, and take a vitamin D supplement.

**Abbreviations and Acronyms:** CD = Crohn disease; DBP = vitamin D binding protein; DC = dendritic cell; ESCEO = European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis; GWAS = genome-wide association study; HR = hazard ratio; ILT = immunoglobulin-like transcript; IOM = Institute of Medicine; IVF = in vitro fertilization; LC-MS/MS = liquid chromatography–tandem mass spectrometry; LPS = lipopolysaccharide; miRNA = microRNA; MS = multiple sclerosis; NHANES = National Health and Nutrition Examination Survey; OR = odds ratio; PTH = parathyroid hormone; PTHrP = parathyroid hormone–related protein; RA = rheumatoid arthritis; RDA = recommended dietary allowance; RCT = randomized controlled trial; RR = risk rate; SE = standard error; SNP = single nucleotide polymorphism; T<sub>H</sub> = T helper cell; VDR = vitamin D receptor; WHI = Women's Health Initiative; 1,25(OH)<sub>2</sub>D = 1,25-dihydroxyvitamin D; 7-DHC = 7-dehydrocholesterol; 25(OH)D = 25-hydroxyvitamin D

**Grant Support:** This work was supported, in part, by grant UL-1-RR-25711 from the National Institutes of Health and by the Mushroom Council.

**Potential Competing Interests:** Dr Holick is a consultant for Quest Diagnostics, Ontometrics, Vital Choice, Merck, and Bioceuticals.

**Correspondence:** Address to Michael F. Holick, PhD, MD, Boston University School of Medicine, 85 E Newton St, M-1013, Boston, MA 02118 (mfholick@bu.edu).

## REFERENCES

- Holick MF. Vitamin D: extraskeletal health. *Rheum Dis Clin North Am*. 2012;38(1):141-160.
- Hossein-nezhad A, Holick MF. Optimize dietary intake of vitamin D: an epigenetic perspective. *Curr Opin Clin Nutr Metab Care*. 2012;15(6):567-579.
- Smit E, Crespo CJ, Michael Y, et al. The effect of vitamin D and frailty on mortality among non-institutionalized US older adults. *Eur J Clin Nutr*. 2012;66(9):1024-1028.
- Holick MF. Nutrition: D-iabetes and D-eath D-efying vitamin D. *Nat Rev Endocrinol*. 2012;8(7):388-390.
- Lucas RM, Ponsonby AL, Pasco JA, Morley R. Future health implications of prenatal and early-life vitamin D status. *Nutr Rev*. 2008;66(12):710-720.
- Brannon PM, Picciano MF. Vitamin D in pregnancy and lactation in humans. *Annu Rev Nutr*. 2011;31:89-115.
- Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357(3):266-281.
- Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr*. 2004;80(6, suppl):1678S-1688S.
- Holick MF, MacLaughlin JA, Clark MB, et al. Photosynthesis of previtamin D<sub>3</sub> in human skin and the physiologic consequences. *Science*. 1980;210(4466):203-205.
- Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D<sub>3</sub>: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D<sub>3</sub> synthesis in human skin. *J Clin Endocrinol Metab*. 1988;67(2):373-378.
- Holick MF, Chen TC, Lu Z, Sauter E. Vitamin D and skin physiology: a D-lightful story. *J Bone Miner Res*. 2007;22(suppl 2):V28-V33.
- Negri AL. Proximal tubule endocytic apparatus as the specific renal uptake mechanism for vitamin D-binding protein/25-(OH)D<sub>3</sub> complex. *Nephrology*. 2006;11(6):510-515.
- Shimada T, Kakitani M, Yamazaki Y, et al. Targeted ablation of Fgf23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *J Clin Invest*. 2004;113(4):561-568.
- Nair R, Maseeh A. Vitamin D: the "sunshine" vitamin. *J Pharmacol Pharmacother*. 2012;3(2):118-126.
- Holick MF. The D-lightful vitamin D for child health. *JPEN J Parenter Enteral Nutr*. 2012;36(1, suppl):9S-19S.
- Nagpal S, Na S, Rathnachalam R. Noncalcemic actions of vitamin D receptor ligands. *Endocr Rev*. 2005;26(5):662-687.
- Hossein-nezhad A, Spira A, Holick MF. Influence of vitamin D status and vitamin D<sub>3</sub> supplementation on genome wide expression of white blood cells: a randomized double-blind clinical trial. *PLoS One*. 2013;8(3):e58725.
- Kitazawa S, Kajimoto K, Kondo T, Kitazawa R. Vitamin D<sub>3</sub> supports osteoclastogenesis via functional vitamin D response element of human RANKL gene promoter. *J Cell Biochem*. 2003;89(4):771-777.
- Cui X, Pelekanos M, Liu PY, Bume TH, McGrath JJ, Eyles DW. The vitamin D receptor in dopamine neurons; its presence in human substantia nigra and its ontogenesis in rat midbrain. *Neuroscience*. 2013;236:77-87.
- Gonzalez-Parra E, Rojas-Rivera J, Tunon J, Praga M, Ortiz A, Egidio J. Vitamin D receptor activation and cardiovascular disease. *Nephrol Dial Transplant*. 2012;27(suppl 4):iv17-iv21.
- Nowak R, Szota J, Mazurek U. Vitamin D receptor gene (VDR) transcripts in bone, cartilage, muscles and blood and microarray analysis of vitamin D responsive genes expression in paravertebral muscles of juvenile and adolescent idiopathic scoliosis patients. *BMC Musculoskelet Disord*. 2012;13:259.
- Doig CL, Singh PK, Dhiman VK, et al. Recruitment of NCOR1 to VDR target genes is enhanced in prostate cancer cells and associates with altered DNA methylation patterns. *Carcinogenesis*. 2013;34(2):248-256.

23. Alimirah F, Peng X, Yuan L, et al. Crosstalk between the peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) and the vitamin D receptor (VDR) in human breast cancer cells: PPAR $\gamma$  binds to VDR and inhibits 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> mediated transactivation. *Exp Cell Res*. 2012;318(19):2490-2497.
24. Nagy L, Szanto A, Szatmari I, Szeles L. Nuclear hormone receptors enable macrophages and dendritic cells to sense their lipid environment and shape their immune response. *Physiol Rev*. 2012;92(2):739-789.
25. Kaludjerovic J, Vieth R. Relationship between vitamin D during perinatal development and health. *J Midwifery Womens Health*. 2010;55(6):550-560.
26. Adams JS, Hewison M. Update in vitamin D. *J Clin Endocrinol Metab*. 2010;95(2):471-478.
27. Shin JS, Choi MY, Longtine MS, Nelson DM. Vitamin D effects on pregnancy and the placenta. *Placenta*. 2010;31(12):1027-1034.
28. Adams JS, Hewison M. Extrarenal expression of the 25-hydroxyvitamin D-1-hydroxylase. *Arch Biochem Biophys*. 2012;523(1):95-102.
29. Grayson R, Hewison M. Vitamin D and human pregnancy. *Fetal Matern Med Rev*. 2011;22(1):67-90.
30. Powe CE, Seely EW, Rana S, et al. First trimester vitamin D, vitamin D binding protein, and subsequent preeclampsia. *Hypertension*. 2010;56(4):758-763.
31. Zella LA, Shevde NK, Hollis BW, Cooke NE, Pike JW. Vitamin D-binding protein influences total circulating levels of 1,25-dihydroxyvitamin D<sub>3</sub> but does not directly modulate the bioactive levels of the hormone in vivo. *Endocrinology*. 2008;149(7):3656-3667.
32. Liu NQ, Hewison M. Vitamin D, the placenta and pregnancy. *Arch Biochem Biophys*. 2012;523(1):37-47.
33. Ardawi MS, Nasrat HA, BA'Aqueel HS. Calcium-regulating hormones and parathyroid hormone-related peptide in normal human pregnancy and postpartum: a longitudinal study. *Eur J Endocrinol*. 1997;137(4):402-409.
34. Weisser J, Riemer S, Schmidt M, et al. Four distinct chondrocyte populations in the fetal bovine growth plate: highest expression levels of PTH/PTHrP receptor, Indian hedgehog, and MMP-13 in hypertrophic chondrocytes and their suppression by PTH (1-34) and PTHrP (1-40). *Exp Cell Res*. 2002;279(1):1-13.
35. Kovacs CS, Kronenberg HM. Maternal-fetal calcium and bone metabolism during pregnancy, puerperium, and lactation. *Endocr Rev*. 1997;18(6):832-872.
36. Mahadevan S, Kumaravel V, Bharath R. Calcium and bone disorders in pregnancy. *Indian J Endocrinol Metab*. 2012;16(3):358-363.
37. Kovacs CS. Vitamin D in pregnancy and lactation: maternal, fetal, and neonatal outcomes from human and animal studies. *Am J Clin Nutr*. 2008;88(2):520S-528S.
38. Kovacs CS. Calcium and bone metabolism disorders during pregnancy and lactation. *Endocrinol Metab Clin North Am*. 2011;40(4):795-826.
39. Moller UK, Streyms S, Mosekilde L, et al. Changes in calcitropic hormones, bone markers and insulin-like growth factor I (IGF-I) during pregnancy and postpartum: a controlled cohort study. *Osteoporos Int*. 2013;24(4):1307-1320.
40. Zhong Y, Armbrrecht HJ, Christakos S. Calcitonin, a regulator of the 25-hydroxyvitamin D<sub>3</sub> 1 $\alpha$ -hydroxylase gene. *J Biol Chem*. 2009;284(17):11059-11069.
41. Kovacs CS. Bone development in the fetus and neonate: role of the calcitropic hormones. *Curr Osteoporos Rep*. 2011;9(4):274-283.
42. Jones G, Kano K, Yamada S, Furusawa T, Takayama H, Suda T. Identification of 24,25,26,27-tetranor-23-hydroxyvitamin D<sub>3</sub> as a product of the renal metabolism of 24,25-dihydroxyvitamin D<sub>3</sub>. *Biochemistry*. 1984;23(16):3749-3754.
43. Papapetrou PD. The interrelationship of serum 1,25-dihydroxyvitamin D, 25-hydroxyvitamin D and 24,25-dihydroxyvitamin D in pregnancy at term: a meta-analysis. *Hormones*. 2010;9(2):136-144.
44. Novakovic B, Sibson M, Ng HK, et al. Placenta-specific methylation of the vitamin D 24-hydroxylase gene: implications for feedback autoregulation of active vitamin D levels at the fetomaternal interface. *J Biol Chem*. 2009;284(22):14838-14848.
45. Dror DK. Vitamin D status during pregnancy: maternal, fetal, and postnatal outcomes. *Curr Opin Obstet Gynecol*. 2011;23(6):422-426.
46. Morris GS, Zhou Q, Hegsted M, Keenan MJ. Maternal consumption of a low vitamin D diet retards metabolic and contractile development in the neonatal rat heart. *J Mol Cell Cardiol*. 1995;27(6):1245-1250.
47. Kim BG, Chang SK, Kim SM, Hwang JS, Jung JW. Dilated cardiomyopathy in a 2 month-old infant: a severe form of hypocalcemia with vitamin D deficient rickets. *Korean Circ J*. 2010;40(4):201-203.
48. Jansson T, Powell TL. Role of the placenta in fetal programming: underlying mechanisms and potential interventional approaches. *Clin Sci*. 2007;113(1):1-13.
49. Evans KN, Bulmer JN, Kilby MD, Hewison M. Vitamin D and placental-decidual function. *J Soc Gynecol Invest*. 2004;11(5):263-271.
50. Ma R, Gu Y, Zhao S, Sun J, Groome LJ, Wang Y. Expressions of vitamin D metabolic components VDBP, CYP2R1, CYP27B1, CYP24A1, and VDR in placentas from normal and preeclamptic pregnancies. *Am J Physiol Endocrinol Metab*. 2012;303(7):E928-E935.
51. Mlaka N, Makrakis J, Parkington HC, Tare M, Morley R, Black MJ. Vitamin D deficiency during pregnancy and lactation stimulates nephrogenesis in rat offspring. *Pediatr Nephrol*. 2008;23(1):55-61.
52. Stefanska B, Karlic H, Varga F, Fabianowska-Majewska K, Haslberger A. Epigenetic mechanisms in anti-cancer actions of bioactive food components: the implications in cancer prevention. *Br J Pharmacol*. 2012;167(2):279-297.
53. Barker DJ, Eriksson JG, Forsen T, Osmond C. Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol*. 2002;31(6):1235-1239.
54. Karlic H, Varga F. Impact of vitamin D metabolism on clinical epigenetics. *Clin Epigenetics*. 2011;2(1):55-61.
55. Martin R, Harvey NC, Crozier SR, et al. Placental calcium transporter (PMCA3) gene expression predicts intrauterine bone mineral accrual. *Bone*. 2007;40(5):1203-1208.
56. Lapiolonne A. Vitamin D deficiency during pregnancy may impair maternal and fetal outcomes. *Med Hypotheses*. 2010;74(1):71-75.
57. Tekes K, Gyenge M, Hantos M, Csaba G. Transgenerational hormonal imprinting caused by vitamin A and vitamin D treatment of newborn rats: alterations in the biogenic amine contents of the adult brain. *Brain Dev*. 2009;31(9):666-670.
58. Milovanovic M, Heine G, Hallatschek W, Opitz B, Radbruch A, Worm M. Vitamin D receptor binds to the epsilon gemline gene promoter and exhibits transrepressive activity. *J Allergy Clin Immunol*. 2010;126(5):1016-1023.
59. Kelly JL, Drake MT, Fredericksen ZS, et al. Early life sun exposure, vitamin D-related gene variants, and risk of non-Hodgkin lymphoma. *Cancer Causes Control*. 2012;23(7):1017-1029.
60. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011;96(7):1911-1930.
61. Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr*. 2006;84(1):18-28.
62. Ross AC, Manson JE, Abrams SA, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from

- the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab*. 2011;96(1):53-58.
63. Hewison M, Adams JS. Vitamin D insufficiency and skeletal development in utero. *J Bone Miner Res*. 2010;25(1):11-13.
  64. Sanders KM, Nicholson GC, Ebeling PR. Is high dose vitamin D harmful? *Calcif Tissue Int*. 2013;92(2):191-206.
  65. Mahon P, Harvey N, Crozier S, et al. Low maternal vitamin D status and fetal bone development: cohort study. *J Bone Miner Res*. 2010;25(1):14-19.
  66. Holick MF, Lim R, Dighe AS. Case records of the Massachusetts General Hospital: case 3-2009: a 9-month-old boy with seizures. *N Engl J Med*. 2009;360(4):398-407.
  67. Nowson CA, McGrath JJ, Ebeling PR, et al. Vitamin D and health in adults in Australia and New Zealand: a position statement. *Med J Aust*. 2012;196(11):686-687.
  68. Sai AJ, Walters RW, Fang X, Gallagher JC. Relationship between vitamin D, parathyroid hormone, and bone health. *J Clin Endocrinol Metab*. 2011;96(3):E436-E446.
  69. Bischoff-Ferrari HA, Dietrich T, Orav EJ, et al. Higher 25-hydroxyvitamin D concentrations are associated with better lower-extremity function in both active and inactive persons aged  $\geq 60$  y. *Am J Clin Nutr*. 2004;80(3):752-758.
  70. Muir SW, Montero-Odasso M. Effect of vitamin D supplementation on muscle strength, gait and balance in older adults: a systematic review and meta-analysis. *J Am Geriatr Soc*. 2011;59(12):2291-2300.
  71. Pramyothin P, Holick MF. Vitamin D supplementation: guidelines and evidence for subclinical deficiency. *Curr Opin Gastroenterol*. 2012;28(2):139-150.
  72. Pfeifer M, Begerow B, Minne HW, Suppan K, Fahrleitner-Pammer A, Dobnig H. Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in community-dwelling older individuals. *Osteoporos Int*. 2009;20(2):315-322.
  73. Bischoff-Ferrari HA, Shao A, Dawson-Hughes B, Hathcock J, Giovannucci E, Willett WC. Benefit-risk assessment of vitamin D supplementation. *Osteoporos Int*. 2010;21(7):1121-1132.
  74. Goswami R, Vatsa M, Sreenivas V, et al. Skeletal muscle strength in young Asian Indian females after vitamin D and calcium supplementation: a double-blind randomized controlled clinical trial. *J Clin Endocrinol Metab*. 2012;97(12):4709-4716.
  75. Abrahamsen B, Masud T, Avenell A, et al. Patient level pooled analysis of 68 500 patients from seven major vitamin D fracture trials in US and Europe. *BMJ*. 2010;12(340):b5463.
  76. Elders PJ, Lips P, Netelenbos JC, et al. Long-term effect of calcium supplementation on bone loss in perimenopausal women. *J Bone Miner Res*. 1994;9(7):963-970.
  77. Reid IR, Mason B, Home A, et al. Randomized controlled trial of calcium in healthy older women. *Am J Med*. 2006;119(9):777-785.
  78. Bischoff-Ferrari HA, Willett WC, Orav EJ, et al. A pooled analysis of vitamin D dose requirements for fracture prevention. *N Engl J Med*. 2012;367(1):40-49.
  79. Priemel M, von Dörmann C, Klatte TO, et al. Bone mineralization defects and vitamin D deficiency: histomorphometric analysis of iliac crest bone biopsies and circulating 25-hydroxyvitamin D in 675 patients. *J Bone Miner Res*. 2010;25(2):305-312.
  80. Bergman GJ, Fan T, McFetridge JT, Sen SS. Efficacy of vitamin D<sub>3</sub> supplementation in preventing fractures in elderly women: a meta-analysis. *Curr Med Res Opin*. 2010;26(5):1193-1201.
  81. Bischoff-Ferrari HA, Willett WC, Wong JB, et al. Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials. *Arch Intern Med*. 2009;169(6):551-561.
  82. Avenell A, Gillespie WJ, Gillespie LD, O'Connell D. Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis. *Cochrane Database Syst Rev*. 2009;(2):CD000227.
  83. Moyer VA, U.S. Preventive Services Task Force. Vitamin D and calcium supplementation to prevent fractures in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2013;158(9):691-696.
  84. Bolland MJ, Grey A, Gamble GD, Reid IR. Calcium and vitamin D supplements and health outcomes: a reanalysis of the Women's Health Initiative (WHI) limited access data set. *Am J Clin Nutr*. 2011;94(4):1144-1149.
  85. Winzenberg T, Powell S, Shaw KA, Jones G. Effects of vitamin D supplementation on bone density in healthy children: systematic review and meta-analysis. *BMJ*. 2011;342:c7254.
  86. Huncharek M, Muscat J, Kupelnick B. Impact of dairy products and dietary calcium on bone-mineral content in children: results of a meta-analysis. *Bone*. 2008;43(2):312-321.
  87. Rizzoli R, Boonen S, Brandi ML, et al. Vitamin D supplementation in elderly or postmenopausal women: a 2013 update of the 2008 recommendations from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Curr Med Res Opin*. 2013;29(4):305-313.
  88. Thome-Lyman A, Fawzi WW. Vitamin D during pregnancy and maternal, neonatal and infant health outcomes: a systematic review and meta-analysis. *Paediatr Perinat Epidemiol*. 2012;1:75-90.
  89. De-Regil LM, Palacios C, Ansary A, Kulier R, Pena-Rosas JP. Vitamin D supplementation for women during pregnancy. *Cochrane Database Syst Rev*. 2012;(2):CD008873.
  90. Lai JK, Lucas RM, Clements MS, Roddam AVW, Banks E. Hip fracture risk in relation to vitamin D supplementation and serum 25-hydroxyvitamin D levels: a systematic review and meta-analysis of randomised controlled trials and observational studies. *BMC Public Health*. 2010;10(331):1471-2458.
  91. Izaks GJ. Fracture prevention with vitamin D supplementation: considering the inconsistent results. *BMC Musculoskelet Disord*. 2007;8:26.
  92. Jackson C, Gaugris S, Sen SS, Hosking D. The effect of cholecalciferol (vitamin D<sub>3</sub>) on the risk of fall and fracture: a meta-analysis. *QJM*. 2007;100(4):185-192.
  93. Boonen S, Lips P, Bouillon R, Bischoff-Ferrari HA, Vanderschueren D, Haentjens P. Need for additional calcium to reduce the risk of hip fracture with vitamin D supplementation: evidence from a comparative metaanalysis of randomized controlled trials. *J Clin Endocrinol Metab*. 2007;92(4):1415-1423.
  94. Avenell A, Gillespie WJ, Gillespie LD, O'Connell DL. Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis. *Cochrane Database Syst Rev*. 2005;(3):CD000227.
  95. Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, et al. Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ*. 2009;339:b3692.
  96. Kalyani RR, Stein B, Valiyil R, Manno R, Maynard JW, Crews DC. Vitamin D treatment for the prevention of falls in older adults: systematic review and meta-analysis. *J Am Geriatr Soc*. 2010;58(7):1299-1310.
  97. Chung M, Lee J, Terasawa T, Lau J, Trikalinos TA. Vitamin D with or without calcium supplementation for prevention of cancer and fractures: an updated meta-analysis for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2011;155(12):827-838.
  98. Buttigliero C, Monagheddu C, Petroni P, et al. Prognostic role of vitamin D status and efficacy of vitamin D supplementation in cancer patients: a systematic review. *Oncologist*. 2011;16(9):1215-1227.
  99. Bjelakovic G, Gluud LL, Nikolova D, et al. Vitamin D supplementation for prevention of mortality in adults. *Cochrane Database Syst Rev*. 2011;(7):CD007470.
  100. Irlam JH, Visser MM, Rollins NN, Siegfried N. Micronutrient supplementation in children and adults with HIV infection. *Cochrane Database Syst Rev*. 2010;(12):CD003650.
  101. Autier P, Gandini S, Mullie P. A systematic review: influence of vitamin D supplementation on serum 25-hydroxyvitamin D concentration. *J Clin Endocrinol Metab*. 2012;97(8):2606-2613.


102. Bjorkman M, Sorva A, Tilvis R. Responses of parathyroid hormone to vitamin D supplementation: a systematic review of clinical trials. *Arch Gerontol Geriatr.* 2009;48(2):160-166.
103. Tripkovic L, Lambert H, Hart K, et al. Comparison of vitamin D<sub>2</sub> and vitamin D<sub>3</sub> supplementation in raising serum 25-hydroxyvitamin D status: a systematic review and meta-analysis. *Am J Clin Nutr.* 2012;95(6):1357-1364.
104. Kandula P, Dobre M, Schold JD, Schreiber MJ Jr, Mehrotra R, Navaneethan SD. Vitamin D supplementation in chronic kidney disease: a systematic review and meta-analysis of observational studies and randomized controlled trials. *Clin J Am Soc Nephrol.* 2011;6(1):50-62.
105. Song Y, Wang L, Pittas AG, et al. Blood 25-hydroxy vitamin D levels and incident type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care.* 2013;36(5):1422-1428.
106. George PS, Pearson ER, Witham MD. Effect of vitamin D supplementation on glycaemic control and insulin resistance: a systematic review and meta-analysis. *Diabet Med.* 2012;29(8):e142-e150.
107. Bath-Hextall FJ, Jenkinson C, Humphreys R, Williams HC. Dietary supplements for established atopic eczema. *Cochrane Database Syst Rev.* 2012;(2):CD005205.
108. Annweiler C, Schott AM, Bernut G, Fantino B, Beauchet O. Vitamin D-related changes in physical performance: a systematic review. *J Nutr Health Aging.* 2009;13(10):893-898.
109. Wang L, Manson JE, Song Y, Sesso HD. Systematic review: vitamin D and calcium supplementation in prevention of cardiovascular events. *Ann Intern Med.* 2010;152(5):315-323.
110. Pittas AG, Chung M, Trikalinos T, et al. Systematic review: vitamin D and cardiometabolic outcomes. *Ann Intern Med.* 2010;152(5):307-314.
111. Fergusson JH, Chang AB. Vitamin D supplementation for cystic fibrosis. *Cochrane Database Syst Rev.* 2009;(4):CD007398.
112. Abba K, Sudarsanam TD, Grobler L, Volmink J. Nutritional supplements for people being treated for active tuberculosis. *Cochrane Database Syst Rev.* 2008;(4):CD006086.
113. Autier P, Gandini S. Vitamin D supplementation and total mortality: a meta-analysis of randomized controlled trials. *Arch Intern Med.* 2007;167(16):1730-1737.
114. Hamis DM, Go VL. Vitamin D and colon carcinogenesis. *J Nutr.* 2004;134(12, suppl):3463S-3471S.
115. Shui IM, Mucci LA, Kraft P, et al. Vitamin D-related genetic variation, plasma vitamin D, and risk of lethal prostate cancer: a prospective nested case-control study. *J Natl Cancer Inst.* 2012;104(9):690-699.
116. Giangreco AA, Nonn L. The sum of many small changes: microRNAs are specifically and potentially globally altered by vitamin D(3) metabolites. *J Steroid Biochem Mol Biol.* 2013 Jan 16 [Epub ahead of print]. <http://dx.doi.org/10.1016/j.jsbmb.2013.01.001>.
117. Holick MF. Vitamin D and sunlight: strategies for cancer prevention and other health benefits. *Clin J Am Soc Nephrol.* 2008;3(5):1548-1554.
118. Freedman DM, Looker AC, Abnet CC, Linet MS, Graubard BI. Serum 25-hydroxyvitamin D and cancer mortality in the NHANES III study (1988-2006). *Cancer Res.* 2010;70(21):8587-8597.
119. Garland CF, Gorham ED, Mohr SB, et al. Vitamin D and prevention of breast cancer: pooled analysis. *J Steroid Biochem Mol Biol.* 2007;103(3-5):708-711.
120. Grant WB. An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar ultraviolet-B radiation. *Cancer.* 2002;94(6):1867-1875.
121. Lee JE, Li H, Chan AT, et al. Circulating levels of vitamin D and colon and rectal cancer: the Physicians' Health Study and a meta-analysis of prospective studies. *Cancer Prev Res.* 2011;4(5):735-743.
122. Holick MF. Calcium plus vitamin D and the risk of colorectal cancer. *N Engl J Med.* 2006;354(21):2287-2288.
123. Bischoff-Ferrari HA, Willett WC, Wong JB, Giovannucci E, Dietrich T, Dawson-Hughes B. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA.* 2005;293(18):2257-2264.
124. Holick MF. Evidence-based D-bate on health benefits of vitamin D revisited. *Dermatoendocrinol.* 2012;4(2):183-190.
125. Grant WB. Geographic variation of prostate cancer mortality rates in the United States: implications for prostate cancer risk related to vitamin D. *Int J Cancer.* 2004;111(3):470-471.
126. Durup D, Jorgensen HL, Christensen J, Schwarz P, Heegaard AM, Lind B. A reverse J-shaped association of all-cause mortality with serum 25-hydroxyvitamin D in general practice: the CopD study. *J Clin Endocrinol Metab.* 2012;97(8):2644-2652.
127. Vieth R. Enzyme kinetics hypothesis to explain the U-shaped risk curve for prostate cancer vs. 25-hydroxyvitamin D in Nordic countries. *Int J Cancer.* 2004;111(3):468.
128. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Guidelines for preventing and treating vitamin D deficiency and insufficiency revisited. *J Clin Endocrinol Metab.* 2012;97(4):1153-1158.
129. Wactawski-Wende J, Kotchen JM, Anderson GL, et al. Calcium plus vitamin D supplementation and the risk of colorectal cancer. *N Engl J Med.* 2006;354(7):684-696.
130. Lappe JM, Travers-Gustafson D, Davies KM, Recker RR, Heaney RP. Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial. *Am J Clin Nutr.* 2007;85(6):1586-1591.
131. Skinner HG, Michaud DS, Giovannucci E, Willett WC, Colditz GA, Fuchs CS. Vitamin D intake and the risk for pancreatic cancer in two cohort studies. *Cancer Epidemiol Biomarkers Prev.* 2006;15(9):1688-1695.
132. Stolzenberg-Solomon RZ, Hayes RB, Horst RL, Anderson KE, Hollis BW, Silverman DT. Serum vitamin D and risk of pancreatic cancer in the Prostate, Lung, Colorectal, and Ovarian Screening Trial. *Cancer Res.* 2009;69(4):1439-1447.
133. Wolpin BI, Ng K, Bao Y, et al. Plasma 25-hydroxyvitamin D and risk of pancreatic cancer. *Cancer Epidemiol Biomarkers Prev.* 2012;21(1):82-91.
134. Holick MF. Vitamin D, sunlight and cancer connection. *Anti-cancer Agents Med Chem.* 2013;13(1):70-82.
135. Grant WB. An estimate of the global reduction in mortality rates through doubling vitamin D levels. *Eur J Clin Nutr.* 2011;65(9):1016-1026.
136. Grant WB. Lower vitamin-D production from solar ultraviolet-B irradiance may explain some differences in cancer survival rates. *J Natl Med Assoc.* 2006;98(3):357-364.
137. Helzlsouer KJ. Overview of the Cohort Consortium Vitamin D Pooling Project of Rarer Cancers. *Am J Epidemiol.* 2010;172(1):4-9.
138. Grant WB. Ecological studies of the UV-B-vitamin D-cancer hypothesis. *Anticancer Res.* 2012;32(1):223-236.
139. Pilz S, Tomaschitz A, Marz W, et al. Vitamin D, cardiovascular disease and mortality. *Clin Endocrinol.* 2011;75(5):575-584.
140. Luxwolda MF, Kuipers RS, Kema IP, Dijk-Brouwer DA, Muskiet FA. Traditionally living populations in East Africa have a mean serum 25-hydroxyvitamin D concentration of 115 nmol/l. *Br J Nutr.* 2012;108(9):1557-1561.
141. Melamed ML, Michos ED, Post W, Astor B. 25-Hydroxyvitamin D levels and the risk of mortality in the general population. *Arch Intern Med.* 2008;168(15):1629-1637.
142. Zittermann A, Iodice S, Pilz S, Grant WB, Bagnardi V, Gandini S. Vitamin D deficiency and mortality risk in the general population: a meta-analysis of prospective cohort studies. *Am J Clin Nutr.* 2012;95(1):91-100.
143. Ma Y, Zhang P, Wang F, Yang J, Liu Z, Qin H. Association between vitamin D and risk of colorectal cancer: a systematic review of prospective studies. *J Clin Oncol.* 2011;29(28):3775-3782.


144. Pilz S, Iodice S, Zittermann A, Grant WB, Gandini S. Vitamin D status and mortality risk in CKD: a meta-analysis of prospective studies. *Am J Kidney Dis*. 2011;58(3):374-382.
145. Rejnmark L, Avenell A, Masud T, et al. Vitamin D with calcium reduces mortality: patient level pooled analysis of 70,528 patients from eight major vitamin D trials. *J Clin Endocrinol Metab*. 2012;97(8):2670-2681.
146. Thomas GN, ó Hartaigh B, Bosch JA, et al. Vitamin D levels predict all-cause and cardiovascular disease mortality in subjects with the metabolic syndrome: the Ludwigshafen Risk and Cardiovascular Health (LURIC) Study. *Diabetes Care*. 2012;35(5):1158-1164.
147. Doherty TM, Tang W, Dascalos S, et al. Ethnic origin and serum levels of 1,25-dihydroxyvitamin D<sub>3</sub> are independent predictors of coronary calcium mass measured by electron-beam computed tomography. *Circulation*. 1997;96(5):1477-1481.
148. Watson KE, Abrolat ML, Malone LL, et al. Active serum vitamin D levels are inversely correlated with coronary calcification. *Circulation*. 1997;96(6):1755-1760.
149. Gigolini M, Iagulli MP, Miconi V, Galiotto M, Lombardi S, Targher G. Serum 25-hydroxyvitamin D<sub>3</sub> concentrations and prevalence of cardiovascular disease among type 2 diabetic patients. *Diabetes Care*. 2006;29(3):722-724.
150. Arson Y, Itzhaky D, Mosseri M, et al. Vitamin D inflammatory cytokines and coronary events: a comprehensive review [published online January 12, 2013]. *Clin Rev Allergy Immunol*. <http://dx.doi.org/10.1007/s12016-013-8356-0>.
151. Grundmann M, Haidar M, Placzko S, et al. Vitamin D improves the angiogenic properties of endothelial progenitor cells. *Am J Physiol Cell Physiol*. 2012;303(9):29.
152. Wang TJ, Pencina MJ, Booth SL, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation*. 2008;117(4):503-511.
153. Anderson JL, May HT, Home BD, et al. Relation of vitamin D deficiency to cardiovascular risk factors, disease status, and incident events in a general healthcare population. *Am J Cardiol*. 2010;106(7):963-968.
154. Tamez H, Kalim S, Thadhani RI. Does vitamin D modulate blood pressure? *Curr Opin Nephrol Hypertens*. 2013;22(2):204-209.
155. Dong Y, Stallmann-Jorgensen IS, Pollock NK, et al. A 16-week randomized clinical trial of 2000 international units daily vitamin D<sub>3</sub> supplementation in black youth: 25-hydroxyvitamin D, adiposity, and arterial stiffness. *J Clin Endocrinol Metab*. 2010;95(10):4584-4591.
156. Kumar J, Muntner P, Kaskel FJ, Hailpern SM, Melamed ML. Prevalence and associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001-2004. *Pediatrics*. 2009;124(3):e362-e370.
157. Forouhi NG, Ye Z, Rickard AP, et al. Circulating 25-hydroxyvitamin D concentration and the risk of type 2 diabetes: results from the European Prospective Investigation into Cancer (EPIC)-Norfolk cohort and updated meta-analysis of prospective studies. *Diabetologia*. 2012;55(8):2173-2182.
158. Welsh P, Doolin O, McConnachie A, et al. Circulating 25OHD, dietary vitamin D, PTH, and calcium associations with incident cardiovascular disease and mortality: the MIDSPAN Family Study. *J Clin Endocrinol Metab*. 2012;97(12):4578-4587.
159. Agmon-Levin N, Theodor E, Segal RM, Shoenfeld Y. Vitamin D in systemic and organ-specific autoimmune diseases [published online December 14, 2012]. *Clin Rev Allergy Immunol*. <http://dx.doi.org/10.1007/s12016-012-8342-y>.
160. Higgins MJ, Mackie SL, Thalayasingam N, Bingham SJ, Hamilton J, Kelly CA. The effect of vitamin D levels on the assessment of disease activity in rheumatoid arthritis [published online January 23, 2013]. *Clin Rheumatol*. <http://dx.doi.org/10.1007/s10067-013-2174-x>.
161. Szekeley JI, Pataki A. Effects of vitamin D on immune disorders with special regard to asthma, COPD and autoimmune diseases: a short review. *Expert Rev Respir Med*. 2012;6(6):683-704.
162. Cantoma MT, Mahon BD. Mounting evidence for vitamin D as an environmental factor affecting autoimmune disease prevalence. *Exp Biol Med*. 2004;229(11):1136-1142.
163. Villaggio B, Soldano S, Cutolo M. 1,25-dihydroxyvitamin D<sub>3</sub> downregulates aromatase expression and inflammatory cytokines in human macrophages. *Clin Exp Rheumatol*. 2012;30(6):934-938.
164. Merino LA, Curtis J, Mikuls TR, Cerhan JR, Criswell LA, Saag KG. Vitamin D intake is inversely associated with rheumatoid arthritis: results from the Iowa Women's Health Study. *Arthritis Rheum*. 2004;50(1):72-77.
165. Andjelkovic Z, Vojinovic J, Pejnovic N, et al. Disease modifying and immunomodulatory effects of high dose 1,25(OH)<sub>2</sub>D<sub>3</sub> in rheumatoid arthritis patients. *Clin Exp Rheumatol*. 1999;17(4):453-456.
166. Sen D, Ranganathan P. Vitamin D in rheumatoid arthritis: panacea or placebo? *Discov Med*. 2012;14(78):311-319.
167. Haga HJ, Schmedes A, Naderi Y, Moreno AM, Peen E. Severe deficiency of 25-hydroxyvitamin D<sub>3</sub> (25-OH-D<sub>3</sub>) is associated with high disease activity of rheumatoid arthritis. *Clin Rheumatol*. 2013;15:15.
168. Schwalfenberg GK. Solar radiation and vitamin D: mitigating environmental factors in autoimmune disease. *J Environ Public Health*. 2012;2012:619381.
169. Knip M, Akerblom HK. Early nutrition and later diabetes risk. *Adv Exp Med Biol*. 2005;569:142-150.
170. Stene LC, Ultnsen J, Magnus P, Joner G. Use of cod liver oil during pregnancy associated with lower risk of type 1 diabetes in the offspring. *Diabetologia*. 2000;43(9):1093-1098.
171. Hypponen E, Laara E, Reunanen A, Jarvelin MR, Virtanen SM. Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet*. 2001;358(9292):1500-1503.
172. Rajakumar K, de las Heras J, Lee S, Holick MF, Arslanian SA. 25-Hydroxyvitamin D concentrations and in vivo insulin sensitivity and  $\beta$ -cell function relative to insulin sensitivity in black and white youth. *Diabetes Care*. 2012;35(3):627-633.
173. Simon KC, Munger KL, Ascherio A. Vitamin D and multiple sclerosis: epidemiology, immunology, and genetics. *Curr Opin Neurol*. 2012;25(3):246-251.
174. Munger KL, Zhang SM, O'Reilly E, et al. Vitamin D intake and incidence of multiple sclerosis. *Neurology*. 2004;62(1):60-65.
175. Runia TF, Hop WC, de Rijke YB, Buijvec D, Hintzen RQ. Lower serum vitamin D levels are associated with a higher relapse risk in multiple sclerosis. *Neurology*. 2012;79(3):261-266.
176. Weinstock-Guttman B, Mehta BK, Ramanathan M, et al. Vitamin D and multiple sclerosis. *Neurologist*. 2012;18(4):179-183.
177. Kimball SM, Ursell MR, O'Connor P, Vieth R. Safety of vitamin D<sub>3</sub> in adults with multiple sclerosis. *Am J Clin Nutr*. 2007;86(3):645-651.
178. Stein MS, Liu Y, Gray OM, et al. A randomized trial of high-dose vitamin D<sub>2</sub> in relapsing-remitting multiple sclerosis. *Neurology*. 2011;77(17):1611-1618.
179. Burton JM, Kimball S, Vieth R, et al. A phase III dose-escalation trial of vitamin D<sub>3</sub> and calcium in multiple sclerosis. *Neurology*. 2010;74(23):1852-1859.
180. Ramagopalan SV, Heger A, Berlanga AJ, et al. A ChIP-seq defined genome-wide map of vitamin D receptor binding: associations with disease and evolution. *Genome Res*. 2010;20(10):1352-1360.
181. Jorgensen SP, Agnholt J, Glerup H, et al. Clinical trial: vitamin D<sub>3</sub> treatment in Crohn's disease: a randomized double-blind placebo-controlled study. *Aliment Pharmacol Ther*. 2010;32(3):377-383.
182. Bartels LE, Jorgensen SP, Bendix M, et al. 25-Hydroxyvitamin D<sub>3</sub> modulates dendritic cell phenotype and function in Crohn's disease. *Inflammopharmacology*. 2013;21(2):177-186.
183. Holick MF. Resurrection of vitamin D deficiency and rickets. *J Clin Invest*. 2006;116(8):2062-2072.

184. Camargo CA Jr, Ingham T, Wickens K, et al. Cord-blood 25-hydroxyvitamin D levels and risk of respiratory infection, wheezing, and asthma. *Pediatrics*. 2011;127(1):e180-e187.
185. Sabetta JR, DePetrillo P, Cipriani RJ, Smardin J, Burns LA, Landry ML. Serum 25-hydroxyvitamin D and the incidence of acute viral respiratory tract infections in healthy adults. *PLoS One*. 2010;5(6):0011088.
186. Urashima M, Segawa T, Okazaki M, Kurihara M, Wada Y, Ida H. Randomized trial of vitamin D supplementation to prevent seasonal influenza A in schoolchildren. *Am J Clin Nutr*. 2010;91(5):1255-1260.
187. Sundar IK, Rahman I. Vitamin D and susceptibility of chronic lung diseases: role of epigenetics. *Front Pharmacol*. 2011;2:50.
188. Wjst M. Is vitamin D supplementation responsible for the allergy pandemic? *Curr Opin Allergy Clin Immunol*. 2012;12(3):257-262.
189. Tolppanen AM, Williams D, Henderson J, Lawlor DA. Serum 25-hydroxyvitamin D and ionised calcium in relation to lung function and allergen skin tests. *Eur J Clin Nutr*. 2011;65(4):493-500.
190. Jones AP, Palmer D, Zhang G, Prescott SL. Cord blood 25-hydroxyvitamin D<sub>3</sub> and allergic disease during infancy. *Pediatrics*. 2012;130(5):e1128-e1135.
191. Rochat MK, Ege MJ, Plabst D, et al. Maternal vitamin D intake during pregnancy increases gene expression of ILT3 and ILT4 in cord blood. *Clin Exp Allergy*. 2010;40(5):786-794.
192. Eyles DW, Borne TH, McGrath JJ. Vitamin D, effects on brain development, adult brain function and the links between low levels of vitamin D and neuropsychiatric disease. *Front Neuroendocrinol*. 2013;34(1):47-64.
193. Bartoccini E, Marini F, Damaskopoulou E, et al. Nuclear lipid microdomains regulate nuclear vitamin D<sub>3</sub> uptake and influence embryonic hippocampal cell differentiation. *Mol Biol Cell*. 2011;22(17):3022-3031.
194. Gale CR, Robinson SM, Harvey NC, et al. Maternal vitamin D status during pregnancy and child outcomes. *Eur J Clin Nutr*. 2008;62(1):68-77.
195. O'Loan J, Eyles DW, Kesby J, Ko P, McGrath JJ, Borne TH. Vitamin D deficiency during various stages of pregnancy in the rat: its impact on development and behaviour in adult offspring. *Psychoneuroendocrinology*. 2007;32(3):227-234.
196. Becker A, Eyles DW, McGrath JJ, Grecksch G. Transient prenatal vitamin D deficiency is associated with subtle alterations in learning and memory functions in adult rats. *Behav Brain Res*. 2005;161(2):306-312.
197. McGrath JJ, Saari K, Hakko H, et al. Vitamin D supplementation during the first year of life and risk of schizophrenia: a Finnish birth cohort study. *Schizophr Res*. 2004;67(2-3):237-245.
198. Newschaffer CJ, Croen LA, Daniels J, et al. The epidemiology of autism spectrum disorders. *Annu Rev Public Health*. 2007;28:235-258.
199. Yu J, Gattioni-Celli M, Zhu H, et al. Vitamin D<sub>3</sub>-enriched diet correlates with a decrease of amyloid plaques in the brain of AβPP transgenic mice. *J Alzheimers Dis*. 2011;25(2):295-307.
200. Whitehouse AJ, Holt BJ, Serralha M, Holt PG, Kusel MM, Hart PH. Maternal serum vitamin D levels during pregnancy and offspring neurocognitive development. *Pediatrics*. 2012;129(3):485-493.
201. Groves NJ, Kesby JP, Eyles DW, McGrath JJ, Mackay-Sim A, Borne TH. Adult vitamin D deficiency leads to behavioural and brain neurochemical alterations in C57BL/6j and BALB/c mice. *Behav Brain Res*. 2013;241:120-131.
202. Maddock J, Berry DJ, Geoffrey MC, Power C, Hypponen E. Vitamin D and common mental disorders in mid-life: cross-sectional and prospective findings. *Clin Nutr*. 2013;32(13):00030-00037.
203. Anglin RE, Samaan Z, Walter SD, McDonald SD. Vitamin D deficiency and depression in adults: systematic review and meta-analysis. *Br J Psychiatry*. 2013;202:100-107.
204. Hoogendijk WJ, Lips P, Dik MG, Deeg DJ, Beekman AT, Penninx BW. Depression is associated with decreased 25-hydroxyvitamin D and increased parathyroid hormone levels in older adults. *Arch Gen Psychiatry*. 2008;65(5):508-512.
205. Armstrong DJ, Meenagh GK, Bickle I, Lee AS, Curran ES, Finch MB. Vitamin D deficiency is associated with anxiety and depression in fibromyalgia. *Clin Rheumatol*. 2007;26(4):551-554.
206. Jorde R, Waterloo K, Saleh F, Haug E, Svartberg J. Neuropsychological function in relation to serum parathyroid hormone and serum 25-hydroxyvitamin D levels: the Tromso study. *J Neurol*. 2006;253(4):464-470.
207. Lansdowne AT, Provost SC. Vitamin D<sub>3</sub> enhances mood in healthy subjects during winter. *Psychopharmacology*. 1998;135(4):319-323.
208. Gloth FM III, Alam W, Hollis B. Vitamin D vs broad spectrum phototherapy in the treatment of seasonal affective disorder. *J Nutr Health Aging*. 1999;3(1):5-7.
209. Wei SQ, Qi HP, Luo ZC, Fraser WD. Maternal vitamin D status and adverse pregnancy outcomes: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med*. 2013;26(9):889-899.
210. Merewood A, Mehta SD, Chen TC, Bauchner H, Holick MF. Association between vitamin D deficiency and primary cesarean section. *J Clin Endocrinol Metab*. 2009;94(3):940-945.
211. Poel YH, Hummel P, Lips P, Stam F, van der Ploeg T, Simsek S. Vitamin D and gestational diabetes: a systematic review and meta-analysis. *Eur J Intern Med*. 2012;23(5):465-469.
212. Hollis BW, Horst RL. The assessment of circulating 25(OH)D and 1,25(OH)<sub>2</sub>D: where we are and where we are going. *J Steroid Biochem Mol Biol*. 2007;103(3-5):473-476.
213. Schottker B, Jansen EH, Haug U, Schomburg L, Kohrle J, Brenner H. Standardization of misleading immunoassay based 25-hydroxyvitamin D levels with liquid chromatography tandem-mass spectrometry in a large cohort study. *PLoS One*. 2012;7(11):e48774.
214. Binkley N, Krueger D, Cowgill CS, et al. Assay variation confounds the diagnosis of hypovitaminosis D: a call for standardization. *J Clin Endocrinol Metab*. 2004;89(7):3152-3157.
215. Carter GD, Carter R, Jones J, Berry J. How accurate are assays for 25-hydroxyvitamin D? data from the international vitamin D external quality assessment scheme. *Clin Chem*. 2004;50(11):2195-2197.
216. Singh RJ, Taylor RL, Reddy GS, Grebe SK. C-3 epimers can account for a significant proportion of total circulating 25-hydroxyvitamin D in infants, complicating accurate measurement and interpretation of vitamin D status. *J Clin Endocrinol Metab*. 2006;91(8):3055-3061.
217. Looker AC, Johnson CL, Lacher DA, Pfeiffer CM, Schleicher RL, Sempos CT. Vitamin D status: United States, 2001-2006. *NCHS Data Brief*. 2011;59:1-8.
218. Centers for Disease Control and Prevention (CDC). Centers for Disease Control and Prevention (CDC). Centers for Disease Control and Prevention (CDC). Centers for Disease Control and Prevention (CDC). Analytical note for NHANES 2000-2006 and NHANES III (1988-1994) 25-hydroxyvitamin D analysis. [http://www.cdc.gov/nchs/data/nhanes/nhanes3/VitaminD\\_analyticnote.pdf](http://www.cdc.gov/nchs/data/nhanes/nhanes3/VitaminD_analyticnote.pdf). Revised November 2010. Accessed June 6, 2013.
219. Carter GD. 25-Hydroxyvitamin D assays: the quest for accuracy. *Clin Chem*. 2009;55(7):1300-1302.
220. Sempos CT, Vesper HW, Phinney KW, Thienpont LM, Coates PM. Vitamin D status as an international issue: national surveys and the problem of standardization. *Scand J Clin Lab Invest Suppl*. 2012;243:32-40.
221. Yetley EA, Pfeiffer CM, Schleicher RL, et al. NHANES monitoring of serum 25-hydroxyvitamin D: a roundtable summary. *J Nutr*. 2010;140(11):2030S-2045S.
222. Fraser WD, Milan AM. Vitamin D assays: past and present debates, difficulties, and developments. *Calif Tissue Int*. 2013;92(2):118-127.

223. Glendenning P, Taranto M, Noble JM, et al. Current assays overestimate 25-hydroxyvitamin D<sub>3</sub> and underestimate 25-hydroxyvitamin D<sub>2</sub> compared with HPLC: need for assay-specific decision limits and metabolite-specific assays. *Ann Clin Biochem*. 2006;43(pt 1):23-30.
224. Thandrayen K, Pettifor JM. Maternal vitamin D status: implications for the development of infantile nutritional rickets. *Rheum Dis Clin North Am*. 2012;38(1):61-79.
225. Shivane VK, Sarathi V, Bandgar T, Menon P, Shah NS. High prevalence of hypovitaminosis D in young healthy adults from the western part of India. *Postgrad Med J*. 2011; 87(1030):514-518.
226. van Schoor NM, Lips P. Worldwide vitamin D status. *Best Pract Res Clin Endocrinol Metab*. 2011;25(4):671-680.
227. Fields J, Trivedi NJ, Horton E, Mechanick JL. Vitamin D in the Persian Gulf: integrative physiology and socioeconomic factors. *Curr Osteoporos Rep*. 2011;9(4):243-250.
228. Ganji V, Zhang X, Tangpricha V. Serum 25-hydroxyvitamin D concentrations and prevalence estimates of hypovitaminosis D in the U.S. population based on assay-adjusted data. *J Nutr*. 2012;142(3):498-507.
229. Greene-Finestone LS, Berger C, de Groh M, et al. 25-Hydroxyvitamin D in Canadian adults: biological, environmental, and behavioral correlates. *Osteoporos Int*. 2011;22(5): 1389-1399.
230. Gonzalez-Gross M, Valtuena J, Breidenassel C, et al. Vitamin D status among adolescents in Europe: the Healthy Lifestyle in Europe by Nutrition in Adolescence study. *Br J Nutr*. 2012; 107(5):755-764.
231. Daly RM, Gagnon C, Lu ZX, et al. Prevalence of vitamin D deficiency and its determinants in Australian adults aged 25 years and older: a national, population-based study. *Clin Endocrinol*. 2012;77(1):26-35.
232. Clemens TL, Adams JS, Henderson SL, Holick MF. Increased skin pigment reduces the capacity of skin to synthesise vitamin D<sub>3</sub>. *Lancet*. 1982;1(8263):74-76.
233. Manicourt DH, Devogelaer JP. Urban tropospheric ozone increases the prevalence of vitamin D deficiency among Belgian postmenopausal women with outdoor activities during summer. *J Clin Endocrinol Metab*. 2008;93(10): 3893-3899.
234. Holick MF. Vitamin D: evolutionary, physiological and health perspectives. *Curr Drug Targets*. 2011;12(1):4-18.
235. van den Berg G, van Eijsden M, Vrijkotte TG, Gemke RJ. Sub-optimal maternal vitamin D status and low education level as determinants of small-for-gestational-age birth weight. *Eur J Nutr*. 2013;52(1):273-279.
236. Godar DE, Pope SJ, Grant WB, Holick MF. Solar UV doses of young Americans and vitamin D<sub>3</sub> production. *Environ Health Perspect*. 2012;120(1):139-143.
237. MacLaughlin J, Holick MF. Aging decreases the capacity of human skin to produce vitamin D<sub>3</sub>. *J Clin Invest*. 1985;76(4): 1536-1538.
238. Levin GP, Robinson-Cohen C, de Boer JH, et al. Genetic variants and associations of 25-hydroxyvitamin D concentrations with major clinical outcomes. *JAMA*. 2012;308(18):1898-1905.
239. Signorello LB, Shi J, Cai Q, et al. Common variation in vitamin D pathway genes predicts circulating 25-hydroxyvitamin D levels among African Americans. *PLoS One*. 2011;6(12): e28623.
240. Ahn J, Yu K, Stolzenberg-Solomon R, et al. Genome-wide association study of circulating vitamin D levels. *Hum Mol Genet*. 2010;19(13):2739-2745.
241. Wang TJ, Zhang F, Richards JB, et al. Common genetic determinants of vitamin D insufficiency: a genome-wide association study. *Lancet*. 2010;376(9736):180-188.
242. Engelman CD, Fingerlin TE, Langefeld CD, et al. Genetic and environmental determinants of 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D levels in Hispanic and African Americans. *J Clin Endocrinol Metab*. 2008;93(9):3381-3388.
243. Zhu H, Wang X, Shi H, et al. A genome-wide methylation study of severe vitamin D deficiency in African American adolescents. *J Pediatr*. 2013;162(5):1004-1009.
244. Hiraki LT, Major JM, Chen C, et al. Exploring the genetic architecture of circulating 25-hydroxyvitamin D. *Genet Epidemiol*. 2013;37(1):92-98.
245. Schlingmann KP, Kaufmann M, Weber S, et al. Mutations in CYP24A1 and idiopathic infantile hypercalcemia. *N Engl J Med*. 2011;365(5):410-421.
246. Kennedy C, Bajdik CD, Willemze R, De Gruij FR, Bouwes Bavinck JN. The influence of painful sunburns and lifetime sun exposure on the risk of actinic keratoses, seborrheic warts, melanocytic nevi, atypical nevi, and skin cancer. *J Invest Dermatol*. 2003;120(6):1087-1093.
247. Garland FC, White MR, Garland CF, Shaw E, Gorham ED. Occupational sunlight exposure and melanoma in the U.S. Navy. *Arch Environ Health*. 1990;45(5):261-267.
248. Tangpricha V, Turner A, Spina C, Decastro S, Chen TC, Holick MF. Tanning is associated with optimal vitamin D status (serum 25-hydroxyvitamin D concentration) and higher bone mineral density. *Am J Clin Nutr*. 2004;80(6):1645-1649.
249. Grant WB. In defense of the sun: an estimate of changes in mortality rates in the United States if mean serum 25-hydroxyvitamin D levels were raised to 45 ng/mL by solar ultraviolet-B irradiance. *Dermatoendocrinol*. 2009;1(4): 207-214.
250. Dabai NS, Pramyothin P, Holick MF. The effect of ultraviolet radiation from a novel portable fluorescent lamp on serum 25-hydroxyvitamin D<sub>3</sub> levels in healthy adults with Fitzpatrick skin types II and III. *Photodermatol Photoimmunol Photomed*. 2012;28(6):307-311.
251. Moore C, Murphy MM, Keast DR, Holick MF. Vitamin D intake in the United States. *J Am Diet Assoc*. 2004;104(6): 980-983.
252. Black LJ, Seamans KM, Cashman KD, Kiely M. An updated systematic review and meta-analysis of the efficacy of vitamin D food fortification. *J Nutr*. 2012;142(6): 1102-1108.
253. Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr*. 2003;77(1):204-210.
254. Demetriou ET, Travison TG, Holick MF. Treatment with 50,000 IU vitamin D<sub>2</sub> every other week and effect on serum 25-hydroxyvitamin D<sub>2</sub>, 25-hydroxyvitamin D<sub>3</sub>, and total 25-hydroxyvitamin D in a clinical setting. *Endocr Pract*. 2012; 18(3):399-402.
255. Pietras SM, Obayan BK, Cai MH, Holick MF. Vitamin D<sub>2</sub> treatment for vitamin D deficiency and insufficiency for up to 6 years. *Arch Intern Med*. 2009;169(19):1806-1808.
256. Sanders KM, Stuart AL, Williamson EJ, et al. Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *JAMA*. 2010; 303(18):1815-1822.
257. Heikinheimo RJ, Inkovaara JA, Harju EJ, et al. Annual injection of vitamin D and fractures of aged bones. *Calcif Tissue Int*. 1992;51(2):105-110.
258. Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D<sub>3</sub> (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomised double blind controlled trial. *BMJ*. 2003; 326(7387):469.
259. Lee JM, Smith JR, Philipp BL, Chen TC, Mathieu J, Holick MF. Vitamin D deficiency in a healthy group of mothers and newborn infants. *Clin Pediatr*. 2007;46(1):42-44.
260. Hollis BW, Wagner CL. Vitamin D and pregnancy: skeletal effects, nonskeletal effects, and birth outcomes. *Calcif Tissue Int*. 2013;92(2):128-139.
261. Hollis BW, Wagner CL. Vitamin D requirements during lactation: high-dose maternal supplementation as therapy to

- prevent hypovitaminosis D for both the mother and the nursing infant. *Am J Clin Nutr*. 2004;80(6, suppl):1752S-1758S.
262. Chandra P, Wolfenden LL, Ziegler TR, et al. Treatment of vitamin D deficiency with UV light in patients with malabsorption syndromes: a case series. *Photodermatol Photoimmunol Photomed*. 2007;23(5):179-185.
263. Biancuzzo RM, Clarke N, Reitz RE, Trivison TG, Holick MF. Serum concentrations of 1,25-dihydroxyvitamin D<sub>2</sub> and 1,25-dihydroxyvitamin D<sub>3</sub> in response to vitamin D<sub>2</sub> and vitamin D<sub>3</sub> supplementation. *J Clin Endocrinol Metab*. 2013; 98(3):973-979.
264. Holick MF, Biancuzzo RM, Chen TC, et al. Vitamin D<sub>2</sub> is as effective as vitamin D<sub>3</sub> in maintaining circulating concentrations of 25-hydroxyvitamin D. *J Clin Endocrinol Metab*. 2008; 93(3):677-681.
265. Gordon CM, Williams AL, Feldman HA, et al. Treatment of hypovitaminosis D in infants and toddlers. *J Clin Endocrinol Metab*. 2008;93(7):2716-2721.
266. Thacher TD, Fischer PR, Obadofin MO, Levine MA, Singh RJ, Pettifor JM. Comparison of metabolism of vitamins D<sub>2</sub> and D<sub>3</sub> in children with nutritional rickets. *J Bone Miner Res*. 2010; 25(9):1988-1995.
267. Keegan R, Lu Z, Bogusz J, Holick M. Photobiology of vitamin D in mushrooms and its bioavailability in humans. *Dermato-Endocrinology*. 2013;5:1.
268. Hathcock JN, Shao A, Vieth R, Heaney R. Risk assessment for vitamin D. *Am J Clin Nutr*. 2007;85(1):6-18.
269. Rajakumar K, Reis EC, Holick MF. Dosing error with over-the-counter vitamin D supplement: a risk for vitamin D toxicity in infants. *Clin Pediatr*. 2013;52(1):82-85.





**HB 356**  
**Prevention of Disease**  
**Decreasing the Alaska Healthcare Liability**

## **\$3.8 BILLION**

**\$3.8 Billion is the amount of our PERS/TRS unfunded liability attributable to healthcare according to the Department of Administration.**

**The old estimate for a 2% annual, out-year, healthcare cost increase was used for setting the contribution rates to fully cover anticipated liabilities.**

**So this \$3.8 billion represents the healthcare cost inflation above 2%.**



# Why are we here?

*The State of Alaska is a significant health care consumer.*

<b>Active plan</b>	<b>17,144 members</b> (includes dependents)	<b>\$111 million</b> total spend in FY13
<b>Retiree plan</b>	<b>64,237 members</b> (includes dependents) 40% live outside Alaska	<b>\$492 million</b> total spend in FY13
<b>Medicaid</b>	<b>145,279 Alaskans covered (2013)</b> 58% children, 36% adults, 6% seniors	<b>\$1.6 billion</b> total spend in 2013


The state also spends money on health care for inmates, state employees who are members of union health trusts and for state workers' compensation claims.

# Alaska's Health-Care Bill: \$7.5 Billion and Climbing

By Mark A. Foster and Scott Goldsmith


UA Research Summary No. 18 • August 2011

Institute of Social and Economic Research • University of Alaska Anchorage



Health-care spending for Alaskans reached about \$7.5 billion in 2010. For comparison, that's close to half the wellhead value of all the oil produced in Alaska that year. It's also roughly equal to half the wages Alaskans collected in 2010.

## Figure 7. What Do Alaska's Health-Care Dollars Buy? (2010 Spending: \$7.5 Billion)



\*Other personal and professional care and public health activities.

**Source:** Mark A. Foster and Associates estimates, based on Centers for Medicare and Medicaid Services, National Health Expenditure accounts

This is the way we currently contain costs:

1. Insurance focus on constraints to healthcare providers
  - A) limitations on coverage of diseases or conditions
  - B) limitations on compensation for procedures
  
2. Department of Labor work rules to prevent accidents
  - A) limitations on time on job or length of working day
  - B) safe working condition requirements
  
- 3). Department of Health and Social Services Programs to prevent disease
  - A) voluntary childhood vaccination
  - B) voluntary lifestyle change information
  - C) prescription drug safety
  - D) early detection of disease
  
- 4). DEC and DNR (Agriculture) - Food safety requirements
  - A) shellfish PSP testing etc.
  - B) pasteurized milk requirement etc.
  - C) restaurant and food processor inspections
  
- 5). Department of Public Safety - enforcement of laws
  - A) illegal drugs
  - B) smoking in close public places
  - C) driving under the influence of alcohol etc.



The Question:

How CAN WE AVOID diseases and PREVENT illness instead of just reacting to and paying for SICKCARE?

**The following slides demonstrate that we can reduce healthcare costs by initiating policies to avoid diseases with scientifically documented strategies.**



4th Quarter Report 2013

ACTIVE PLAN

## Aggregate Risk Profile

Member Information			
Member Count	17338	Avg Forecasted Cost	\$6,670
Avg Age	35	Avg Total Cost	\$6,774
Percent Female	51%	Avg Forecasted Risk Index	1.09
Avg Months Enrolled	11	%/w Acute Impact Score >= 95	1.06%
		%/w Chronic Impact Score >= 95	5.38%
		%/w Motivation Rank >= 95	4.83%

Aggregate Risk Summary				
Risk Drivers	# Members	Avg Risk Contribution	Contribution to Forecast	Risk Contribution
Demographics	17338	SENIORS SKIN, FRACTURES, FALLS \$345	\$5,987,784	5.18% <b>Y</b>
Acute Respiratory Disorders	2880	TUBERCULOSIS \$1,129	\$3,251,578	2.81% <b>X</b>
Arrhythmia Disorders	220	\$3,382	\$744,142	0.64%
CHF Conditions	673	CHRONIC HEART FAILURE \$3,181	\$2,140,732	1.85% <b>X</b>
Cerebral Vascular Disorder	247	\$4,199	\$1,037,177	0.90%
Chronic Respiratory Disorders	1728	UPPER RESPIRATORY TRACT \$2,318	\$4,005,542	3.46% <b>X</b>
Coronary Artery Related Conditions	1655	CORONARY HEART DISEASE \$2,958	\$4,895,538	4.23% <b>X</b>
Dermatological Disorder	3075	\$1,272	\$3,911,805	3.38%
Diabetic Disorders	882	TYPE 1 AND TYPE 2 \$5,932	\$5,231,998	4.52% <b>X</b>
Female Reproductive Conditions	341	PRETERM BIRTHS \$2,510	\$856,045	0.74% <b>X</b>
Gastrointestinal Disorders	2351	COLORECTAL CANCER \$2,011	\$4,728,854	4.09% <b>X</b>
Heart Related Conditions	180	\$5,628	\$1,013,046	0.88%
Hypertension	1527	BLOOD PRESSURE \$1,983	\$3,028,315	2.62% <b>X</b>
Hypotensive Drugs	1784	\$2,207	\$3,937,924	3.41%
Major Infection Related Conditions	2950	MRSA \$2,023	\$5,968,497	5.16% <b>X</b>
Metabolic Conditions	3077	FIBROMYALGIA \$2,680	\$8,247,421	7.13% <b>X</b>
Minor Infection Related Conditions	3704	\$1,340	\$4,965,037	4.29%
Miscellaneous Conditions	4750	\$2,303	\$10,940,696	9.46%
Musculo-skeletal Disorders	5173	INFANT MUSCLE, SENIOR FALLS \$2,206	\$11,409,047	9.87% <b>X</b>
Myocardial Infarction Related Conditions	271	\$5,315	\$1,440,328	1.25%
Neonatal Issues	255	ALITISM, HEART PROGRAMMING \$935	\$238,299	0.21% <b>X</b>
Neoplastic Related Conditions	638	\$4,332	\$2,763,900	2.39%
Neurological Disorder	3770	ALZHEIMER'S \$1,435	\$5,409,047	4.68% <b>X</b>
Non-specific condition	5561	\$140	\$780,877	0.68%
Pneumonia	243	UPPER RESPIRATORY TRACT \$3,822	\$928,744	0.80% <b>X</b>
Psychological Disorder	2688	S.A.D. AND DEPRESSION \$2,771	\$7,447,883	6.44% <b>X</b>
Renal Disorders	309	\$15,145	\$4,679,794	4.05%
Trauma Related Condition	1822	TRAUMATIC BRAIN INJURY \$1,455	\$2,651,019	2.29% <b>X</b>
Urinary Disorders	1381	\$2,170	\$2,996,360	2.59%

% total diseases directly related to Vitamin D status = 66.08 %

# RETIREE PLAN



4th Quarter Report 2013

## Aggregate Risk Profile


Member Information			
Member Count	65376	Avg Forecasted Cost	\$15,666
Avg Age	63	Avg Total Cost	\$17,726
Percent Female	54%	Avg Forecasted Risk Index	2.56
Avg Months Enrolled	12	%/w Acute Impact Score >= 95	5.47%
		%/w Chronic Impact Score >= 95	16.63%
		%/w Motivation Rank >= 95	9.70%

Aggregate Risk Summary				
Risk Drivers	# Members	Avg Risk Contribution	Contribution to Forecast	Risk Contribution
Demographics	65376	SENIORS SKIN, FRACTURES, FALLS \$454	\$29,702,943	2.90% <b>X</b>
Acute Respiratory Disorders	9520	TUBERCULOSIS \$1,669	\$15,893,121	1.55% <b>X</b>
Arrhythmia Disorders	5170	\$2,860	\$14,786,038	1.44%
CHF Conditions	10658	CHRONIC HEART FAILURE \$2,758	\$29,389,999	2.87% <b>X</b>
Cerebral Vascular Disorder	5021	\$3,726	\$18,710,595	1.83%
Chronic Respiratory Disorders	11241	UPPER RESPIRATORY TRACT \$3,093	\$34,763,411	3.39% <b>X</b>
Coronary Artery Related Conditions	24057	CORONARY HEART DISEASE \$2,900	\$69,776,210	6.81% <b>X</b>
Dermatological Disorder	15979	\$1,958	\$31,281,265	3.05%
Diabetic Disorders	10689	TYPE 1 AND TYPE 2 \$5,966	\$63,771,119	6.23% <b>X</b>
Female Reproductive Conditions	103	PRETERM BIRTHS \$1,489	\$153,413	0.01% <b>X</b>
Gastrointestinal Disorders	18753	COLORECTAL CANCER \$2,146	\$40,246,314	3.93% <b>X</b>
Heart Related Conditions	4346	\$4,374	\$19,007,254	1.86%
Hypertension	21394	BLOOD PRESSURE \$1,623	\$34,713,887	3.39% <b>X</b>
Hypotensive Drugs	21282	\$1,869	\$39,772,355	3.88%
Major Infection Related Conditions	13879	MRSA \$3,007	\$41,731,745	4.07% <b>X</b>
Metabolic Conditions	28763	FIBROMYALGIA \$2,545	\$73,213,862	7.15% <b>X</b>
Minor Infection Related Conditions	14339	\$1,726	\$24,754,410	2.42%
Miscellaneous Conditions	32888	\$2,762	\$90,833,634	8.87%
Musculo-skeletal Disorders	32886	INFANT MUSCLE, SENIOR FALLS \$2,498	\$82,154,255	8.02% <b>X</b>
Myocardial Infarction Related Conditions	5796	\$3,424	\$19,844,208	1.94%
Neonatal Issues	63	AUTISM, HEART PROGRAMMING \$1,504	\$94,751	0.01% <b>X</b>
Neoplastic Related Conditions	7447	\$3,928	\$29,253,071	2.86%
Neurological Disorder	27775	ALZHEIMER'S \$1,561	\$43,366,164	4.23% <b>X</b>
Non-specific condition	12687	\$113	\$1,429,752	0.14%
Pneumonia	1619	UPPER RESPIRATORY TRACT \$5,098	\$8,254,358	0.81% <b>X</b>
Psychological Disorder	16031	S.A.D. AND DEPRESSION \$2,659	\$42,623,390	4.16% <b>X</b>
Renal Disorders	4750	\$16,668	\$79,174,484	7.73%
Trauma Related Condition	8749	TRAUMATIC BRAIN INJURY \$1,904	\$16,660,268	1.63% <b>X</b>
Urinary Disorders	11515	\$2,504	\$28,836,425	2.82%

% total diseases directly related to Vitamin D status = 61.16%  
AlaskaCare Retiree Plan - 4th Quarter Report 2013

Notes in red by office of Rep. Seaton

# Meta-analysis of breast cancer risk



Dose–response gradient of risk of breast cancer according to serum 25-hydroxyvitamin D concentration, pooled analysis.

Active State Of Alaska employees, Retirees and dependents – **83,000**

**Female percentage of AK employees and retirees: 53% = 43,990**

Incidence of Breast Cancer per year in AK - **125 per 100,000 (.0125)**

Average cost of annual medical expenditures directly attributable to Breast  
Cancer – **\$11,000**

=


Per year AK State Cost for Breast Cancer: **\$ 6,048,625**

**50% reduction with vitamin D**

**Per Year Savings with vitamin D:**

**\$3,024,312**

# Meta-analysis



**Figure 1.** Dose–response gradient for colorectal cancer according to serum 25(OH)D concentration, all five studies combined.<sup>1,4–7</sup> The five points are the odds ratios for each quintile of 25(OH)D based on combined data from the five studies.

**Active State Of Alaska employees, Retirees and dependents – 83,000**

**Incidence of Colorectal Cancer per year in AK - 43 per 100,000 (.0043)**

**Average cost of annual medical expenditures directly attributable to Colon Cancer  
– \$11,000**

**=**

**AK State Cost for Colorectal Cancer per year \$ 3,925,900**

**50% per year savings with vitamin D**

**\$1,962,950**

**(meta-analysis Gorham et. al.)**

## Diabetes Incidence: Comparing NHANES and D\*action (18+ years)

In a comparison of data from the National Health and Nutrition Examination Survey (NHANES), 2005-2006, and GrassrootsHealth D\*action participants in the United States, we found the following:

Incidence of Diabetes:

NHANES: 8.5/1,000 person-years


D\*action: 0.9/1,000 person-years

A full 90% reduction in incidence - before adjusting for co-factors.

(Both groups had a similar average BMI, within 3 points.)


Rate Ratio = 9.7 (P=0.0002)

Chart Date: 8/6/13



# TYPE 2 DIABETES

Diabetes Incidence:  
Comparing NHANES\* and D\*action (18+ years)



NHANES Average 21 ng/ml

D\*Action Average 48 ng/ml

\*DATA SOURCE: NATIONAL HEALTH AND NUTRITION EXAMINATION SURVEY (NHANES), 2005-2006

\*\*RATE RATIO = 9.7 (P=0.0002)

Vitamin **D**\*action



**GrassrootsHealth**  
A Public Health Promotion Organization

[www.grassrootshealth.net](http://www.grassrootshealth.net)

Chart Date 8/6/13

© 2013 GrassrootsHealth

Preliminary data, not yet published.

**Active State Of Alaska employees, Retirees and dependents – 83,000**

**Employees, Retirees and dependents minus those with Diabetes already – 71,143**

**New incidences of diabetes per year – 8.5 per 1,000 per year (.0085)**

**Average cost of annual medical expenditures directly attributable to diabetes – \$7,900**

**=**

**Current Diabetes Cost per year= \$4,777,252**

**Per year Savings at 90% reduction = \$4,299,527**

**(GrassrootsHealth D\*Action study)**

**Per year Savings at 38% reduction = \$1,815,356**

**(Meta-analysis of prospective studies - Song et.al.)**

Select a summary ▼

search go

news: Fetal and

## Alaska

Find maternal and infant health data on a state level, or by county or city. Narrow your results or compare with another region.

Location: Alaska edit

---

Topic: Preterm by race/ethnicity edit

---


Format: Bar Graph edit

search reset

view tray

slides (0)


### Preterm by race/ethnicity: Alaska, 2009-2011 Average



- **Approximate number of births per year in Alaska = 11,000**
- **Assumption: 500 births per year to State of Alaska Employees, Retirees and Dependents**
- **For each 500 pregnancies in the Alaska insured and dependent category with vitamin D sufficiency 25 preterm births avoided**
- **Savings to the state by avoiding 25 preterm births = \$1,375,000**

## Upper Respiratory Tract Infections


Recently, a study was conducted with seven hundred forty-three children ages 3-15 in a Canadian Hutterite Community. **The findings of the study show that children with higher vitamin D blood levels had a 50% lower relative risk of contracting an Upper Respiratory Tract infection.** Those children at the United States national average of 21 ng/ml vitamin D levels were at a 70% greater risk of contracting respiratory infections. Illnesses such as RTI's are commonly a factor in children's absences from school. Making sure your child has sufficient vitamin D will not only increase their health, but will lead to less school absences due to illness.



*Low Serum 25 Hydroxyvitamin D level and Risk of Upper Respiratory tract infection in Children and Adolescents Science et. al. Journal of Clinical Infectious Diseases, August 2013 volume 57.*

Prepared by the office of Representative Paul Seaton

## Inflammatory gum disease



**HOUSE BILL NO. 356**

**IN THE LEGISLATURE OF THE STATE OF ALASKA**

**TWENTY-EIGHTH LEGISLATURE - SECOND SESSION**

**BY REPRESENTATIVE SEATON**

**Introduced: 2/26/14**

**Referred: Health and Social Services, State Affairs**

**A BILL**

**FOR AN ACT ENTITLED**

1 **"An Act establishing the Advisory Committee on Wellness; and relating to the**  
2 **administration of state group health insurance policies."**

3 **BE IT ENACTED BY THE LEGISLATURE OF THE STATE OF ALASKA:**

4 \* **Section 1.** The uncodified law of the State of Alaska is amended by adding a new section  
5 to read:

6 **LEGISLATIVE FINDINGS AND INTENT.** (a) The legislature finds that

7 (1) individual Alaskans and the state share the burden of increasing health care  
8 and insurance costs resulting from avoidable disease in humans;

9 (2) the state has not adequately focused state policy on avoiding disease and  
10 maintaining good health in the state's population;

11 (3) the state relies on federal one-size-fits-all human health policy  
12 recommendations, creating federal overreach into the state's administration of health policy;

13 (4) the state's human health policies have not incorporated peer-reviewed  
14 scientific studies that illustrate health-related differences between the population of the state


11 providing health care benefits to persons who are covered by a policy of group health  
12 insurance obtained under AS 39.30.090(a) or 39.30.091(a) to a rate that does not exceed two  
13 percent annually.

14 \* **Sec. 2.** AS 39.30.090(a) is amended by adding a new paragraph to read:

15 (13) To the greatest extent legally and reasonably practicable, the  
16 Department of Administration shall work to hold the escalation of health care costs to  
17 less than two percent annually by administering policies of group health insurance  
18 obtained under this subsection in a manner that is likely to reduce the incidence of  
19 disease in the state's population and that facilitates implementation of the  
20 recommendations of the Advisory Committee on Wellness established under  
21 AS 39.30.093.

22 \* **Sec. 3.** AS 39.30.091 is amended by adding a new subsection to read:

23 (b) To the greatest extent legally and reasonably practicable, the Department  
24 of Administration shall work to hold the escalation of health care costs to less than two  
25 percent annually by administering policies of group health insurance obtained under





KENAI PENINSULA BOROUGH SCHOOL DISTRICT

**Office of Superintendent**

Dr. Steve Atwater, Superintendent of Schools  
148 North Binkley Street Soldotna, Alaska 99669-7553  
Phone (907) 714-8888 Fax (907) 262-9132

March 6, 2014

Representative Paul Seaton  
State Capitol, Room 102  
Juneau, AK 99801-1182

Dear Representative Seaton,

I am writing on behalf of the Kenai Peninsula Borough School District to support House Bill 356. If passed, the bill will create an Advisory Committee on Wellness which will make recommendations to the Commissioner of Administration on ways to decrease the incidence of disease in Alaska. The intent of the bill is to implement policies to keep Alaska state employees healthy by preventing disease, rather than waiting to react once people are sick. KPBSD welcomes this proactive approach and recognizes that it is a necessary step in ensuring a healthy workforce as well as reducing the amount of funds dedicated to health care.

The cost of health care for my district's employees is increasing. In FY09 the total spent on employee health care was \$15,376,426 while last year it was \$21,541,380. This 40% increase in four years is alarming and unsustainable. KPBSD is thus pleased to support this bill which can help stem increases in district funds to health care.

Please contact me if you require further information for why KPBSD supports passage of HB 356.

Sincerely,

Steve Atwater, Ph.D.  
Superintendent of Schools

# Alaska State Legislature

State Capitol Room 102  
Juneau, Alaska 99801-1182  
(907) 465-2689  
Fax: (907) 465-3472  
1-800-665-2689




270 W Pioneer Ave. Suite B  
Homer, Alaska 99603  
(907) 235-2921  
Fax: (907) 235-4008  
1-800-665-2689

## REPRESENTATIVE PAUL SEATON HOUSE DISTRICT 30

### Memorandum

**From:** Representative Paul Seaton  
**To:** House State Affairs Committee Chairman, Representative Bob Lynn  
**DATE:** April 10, 2014  
**RE:** HB 356 Hearing Request



I respectfully request that HB 356 be heard in the House State Affairs Committee at your earliest possible convenience.

HB 356 requires that the Commissioner of the Department of Administration implement policies to reduce the incidence of disease in active and retired Alaska state workers to hold the inflation of health care costs to no more than 2% per year.

Attached to this request please find the following documents:

HB 356

HB 356 Sponsor Statement

HB 356 Background material in a single document but separated in categories as follows:

- Alaska Healthcare Costs
- Alaska Active and Retiree Health Plan Data
- Diabetes Cost
- Colorectal Cancer
- Breast Cancer
- Preterm Births
- Upper Respiratory Tract Infections
- Inflammation
- Muscle Function
- Senior Falls and Fractures
- Review of Recent Evidence on Vitamin D and Health
- Cost Savings per year with vitamin D

HB 356 Background material – Mayo Clinic “Vitamin D for Health: A Global Perspective”

HB 356 Support Letter – Kenai Peninsula Borough School District

HB 356 Zero Fiscal Note

Staff Contact: Louie Flora at 465-3923