

ALASKA LEGISLATURE COMMITTEE FILES 1987-1988 8672

5300 SJUD SB 32 (file 1)

872

reflex adjustments in the heart and systemic circulation. Finally, other possibilities, such as desensitization or blockade of peripheral adrenergic receptors, have not been examined.

Although the data on human beings are not adequate to determine how marijuana influences autonomic function, evidence that it does has been obtained. For example, Δ -9-THC appears to reduce a number of autonomic reflexes: After marijuana, the typical changes in heart rate and blood pressure elicited by the Valsalva maneuver (a forced exhalation effort against the closed glottis) are decreased, and so are the reflex circulatory responses to immersion of the hand in cold water (Beaconsfield et al., 1972; Benowitz et al., 1979). However, during chronic administration of Δ -9-THC, no change occurs in the reflex decrease in heart rate caused by infusion of a dose of the vasoconstrictor phenylephrine sufficient to increase the blood pressure (Benowitz and Jones, 1975; Benowitz et al., 1979).

Exercise

Acute exposure to Δ -9-THC modifies exercise performance by human beings. Smoking (20 mg of Δ -9-THC) decreased the duration of exercise but caused no change in any cardiopulmonary parameter at any work load except for heart rate, which increased (Shapiro et al., 1976b).

Other Effects (Plasma Volume, Sodium Retention)

Acute administration of Δ -9-THC would not be expected to have prominent effects on sodium balance or plasma volume. Chronic administration, on the other hand, has distinct effects. With chronic ingestion of large doses of Δ -9-THC there is a consistent gain in body weight and plasma volume, the latter caused by sodium retention (Benowitz and Jones, 1975, 1977a,b). The change in plasma volume seems to be causally related to the decrease in orthostatic hypotension during chronic exposure. The mechanisms responsible for the retention of salt and water have not been explored and may include changes in renal perfusion, inhibition of prostaglandin (a substance that affects blood pressure) synthesis by Δ -9-THC (Burststein and Raz, 1972; Howes and Osgood, 1976), or some modification in pituitary-adrenal function (Birmingham and Bartova, 1976).

Abnormal Heart and Circulation

Although smoking marijuana or the introduction of Δ -9-THC into the body is apparently without deleterious effect on the normal heart and circulation, the possibility is great that the abnormal heart and circulation will not be as tolerant of an agent that speeds up the heart, sometimes unpredictably raises or drops the blood pressure,

and modifies the activities of the autonomic nervous system. Therefore, it is pertinent to examine the prospects that marijuana (or Δ -9-THC) may be harmful in individuals with coronary heart disease, cerebrovascular disease, hypertension, and heart failure. Moreover, it may be important to determine if Δ -9-THC interacts in its effects on the abnormal heart or circulation with other agents that are being administered for therapeutic purposes.

Coronary Heart Disease

Data on this topic are sparse, presumably because of the relatively short time that marijuana has been available in this country. Those who have smoked marijuana are just entering the age when coronary atherosclerosis is common. However, it has been shown both in normal individuals and in individuals with coronary artery disease that the acute administration of Δ -9-THC by smoking or injection can cause changes in the electrocardiogram (ECG) (Johnson and DeMino, 1971; Beaconsfield et al., 1972; Kochar and Hosko, 1973). Premature beats have also been noted. The reasons for the changes are unclear. Also not understood is the contribution of the increase in heart rate itself to the ECG changes and to the premature beats.

In some patients with coronary artery disease, increased catecholamines can induce arrhythmias. It seems likely that in such patients Δ -9-THC could have the same effect. Also, in patients with coronary artery disease a large increase in heart rate can induce angina (pain) and even ischemic damage from insufficient oxygen as a result of an obstructed blood vessel. If Δ -9-THC were to increase heart rate markedly in such patients, and at the same time increase the need for cardiac perfusion because of the increased cardiac work and because of the intensified effect of catecholamines on the heart, it seems reasonable that there could be induction of angina and potentially precipitation of ischemic damage. Furthermore, if Δ -9-THC dulled the appreciation of pain and the appropriate responses to pain, the patient might not take suitable measure to relieve the angina, thereby increasing the risk of damage or arrhythmias.

A decrease in oxygen-carrying capacity of blood because of formation of carboxyhemoglobin could also be troublesome. Exercise tolerance has been reported to decrease in individuals with angina after smoking marijuana; this decrease is in contrast to the unaffected exercise tolerance after smoking a placebo marijuana cigarette (Aronow and Cassidy, 1974). Oral ingestion of Δ -9-THC or smoking marijuana apparently can cause marked hypertension in association with an increase in systemic vascular resistance (Benowitz et al., 1979), which would place the heart with coronary artery disease at risk of damage.

These observations concur in indicating that marijuana and Δ -9-THC increase the work of the heart, often in many ways. The conclusion seems inescapable that this increased work, coupled with stimulation by catecholamines, may tax the heart to the point of clinical hazard.

Cerebrovascular Disease

There are few, if any, indications that Δ -9-THC has direct effects on the cerebral circulation that would be important in patients with cerebrovascular disease. In the occasional patient who develops hypertension after smoking, there would be an increased risk of a cerebral vascular accident (stroke). Also, because Δ -9-THC administered after atropine can cause marked increases in blood pressure, this combination would place the patient with cerebrovascular disease at risk, as would smoking after ingestion of other muscarinic blockers. In some patients, postural hypotension could be a problem, not only for persons with abnormal cerebral circulations, but also with abnormal coronary circulations.

Hypertension

The factors that act to intensify angina would be of importance in hypertensive patients. Although data are lacking on the magnitude of change in blood pressure caused by Δ -9-THC in hypertensives, it seems reasonable to assume that hypertensives smoking marijuana might have a greater increase in blood pressure than normals do. The increase in plasma volume and sodium retention that are associated with chronic exposure to Δ -9-THC could increase blood pressure in hypertensives and the mechanisms responsible for these changes very likely would interfere with the action of a number of antihypertensive medications.

Heart Failure

Because marijuana can cause tachycardia, a decrease in systemic vascular resistance (required for increased cardiac output to sustain blood pressure) and salt and water retention might place patients with severe heart failure at a disadvantage by exposure to Δ -9-THC. Data on such patients are lacking. In older patients treated by Δ -9-THC or who have smoked marijuana for glaucoma or cancer, orthostatic hypotension has been both disabling and a threat of cardiovascular complications (Merritt et al., 1980). However, tolerance to orthostatic hypotension seems to develop during continued intake of Δ -9-THC or continued smoking of marijuana. Dehydration, as during vomiting or diuretic therapy, predisposes to the orthostatic hypotensive effects and resists the development of tolerance because it prevents expansion of blood volume.

Interactions with Cardioactive Drugs

Few studies evaluate interactions between Δ -9-THC and other drugs that act directly or indirectly on the heart. Propranolol usually attenuates the increase in heart rate caused by Δ -9-THC. Atropine

can greatly potentiate the ability of Δ -9-THC to increase systemic arterial pressure (Benowitz and Jones, 1977a,b). A number of possible interactions can be imagined. If a patient were taking a drug that blocked uptake of catecholamines by nerve terminals, then those effects of Δ -9-THC that are mediated by catecholamines would be intensified. Because a great many psychotropic and antihypertensive drugs modify metabolism of neurotransmitters in the central nervous system and periphery, a wide variety of interactions with Δ -9-THC seems possible.

Summary: Cardiovascular System

The smoking of marijuana causes changes in the heart and circulation that are characteristic of stress. But there is no evidence to indicate that it exerts a permanently deleterious effect on the normal cardiovascular system. Neither is there convincing evidence that marijuana would be of particular benefit in treating any of the major forms of cardiovascular disease.

The situation is quite different for those with an abnormal heart or circulation. Evidence abounds that marijuana increases the work of the heart, usually by increasing heart rate, and in some persons by increasing blood pressure. This increase in workload poses a threat to patients with hypertension, cerebrovascular disease, and coronary atherosclerosis. The magnitude and incidence of the threat remains to be determined because marijuana smoking has largely been confined to younger adults who are only now entering the age of serious complications of atherosclerosis on the heart, brain, and peripheral vessels.

Marijuana also can cause postural hypotension. This drop in blood pressure could be hazardous in those individuals with compromised blood flow to the heart or brain, especially if they are volume-depleted (dehydrated) or if other drugs have impaired reflex control of their blood vessels.

Marijuana appears to intensify the effects of the sympathetic nervous system on the heart, an undesirable consequence in patients with coronary artery disease and in those susceptible to arrhythmias. Many of the undesirable effects of marijuana on the cardiovascular system seem to become less severe following chronic exposure. Whether the relative paucity of reports of the ill-effects of marijuana on the abnormal cardiovascular system is a consequence of adaptation to chronic usage or to lack of exposure to marijuana of a population that is sufficiently advanced in years to be susceptible to its untoward effects remains to be determined.

Recommendations for Research

Additional studies are needed both (1) to provide information on the mechanisms responsible for the observed effects of marijuana on the cardiovascular system and (2) to provide new data on the effects of marijuana in patients with known forms of cardiovascular disease.

• The manner in which Δ -9-THC acts on the heart to change the rate and force of contraction needs clarification. Direct effects on the heart are not likely to differ among species, and thus experiments can be planned for a "standard" heart preparation.

• Direct effects on electrical activity, which might relate to reports of changes in electrical activity and production of premature impulses as well as changes in sinus rate, should be evaluated with standard methods and standard preparations.

• Direct effects of Δ -9-THC on vascular smooth muscle should be explored. For this purpose, it would be essential to use some vessels that did, and others that did not, have functioning nerve terminals. It would be important here to include studies on selected coronary vessels and on vessels which play a dominant role in the regulation of systemic vascular resistance.

• A number of related studies are needed before the effects on humans can be explained in full, particularly the effects of Δ -9-THC on the renin-angiotensin system in the kidney, which provides control of arterial pressure, and on the several sequences of prostaglandin metabolism.

Studies also are indicated to obtain new data about the effects of marijuana on:

• persons with hypertension, coronary artery disease, and cerebrovascular disease;

• increases in systemic arterial pressure in low- and high-renin hypertension and the interactions between Δ -9-THC and several classes of antihypertensive medications;

• the interactions between the salt and water-retaining effect of Δ -9-THC and diuretics that could be employed both in hypertensives and those with heart failure.

Additionally, studies should be done on the use of standard monitoring techniques to quantify any effect of marijuana smoking on tendencies toward arrhythmias, and on interactions of Δ -9-THC with drugs that modify synaptic transmission in the central nervous system.

REFERENCES

- Adams, M.D., Earnhardt, J.T., Dewey, W.L., and Harris, L.S. Vasoconstrictor actions of Δ -8- and Δ -9-tetrahydrocannabinol in the rat. J. Pharmacol. Exp. Ther. 196:649-656, 1976.
- Ames, F. A clinical and metabolic study of acute intoxication with cannabis sativa and its role in the model psychoses. J. Ment. Sci. 104:972-999, 1968.
- Aronow, W.S. and Cassidy, J. Effect of marijuana and placebo-marijuana smoking on angina pectoris. N. Engl. J. Med. 291:65-67, 1974.
- Auerbach, O., Stout, A.P., Hammond, E.C., and Garfinkel, L. Changes in bronchial epithelium relation to cigarette smoking and in relation to lung cancer. N. Engl. J. Med. 265:253-267, 1961.

- Beaconsfield, P., Ginsburg, J., and Rainsbury, R. Marijuana smoking: Cardiovascular effects in man and possible mechanisms. N. Engl. J. Med. 287:209-212, 1972.
- Bellville, J.W., Swanson, G.D., and Aqleh, K.A. Respiratory effects of Δ -9-tetrahydrocannabinol. Clin. Pharmacol. Ther. 17:541-548, 1975.
- Benowitz, N.L. and Jones, R.T. Cardiovascular effects of prolonged Δ -9-tetrahydrocannabinol ingestion. Clin. Pharmacol. Ther. 18:287-297, 1975.
- Benowitz, N.L. and Jones, R.T. Effects of Δ -9-tetrahydrocannabinol on drug distribution and metabolism. Clin. Pharmacol. Ther. 22:259-268, 1977a.
- Benowitz, N.L. and Jones, R.T. Prolonged Δ -9-tetrahydrocannabinol ingestion: Effects of sympathomimetic amines and autonomic blockades. Clin. Pharmacol. Ther. 21:336-342, 1977b.
- Benowitz, N.L., Rosenberg, J., Rogers, W., et al. Cardiovascular effects of intravenous Δ -9-tetrahydrocannabinol: Autonomic nervous mechanisms. Clin. Pharmacol. Ther. 25:440-446, 1979.
- Bernstein, J.G., Becker, D., Babor, T.F., and Mendelson, J.H. Physiological assessments: Cardiopulmonary function, pp. 147-160. In Mendelson, J.H., Rossi, A.M., and Meyers, R.E. (eds.) The Use of Marijuana: A Psychological and Physiological Inquiry. New York: Plenum Press, 1974.
- Birmingham, M.K. and Bartova, A. Effects of cannabinol derivatives on blood pressure, body weight, pituitary-adrenal function, and mitochondrial respiration in the rat, pp. 425-438. In Nahas, G.G. (ed.) Marihuana: Chemistry, Biochemistry, and Cellular Effects. New York: Springer-Verlag, 1976.
- Boulougouris, J.C., Panayiotopoulos, C.P., Antypas, E., et al. Effect of chronic hashish use on medical status in 44 users compared with 38 controls. Ann. N.Y. Acad. Sci. 282:168-172, 1976.
- Bright, T.P., Kiplinger, G.F., Brown, D., et al. Effects of beta-adrenergic blockade on marijuana-induced tachycardia, pp. 1737-1744. In Report of the 33rd Annual Scientific Meeting of the Committee on Problems of Drug Dependence. Vol. 2. Washington, D.C.: National Academy of Sciences, 1971.
- Burstein, S. and Raz, A. Inhibition of prostaglandin E_2 biosynthesis by Δ -1-tetrahydrocannabinol. Prostaglandins 2:369-374, 1972.
- Busch, F.W., Seid, D.A., and Wei, E.T. Mutagenic activity of marijuana smoke condensates. Cancer Lett. 6:319-324, 1979.
- Cavero, I., Buckley, J.P., and Jandhyala, B.S. Parasympatholytic activity of (-)-delta-9-tetrahydrocannabinol in mongrel dogs. Eur. J. Pharmacol. 19:301-304, 1972.
- Cavero, I., Solomon, T., Buckley, J.P., and Jandhyala, B.S. Studies on the bradycardia induced by (-)- Δ -9-tetrahydrocannabinol in anesthetized dogs. Eur. J. Pharmacol. 22:263-269, 1973.
- Dornbush, R.L. and Kokkevi, A. Acute effects of cannabis on cognitive, perceptual, and motor performance in chronic hashish users. Ann. N.Y. Acad. Sci. 282:313-322, 1976.

- Drath, D.B., Shorey, J.M., Price, L., and Huber, G.L. Metabolic and functional characteristics of alveolar macrophages recovered from rats exposed to marijuana smoke. Infect. Immun. 25:266-272, 1979.
- Englert, L.F., Ho, B.T., and Taylor, D. The effects of (-)- Δ -9-tetrahydrocannabinol on reserpine induced hypothermia in rats. Br. J. Pharmacol. 49:243-252, 1973.
- Fleischman, R.W., Baker, J.R., and Rosenkrantz, H. Pulmonary pathologic changes in rats exposed to marijuana smoke for 1 year. Toxicol. Appl. Pharmacol. 47:557-566, 1979.
- Gash, A., Karliner, J.S., Janowsky, D., and Lake, C.R. Effects of smoking marijuana on left ventricular performance and plasma norepinephrine: Studies in normal men. Ann. Intern. Med. 89:448-452, 1978.
- Gill, E.W. and Paton, W.D.M. Pharmacological experiments in vitro on the active principles of cannabis, pp. 165-173. In Joyce, C.R.B., Curry, S.H., London, J., and Churchill, A. The Botany and Chemistry of Cannabis, 1970.
- Glatt, H., Ohlsson, A., Agurell, S., and Oesch, F. Δ -1-tetrahydrocannabinol and 1-alpha, 2-alpha-epoxyhexahydrocannabinol: Mutagenicity investigation in the Ames test. Mutat. Res. 66:329-335, 1979.
- Graham, J.D.P. and Li, D.M.F. Cardiovascular and respiratory effects of cannabis in cat and rat. Br. J. Pharmacol. 49:1-10, 1973.
- Hall, J.A.S. Testimony in Marijuana-Hashish Epidemic Hearing of the Committee of the Judiciary U.S. Senate. Washington, D.C.: U.S. Government Printing Office, 1975.
- Hardman, H.F. and Losko, M.J. An overview of the cardiovascular-autonomic action of cannabis, pp. 231-238. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marijuana. New York: Raven Press, 1976.
- Henderson, R.L., Tennant, F.S., and Guerry, R. Respiratory manifestations of hashish smoking. Arch. Otolaryngol. 95:248-251, 1972.
- Hernandez-Bolanos, J., Swenson, E.W., and Coggins, W.J. Preservation of pulmonary function in regular, heavy, long-term marijuana smokers. Am. Rev. Resp. Dis. 113 (Suppl.):100, 1976.
- Ho, B.T., Taylor, D., and Englert, L.F. The effect of repeated administration of (-)- Δ -9-tetrahydrocannabinol on the biosynthesis of brain amines. Res. Commun. Chem. Pathol. Pharmacol. 5:851-854, 1973.
- Ho, B.T. and Johnson, K.M. Sites of neurochemical action of Δ -9-tetrahydrocannabinol: Interaction with reserpine, pp. 367-382. In Nahas, G.G. (ed.) Marijuana: Chemistry, Biochemistry, and Cellular Effects. New York: Springer-Verlag, 1976.
- Hoffmann, D., Brunnemann, K.D., Gori, G.B., and Wynder, E.L. On the carcinogenicity of marijuana smoke, pp. 63-81. In Runeckles, V.C. (ed.) Recent Advances in Phytochemistry New York: Plenum Publishing Corporation, 1975.
- Howes, J. and Osgood, P. The effect of Δ -9-tetrahydrocannabinol on the uptake and release of ^{14}C -dopamine from crude striatal synaptosomal preparations. Neuropharmacology 13:1109-1114, 1974.

- Howes, J.F. and Osgood, P.F. Cannabinoids and the inhibition of prostaglandin synthesis, pp. 415-424. In Nahas, G.G. (ed.) Marihuana: Chemistry, Biochemistry, and Cellular Effects. New York: Springer-Verlag, 1976.
- Huber, G.L., Simmons, G.A., McCarthy, C.R., et al. Depressant effect of marihuana smoke on antibactericidal activity of pulmonary alveolar macrophages. Chest 68:769-773, 1975.
- Huber, G.L., Pochay, V.E., Shea, J.W., et al. An experimental animal model for quantifying the biologic effects of marijuana on the defense system of the lung, pp. 301-328. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979a.
- Huber, G.L., Shea, J.W., Hinds, W.E., et al. The gas phase of marijuana smoke and intrapulmonary bactericidal defenses. Bull. Eur. Physiopath. Resp. 15:491-503, 1979b.
- Huber, G.L., Pochay, V.E., Pereira, W., et al. Marijuana, tetrahydrocannabinol, and pulmonary antibacterial defenses. Chest 77:403-410, 1980.
- Huot, J. Cellular and biochemical alterations induced in vitro by delta-1-tetrahydrocannabinol: Effects on cell proliferation, nucleic acids, plasma cell membrane ATPase, and adenylate cyclase, pp. 313-327. In Nahas, G.G. (ed.) Marihuana: Chemistry, Biochemistry, and Cellular Effects. New York: Springer-Verlag, 1976.
- Joachimoglu, G. Natural and smoked hashish, pp. 2-14. In Wolstenholme, G.E.W., Knight, J., London, J., and Churchill, A. Hashish: Its Chemistry and Pharmacology (CIBA Foundation Study Group 21), 1965.
- Johnson, S. and Domino, E.F. Some cardiovascular effects of marihuana smoking in normal volunteers. Clin. Pharmacol. Ther. 12:762-768, 1971.
- Kanakis, C.J., Pouget, J.M., and Rosen, K.M. The effects of Δ -9-tetrahydrocannabinol (cannabis) on cardiac performance with and without beta blockage. Circulation 53:703-707, 1976.
- Kawasaki, H., Watanabe, S., Oishi, R., and Ueki, S. Effects of Δ -9-tetrahydrocannabinol on the cardiovascular system, and pressor and behavioral responses to brain stimulation in rats. Jpn. J. Pharmacol. 30:493-502, 1980.
- Kochar, M.S. and Hosko, M.J. Electrocardiographic effects of marihuana. JAMA 225:25-27, 1973.
- Leuchtenberger, C. and Leuchtenberger, R. Morphological and cytochemical effects of marijuana cigarette smoke on epithelioid cells of lung explants from mice. Nature 234:227-229, 1971.
- Leuchtenberger, C., Leuchtenberger, R., and Schneider, A. Effects of marijuana and tobacco smoke on human lung physiology. Nature 241:137-139, 1973a.
- Leuchtenberger, C., Leuchtengerger, R., Ritter, U., and Inui, N. Effects of marijuana and tobacco smoke on DNA and chromosomal complement in human lung explants. Nature 242:403-404, 1973b.

- Leuchtenberger, C. and Leuchtenberger, R. Cytological and cytochemical studies of the effect of fresh marijuana cigarette smoke on growth and DNA metabolism of animal and human lung cultures, pp. 595-612. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marijuana. New York: Raven Press, 1976.
- Loewe, S. Pharmacological study, pp. 149-220. In The Marijuana Problem in the City of New York: Sociological, Medical, Psychological and Pharmacological Studies. Lancaster, Pa.: Jacques Cattall Press, 1944.
- Malit, L.A., Johnstone, R.E., Bourke, D.I., et al. Intravenous Δ -9-tetrahydrocannabinol: Effects on ventilatory control and cardiovascular dynamics. Anesthesiology 42:666-673, 1975.
- Martz, R., Brown, D.J., Forney, R.B., et al. Propranolol antagonism of marijuana induced tachycardia. Life Sci. 11:999-1005, 1972.
- Merritt, J.C., Crawford, W.J., Alexander, P.C., et al. Effect of marijuana on intraocular and blood pressure in glaucoma. Ophthalmology 87:222-228, 1980.
- Nowlan, R. and Cohen, S. Tolerance to marijuana: Heart rate and subjective "high." Clin. Pharmacol. Ther. 22:550-556, 1977.
- Perez-Reyes, M., Lipton, M.A., Timmons, M.C., et al. Pharmacology of orally administered Δ -9-tetrahydrocannabinol. Clin. Pharmacol. Ther. 14:48-55, 1973.
- Rosenkrantz, H. and Braude, M.C. Acute, subacute and 23-day chronic marijuana inhalation toxicities in the rat. Toxicol. Appl. Pharmacol. 28:428-441, 1974.
- Roy, P.E., Magnan-Lapointe, F., Huy, N.D., and Boutet, M. Chronic inhalation of marijuana and tobacco in dogs: Pulmonary pathology. Res. Commun. Chem. Pathol. Pharmacol. 14:305-317, 1976.
- Rubin, V. and Comitas, L. Ganja in Jamaica: A Medical Anthropological Study of Chronic Marijuana Use. The Hague: Mouton and Co., 1975.
- Seid, D.A. and Wei, E.T. Mutagenic activity of marijuana smoke condensates. Pharmacologist 21:204, 1979.
- Shapiro, B.J., Tashkin, D.P., and Frank, I.M. Mechanism of increased specific airway conductance with marijuana smoking in healthy young men. Ann. Int. Med. 78:832-833, 1973.
- Shapiro, B.J., Tashkin, D.P., and Frank, I.M. Effects of beta-adrenergic blockade and muscarinic stimulation upon cannabis bronchodilation, pp. 277-286. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marijuana. New York: Raven Press, 1976a.
- Shapiro, B.J., Reiss, S., Sullivan, S.F., Tashkin, D.P., Simmons, M.S., Smith, R.T. Cardiopulmonary effects of marijuana smoking during exercise. Chest 70:441, 1976b.
- Stefanis, C. Biological aspects of cannabis use, pp. 149-178. In Petersen, R.C. (ed.) The National Challenge of Drug Abuse. NIDA Research Monograph No. 19. HEW Publication No. (ADM)78-654. Washington, D.C.: U.S. Government Printing Office, 1978.
- Sulkowski, A., Vachon, L., and Rich, E.S., Jr. Propranolol effects on acute marijuana intoxication in man. Psychopharmacology 52:47-53, 1977.

- Tashkin, D.P., Shapiro, B.J., and Frank, I.M. Acute pulmonary physiologic effects of smoked marijuana and oral Δ -9-tetrahydrocannabinol in healthy young men. N. Engl. J. Med. 289:336-341, 1973.
- Tashkin, D.P., Shapiro, B.J., and Frank, I.M. Acute effects of smoked marijuana and oral Δ -9-tetrahydrocannabinol on specific airway conductance in asthmatic subjects. Am. Rev. Respir. Dis. 109:420-428, 1974.
- Tashkin, D.P., Shapiro, B.J., Lee, Y.E., and Harper, C.E. Effects of smoked marijuana in experimentally induced asthma. Am. Rev. Respir. Dis. 112:377-386, 1975.
- Tashkin, D.P., Shapiro, B.J., Lee, E.Y., Harper, C.E. Subacute effects of heavy marijuana smoking pulmonary function in healthy young males. N. Engl. J. Med. 294:125-129, 1976.
- Tashkin, D.P., Reiss, S., Shapiro, B.J., et al. Bronchial effects of aerosolized Δ -9-tetrahydrocannabinol in healthy and asthmatic subjects. Am. Rev. Respir. Dis. 115:57-65, 1977a.
- Tashkin, D.P., Levisman, J.A., and Abbasi, A.S. Short-term effects of smoked marijuana on left ventricular function in man. Chest 72:20-26, 1977b.
- Tashkin, D.P., Soares, J.R., Hepler, R.S., Shapiro, B.J., and Rachelefsky, G.S. Cannabis, 1977. Ann. Int. Med. 89:539-549, 1978.
- Tashkin, D.P., Calvarese, B.M., Simmons, M.S., and Shapiro, B.J. Respiratory status of seventy-four habitual marijuana smokers. Chest 78:699-706, 1980.
- Tennant, F.S., Preble, M., Prendergast, T.J., and Ventry, P. Medical manifestations associated with hashish. JAMA 216:1965-1969, 1971.
- Tennant, F.S., Guerry, R.L., and Henderson, R.L. Histopathologic and clinical abnormalities of the respiratory system in chronic hashish smokers. Substance and Alcohol Misuse 1:93-100, 1980.
- Truitt, E.B. and Anderson, S.M. Biogenic amine alterations produced in the brain by tetrahydrocannabinols and their metabolites. Ann. N.Y. Acad. Sci. 191:68-73, 1971.
- Vachon, L., FitzGerald, M.X., Solliday, N.H., et al. Single-dose effect of marijuana smoke: Bronchial dynamics and respiratory-center sensitivity in normal subjects. N. Engl. J. Med. 288:985-989, 1973.
- van Went, G.F. Mutagenicity testing of 3 hallucinogens: LSD, psilocybin and Δ -9-THC, using the micronucleus test. Experientia 34:324-325, 1978.
- Vollmer, R.R., Caverio, I., Ertel, R.J., et al. Role of the central autonomic nervous system in the hypotension and bradycardia induced by (-)- Δ -9-trans-tetrahydrocannabinol. J. Pharm. Pharmacol. 26:186-192, 1974.
- Wehner, F.C., Van Rensburg, S.J., and Thiel, P.G. Mutagenicity of marijuana and transkei tobacco smoke condensates in the salmonella/microsome assay. Mutat. Res. 77:135-142, 1980.
- Weil, A.T., Zinberg, N.E., and Nelsen, J.M. Clinical and psychological effects of marijuana in man. Science 162:1234-1242, 1968.

Weiss, J.L., Watanabe, A.M., Lemberger, L., et al. Cardiovascular effects of Δ -9-tetrahydrocannabinol in man. Clin. Pharmacol. Ther. 13:671-684, 1972.

Zwillich, C.W., Doekel, R., Hammill, S., and Weil, J.V. The effects of smoked marijuana on metabolism and respiratory control. Am. Rev. Respir. Dis. 118:885-891, 1978.

EFFECTS OF MARIJUANA ON THE BRAIN

The most clearly established effects of cannabis are upon behavior. These effects, described in Chapter 6, indicate that major actions of cannabinoids are upon the brain. The ways in which marijuana alters the brain to produce its behavioral effects are not known.

Efforts to discover the causes of the behavioral effects have included studies on brain morphology, physiology, and chemistry to be reviewed in this chapter. Effects of marijuana on brain electrical activity and on brain chemistry have been measured, but their significance for brain function is not known because of our limited knowledge of brain-behavior relations. Marijuana causes temporary intoxication and results in changes in brain physiology and chemistry similar to those caused by other intoxicating drugs. Although these kinds of studies may ultimately shed light on the way marijuana produces its behavioral changes, they do not provide answers to important clinical questions. Does marijuana cause long-term changes in the brain that lead to chronic psychiatric or neurological disorders? So far, the studies reviewed below provide no convincing evidence for long-term changes because of use of marijuana.

BRAIN MORPHOLOGY

There is substantial controversy about whether marijuana causes changes in brain structure or in brain cells. Two studies have reported that marijuana produces changes in brain morphology. Both suffer sufficiently from methodologic and interpretational defects that their conclusions cannot be accepted. Furthermore, other studies have not found changes in morphology.

Gross Morphology

Data suggesting that use of marijuana causes brain atrophy were obtained by pneumoencephalography (injection of air into spaces in and surrounding the brain) on 10 users of marijuana who had sought medical attention because of neurologic complaints (Campbell et al., 1971). The size of the largest brain cavities (ventricles) was

measured to determine whether loss of brain tissue had occurred. The authors interpreted their data as showing that atrophy was present.

One of the first critics of this report questioned the interpretation of the radiologic techniques used (Bull, 1971). The results also have been seriously criticized because of the marijuana users studied. They had neurological symptoms or signs sufficient to justify an invasive and painful diagnostic test, but there is no evidence that such neurological complaints occur with greater frequency in users of marijuana than in the general population. Further, Campbell's patients did not only use marijuana, but also used such behavior-altering drugs, as lysergic acid diethylamide (LSD) and amphetamines.

More recent evidence has been provided by computed tomography (CT) scans of the brain. This technique, which is noninvasive, painless, and yields more precise and quantifiable measures of brain atrophy, has replaced pneumoencephalography as a diagnostic test. Using CT methods, two studies failed to find evidence of cerebral atrophy in healthy chronic marijuana users (Co et al., 1977; Keuhle et al., 1977). These latter results suggest that the earlier findings were attributable to the imprecision of conventional pneumoencephalography, or to the fact that a group with neurologic complaints was studied, or to the use of multiple psychoactive drugs by these individuals. This last possibility is reinforced by CT scans of animals who received a variety of psychoactive drugs. Marijuana alone produced no evidence of brain atrophy, whereas other drugs, such as amphetamines, did produce changes (Rumbaugh et al., 1980).

Microscopic Morphology

Three post mortem studies on monkeys in the same laboratory have reported changes in the microscopic morphology of the brain at the ultrastructural level (Harper et al., 1977; Meyers and Heath, 1979; Heath et al., 1980). No similar studies on human beings have been reported. The monkeys received either chronic exposure to marijuana smoke or chronic injections of Δ -9-THC. Changes reported to have occurred in the brains included alteration in synaptic* cleft width, increased density of synaptic cleft material, a decrease in volume of rough endoplasmic reticulum, presence of clumping of synaptic vesicles in axon terminals (where impulses travel away from the cell body), and an increase in intranuclear inclusions. These changes appear dramatic, but they must be interpreted with caution. The three studies are based principally upon examination of two limited brain areas only in three treated monkeys, two receiving marijuana smoke

*A synapse is the region of communication between nerve cells, forming the place where a nervous impulse is transmitted from one nerve cell to another.

and one intravenous Δ -9-THC; a fourth treated animal was added to the last study and more brain areas were analyzed in it (Heath et al., 1980). Further, although the material was evaluated "doubleblind" after electron micrographs had been made, it would appear that fixation, tissue preparation, and photography were carried out before these safeguards against bias were applied. It is possible that unknown but systematic differences occurred between experimental (treated) and control animals in fixation and preparation of tissue or in selection of samples for micrography. In addition, it should be noted that at least one of the changes noted, clumping of synaptic vesicles (Harper et al., 1977), is a normal variant in the synaptic morphology of axon terminals in mammalian brain (Sipe and Moore, 1977) and does not represent a pathological change. Also, these studies have not been replicated and, because the basis for interpretation is such a limited sample, it is concluded that no definitive interpretation can be made at this time. However, the possibility that marijuana may produce chronic, ultra-structural changes in brain has not been ruled out and should be investigated.

NEUROPHYSIOLOGY

One source of information on the mechanisms of action of a drug, such as marijuana, is the study of its physiological effects. Effects of marijuana on the electrical activity of the brain have been demonstrated by means of the electroencephalogram (EEG). The standard, or clinical, EEG measures tiny variations at the scalp of voltages produced by the electrical activity of the brain. Voltage differences between two points on the scalp, or between the scalp and an inactive reference site, are recorded on moving paper, producing a graph of voltage over time. The waves observed are classified according to frequencies as delta, theta, alpha, and beta. While the changes in EEG described below are of interest, their biological significance is unknown.

Acute (Short-Term) Effects in Waking EEG

Ingested marijuana or Δ -9-THC produces rather slight effects on the EEG of an awake subject. Relatively high doses (210 mg Δ -9-THC or its equivalent/day) have failed to produce measurable changes even though marked behavioral effects were observed. The EEG effect most frequently reported in recent studies has been an increased abundance of alpha waves associated with a slight slowing (about 0.25 Hz) of the alpha frequency (Rodin et al., 1970; Volavka et al., 1971; Fink, 1976). However, reduced alpha abundance and increased fast frequency activity (beta) have also been reported (Wikler and Lloyd, 1945; Jones and Stone, 1970). Most studies which report EEG changes have noted that tolerance develops with repeated drug administration. No significance with respect to hazard can be inferred from the effects

of cannabis on the waking EEG. For a further review of this literature, see Fried (1977).

Persistent Effects in Waking EEG

The occurrence of persistent (long-lasting) changes in EEG with use of marijuana would cause concern even if their significance for brain function was unknown. However, in attempting to investigate the question of whether such changes occur, there inevitably arise crucial issues of subject selection. If one selects only chronic marijuana users who are in good health, one may be eliminating systematically those who have been adversely affected by use of the drug and who might have shown EEG changes. On the other hand, if one includes in such studies marijuana users who suffer from various illnesses or behavioral disturbances, one might find abnormalities of the EEG that result from these conditions rather than from the marijuana.

Long-term use of marijuana, either in the modest doses customarily used in this country or the heavy doses of hashish and ganga used by certain studied populations abroad, has not been shown to produce changes in the EEG. No abnormalities were found in the EEG of 10 healthy students who had smoked marijuana regularly for 1 year (Rodin et al., 1970). Another study compared clinical EEG records of 40 hashish users and 40 matched controls in Greece (Fink, 1976). Each record was evaluated independently by four qualified neurologist-electroencephalographers. No differences were observed in the incidence of abnormal records in the users and controls, a result consistent with the absence of significant differences between the two groups in various tests of neurological function.

Essentially, the same negative results were obtained in studies of ganga users in Jamaica (Rubin and Comitas, 1975) and marijuana users in Costa Rica (Karacan et al., 1976). In these later studies subjects were carefully selected to include only those in good health who were functioning adequately in the community. As mentioned above, this method of selection runs the risk of eliminating subjects whose health or behavior were adversely affected by marijuana and who might have shown EEG changes. This methodological difficulty cannot be eliminated in any small sample investigation of marijuana users.

Acute Effects in Event-Related Potentials

One can employ computer averaging to retrieve from the EEG certain information that is not detectable by visual inspection. In this way, the electrical events that follow a stimulus may be studied in subjects who are at rest, asleep, or carrying out certain tasks. These computer-averaged potentials provide clues to the sequential processing of information by the brain.

Although the literature is inconsistent, it is clear that cannabis can produce effects on event-related potentials (EPs) (Herning et al., 1979). Effects on amplitude are more often reported than effects

on latency of the event-related waves. Several studies with inconsistent results have appeared; these inconsistencies result from differences in task, dose, or duration of administration. Thus, EPs in response to sensory stimuli are unaffected or even increased by cannabis if the subject is passive, but are decreased in amplitude if the subject is performing a task. One study found the first negative wave, a component of the auditory EP, was reduced at a dose of 180-210 mg per day, but not at a dose of 70-90 mg per day during acute (1 to 3 days) administration (Herning et al., 1979). After 2 weeks at the higher dosage, this effect was observed only for the more difficult tasks. This study demonstrates differences in marijuana effects on EPs according to dose, duration of administration, and task complexity.

Acute Effects in Sleep EEG

Drugs often produce marked effects on the EEG during sleep, but producing little or no change in the waking EEG. This is the case with marijuana and Δ -9-THC.

In relatively high doses (70-210 mg/day), Δ -9-THC and marijuana extract produced marked effects on sleep EEG (Fainberg et al., 1975, 1976). On initial administration, the time spent in REM sleep* (stage REM duration) was reduced below baseline levels (placebo) by 18 percent and the number of eye movements by 49 percent. Some tolerance (return toward baseline levels) was apparent during the period (12-16 days) of drug administration. On withdrawal, REM duration was increased above baseline by 49 percent and rapid eye movements were increased by 67 percent. While these effects are quite large, their clinical significance is unknown. They were not accompanied by such unusual behavioral changes as hallucinations or disorientation, although there was evidence of withdrawal--irritability, increased reflexes, and mild agitation. With much smaller doses of Δ -9-THC, either a small reduction in REM sleep (Pivik et al., 1972; Freeman, 1974) or no change has been reported (Barratt et al., 1974; Hosko et al., 1973; Prankoff et al., 1973).

Persistent Effects in Sleep EEG

We are not aware of any investigation of sleep in abstinent long-term marijuana users. However, 32 male chronic marijuana users and matched controls were studied in Costa Rica (Karacan et al., 1976). The users habitually smoked 2.5 to 23.3 cigarettes per day (mean = 9.2) and had used the drug for 10 to 27 years; they continued their usual intake during the study. (Costa Rican cigarettes contain about 200 mg

*A stage in sleep during which Rapid Eye Movements may be detected and vivid dreaming usually occurs.

of marijuana). The subjects selected for this study had normal medical, neurologic, and laboratory evaluations.

Sleep was recorded for 8 consecutive nights. Prior to each night's recording, the users described their marijuana intake during the previous 24 hours. This intake was not directly monitored or controlled by the experimenters, because the goal was to observe sleep patterns under "naturalistic" conditions. The subjects were forbidden to use marijuana during the 2-3 hours prior to sleep recording. (For further details of this extensive study, see Karacan et al., 1976.)

All of the major variables derived from visual sleep stage classification were examined. The only statistically significant differences between marijuana users and their matched controls were in one of the sleep latency measures and in REM percentage of total sleep and average REM period length. The differences were quite small and may have been due to the subjects experiencing early withdrawal at the time their sleep was recorded. This is a likely explanation for these findings according to studies described previously (Feinberg et al., 1975, 1976).

The Costa Rican study concluded there was a lack of evidence of... major disturbances of EEG sleep patterns in user subjects studied in situ (Karacan et al., 1976). Thus, long-term marijuana use has not been demonstrated to cause marked and consistent abnormalities of sleep EEG that can be demonstrated in studies with small samples.

Electrophysiological Studies in Animals

Sleep Studies

The findings of several animal studies carried out to investigate the effects of marijuana on EEG differ in some respects to those in human beings. Species differences are thought to be responsible for some of the variations found from species to species. For example, 5 and 10 mg/kg Δ -9-THC administered acutely to rats suppressed REM, reduced slow-wave sleep, and increased wakefulness (Moreton and Davis, 1973). Chronic administration caused an initial suppression of REM, which returned to baseline after 4 days and remained at baseline levels for a further 16 days. In contrast to the human studies, there was no withdrawal increase in REM above baseline during a 10-day withdrawal period. Similar results were obtained in a short-term study that employed intravenous doses of Δ -9-THC (0.5 and 1.0 mg/kg) to rabbits (Fujiimori and Himwich, 1973).

Appreciable qualitative differences in sleep EEG response to Δ -9-THC have also been detected in primates when compared with human studies. When 1.2 mg/kg Δ -9-THC is administered to squirrel monkeys in a single oral dose, daily for 60 days, no significant effects on REM sleep duration occurred; instead, a decrease in EEG stages 3 and 4 was noted (Adams and Barratt, 1975).

EEG Studies in Subcortical Structures

Electrode implantation is rarely possible in man, but is a routine and essential technique for the study of brain electrophysiology in animals. Animal experiments also permit use of higher doses and more prolonged administration than is possible with human subjects. For these reasons, animal experiments can yield important data that cannot be obtained in human studies. In general, EEG recordings after short-term administration of marijuana are similar from surface (cortex) or from deep brain (subcortex) regions. However, after chronic administration of high doses of Δ -9-THC, abnormal recordings have been observed in subcortical regions of some animals, readings not seen in the cortex. Although these findings have not been replicated, they are of particular concern, because they raise the possibility that chronic exposure to high doses of marijuana produces long-lasting effects on brain physiology.

After intravenous administration of a range of Δ -9-THC doses (from 0.05 to 12.8 mg/kg) to rhesus monkeys, a general increase in EEG synchrony was observed; and at higher dose ranges, there were specific EEG changes in the limbic system, frontal cortex, thalamus and fastigial nuclei (Martinez et al., 1972). In this study, the increase in high-voltage activity showed a good dose-response relationship. In a second study, oral dosing of three rhesus monkeys with a crude marijuana extract containing 25 percent Δ -9-THC produced dose-related EEG changes, including slow waves in the hippocampus, amygdala, and septum (Stadnicki et al., 1974). Tolerance to the behavioral and EEG changes occurred with daily treatment, which was stopped after 51 days. Behavioral withdrawal effects were noted, but EEG changes during withdrawal were minimal and there was no evidence of EEG changes persisting beyond the period of Δ -9-THC ingestion.

Two studies that monitored EEG recording from deep brain sites after chronic administration of high doses of marijuana found changes in EEGs from deep brain sites that were not observed in surface areas after drug withdrawal (Fehr et al., 1976; Heath, 1976; Heath et al., 1979). Studies of two rats with electrodes implanted in the anterior neocortex, dorsal hippocampus, and mesencephalic reticular formation 1 year after exposure to 20 mg/kg for 6 months (Fehr et al., 1976) yielded hippocampal recordings with "epileptiform" abnormalities, in contrast to one control and two alcohol-treated animals.

The second study was carried out on thirteen feral-raised rhesus monkeys (Heath 1976; Heath et al., 1979). Ten monkeys had electrodes implanted in deep sites and in brain cortex. Four monkeys were made to smoke marijuana three times a day, 5 days per week for 6 months; two other monkeys with implants were given 0.6 mg/kg Δ -9-THC each day, 5 days per week for 6 months; still other monkeys were used as controls or received smaller doses of marijuana. In three high-dose monkeys, two smoking and one ingesting Δ -9-THC, changes in EEG could be detected in recordings from deep brain sites; the changes continued 7 months after cessation of marijuana exposure. No EEG abnormalities were present in recordings from the brain surface.

One of the major criticisms of both these studies is their use of small numbers of animals. Furthermore, there have been no attempts at replication by other workers. Nevertheless, because these findings provide some of the only evidence for a possible irreversible effect of chronic high doses of marijuana, they are mentioned here with a strong urging for additional studies in an effort to replicate these findings.

EPILEPSY

Because of the effects of marijuana on brain electrical activity, questions have been raised about its association with epilepsy. Two questions are raised in the literature. First, does marijuana produce seizures? Second, does marijuana or a derivative prevent seizures? The first question will be discussed here. The second is reviewed in Chapter 7, which is concerned with the potential therapeutic uses of cannabis.

There are anecdotal reports in the literature that suggest seizures may be induced by marijuana in some persons with a known seizure disorder. A rigorous study, using adequate numbers of patients with documented seizure patterns, has not been done. Reports of experimental animal studies are conflicting and varied (Feeney et al., 1973, 1979; Lemberger, 1980). There are some circumstances in which cannabis administration does not alter certain types of seizures such as the photosensitive seizures in the baboon (Meldrum et al., 1974), and others in which it seems that seizures are induced. A single rabbit that responded to Δ -9-THC administration with seizures was bred to establish a colony of rabbits with similar response (Consroe and Fish, 1981). It will be of considerable interest to determine mechanisms of seizure induction and pharmacologic response patterns in this unusual animal model. However, as described further in Chapter 7, the bulk of the animal literature and some data from human studies suggest that the more prominent effect of marijuana derivatives, especially cannabidiol and cannabidiol, is to decrease rather than increase seizure susceptibility (see Karler and Turkhan, 1981, for review).

NEUROCHEMISTRY

Our knowledge of the effects of marijuana on brain chemistry has come largely from studies in animals. Cannabis and some of its derivatives have been shown to cause chemical effects in the brain, as demonstrated by effects on neurotransmitters and on nucleic acids. The evidence is reviewed below.

Neurotransmitters

The brain is composed of many information-processing networks of nerve cells. Within each of these networks the transfer of informa-

tion from one nerve cell to another is dependent upon chemicals called neurotransmitters. These substances are produced by nerve cells, released when the cells are stimulated and act to alter the excitability of neighboring nerve cells. Neurotransmitters play an essential role in the transmission and processing of information, and it is not surprising that many drugs that alter behavior do so by their actions on neurotransmitters. The understanding of the effects of marijuana on the brain must include knowledge of its effects on neurotransmitter systems.

Several different classes of chemicals act as neurotransmitters. The first chemical to be demonstrated to have this function was acetylcholine, and it is now established that acetylcholine is the neurotransmitter for several nerve cell networks in the brain. A number of studies in animals have examined the effect of marijuana on brain acetylcholine (see Domino, 1981, for a brief review of the extensive literature). The most clear-cut effects have been on acetylcholine turnover, a measure of the level of activity of neurons producing the chemical. Small doses of Δ -9-THC cause a reduction in acetylcholine turnover in the hippocampus (Domino et al., 1978; Revuelta et al., 1978; Domino, 1981) and this results from reduced activity of the acetylcholine neurons. It is noteworthy that the effect is produced by small doses and only by cannabinoids. Administration of physostigmine, a drug that enhances acetylcholine action by partially blocking its breakdown, to five healthy human volunteers (2 hours after ingestion of 20 to 40 mg of Δ -9-THC) produced enhancement of the lethargy and somnolence occurring late in the course of the Δ -9-THC intoxication (Freemon et al., 1975). The results of this study, and others in man and animals (El-Yousef et al., 1973; Low et al., 1973; Drew and Miller, 1974; Freemon et al., 1975), have led to the conclusion that Δ -9-THC acts to inhibit acetylcholine nerve cell networks. The exact nature of this action is not known, but it may be related to the memory deficits produced (Domino, 1981).

There have been studies of cannabinoids on several other neurotransmitters in brain, including catecholamines, serotonin, and gamma aminobutyric acid (Banerjee et al., 1975; Bracs et al., 1975). Although some effects have been reported, they either are produced by a very high dose or are so fragmentary that their implications are unclear. The effects of cannabinoids on neurotransmitters that have been studied to date, other than acetylcholine, are not striking. In particular, there is no evidence for any significant, long-term toxic effect of cannabinoids on any of the nerve cell networks that produce identified neurotransmitters.

Proteins, Enzymes, Nucleic Acids

A very few studies have examined the effects of marijuana on neurochemical variables other than neurotransmitters (Luthra and Rosenkrantz, 1974; Luthra et al., 1975, 1976). After chronic administration to rats either of Δ -9-THC or marijuana smoke (for

periods from 28 to 180 days), these investigators examined brain lipid, protein, and ribonucleic acid (RNA) content. With very high doses of Δ -9-THC (up to 500 mg/kg/day), some decrease in brain protein and RNA was noted; no decrease was noted in lipid content. However, with smaller doses, or administration of marijuana smoke, no consistent or marked changes were noted. The significance of these effects is unknown. Whether additional effects might be observed with more sophisticated and sensitive methods directed to more restricted analytical problems cannot be answered at present.

SUMMARY

There is no persuasive evidence that marijuana causes morphological changes in the brain. Computer tomography studies on users of marijuana reveal no gross changes in brain structure. Electron micrographic studies of monkey brains indicating morphologic changes are methodologically flawed and cannot be used as evidence for an effect of marijuana on brain cell morphology. Clear effects on brain electrical activity in human beings and in animals have been found after drug exposure. These effects have not been demonstrated to persist in human beings after the drug has been discontinued. Studies of EEG from deep brain structures in chronically treated animals have shown changes after the withdrawal of the drug. These limited findings need to be confirmed by further studies. Studies in human beings and animals indicate that, despite the neurophysiologic effects demonstrated in EEG studies, marijuana does not appear to increase epileptic seizure susceptibility. Current evidence has shown marijuana causes some chemical changes in brain: Cannabinoids affect several neurotransmitter systems, especially the cholinergic system. At high doses marijuana also has been shown to affect nucleoprotein synthesis. The significance of these findings for brain function as demonstrated by human behavior or their clinical relevance is unknown.

RECOMMENDATIONS FOR RESEARCH

In view of the widespread use of cannabis, it would be worthwhile to carry out further and more systematic studies of the effects of cannabis on brain structure, chemistry, and electrophysiology. Such studies should be closely correlated with behavior, e.g., learning, psychomotor coordination (see Chapter 6). One useful approach might be to investigate the effects of medium and high doses of cannabis (defined in terms of the patterns of human consumption) on juvenile and adult monkeys during and after long-term exposure. Juvenile monkeys should be included because the immature nervous system may be more sensitive to harmful drug effects; this issue is of great clinical concern, because marijuana use by human beings now begins quite early in life (see Chapter 2). Observations also should be made during long-term abstinence after previous long-term exposure to

determine whether any persistent abnormalities have been produced. A systematic approach to these questions using modern methods of measurement and analysis could extend our present knowledge substantially.

REFERENCES

- Adams, P.M. and Barratt, E.S. Effect of chronic marijuana administration on stages of primate sleepwakefulness. Biol. Psychiatry 10:315-322, 1975.
- Banerjee, S.P., Snyder, S.H., and Mechoulam, R. Cannabinoids: Influence on neurotransmitter uptake in rat brain synaptosomes. J. Pharmacol. Exp. Ther. 194:74-81, 1975.
- Barratt, E.S., Beaver, W., and White, R. The effects of marijuana on human sleep patterns. Biol. Psychiatry 8:47-54, 1974.
- Bracs, P., Jackson, D.M., and Chester, G.B. The effect of Δ -9-tetrahydrocannabinol on brain amine concentration and turnover in whole rat brain and in various regions of the brain. J. Pharm. Pharmacol. 27:713-715, 1975.
- Bull, J. Cerebral atrophy in young cannabis smokers. Lancet 2:1420, 1971.
- Campbell, A.M.G., Evans, M., Thomson, J.L.G., and Williams, M.J. Cerebral atrophy in young cannabis smokers. Lancet 2:1219-1225, 1971.
- Co, B.T., Goodwin, D.N., Gado, M., et al. Absence of cerebral atrophy in chronic cannabis users. JAMA 237:1229-1230, 1977.
- Consroe, P. and Fish, B.S. Behavioral pharmacology of tetrahydrocannabinol convulsions in rabbits. Comm. Psychopharmacol. Comm. Pharmacol. 4:287-291, 1981.
- Domino, E.F., Donelson, A.C., and Tuttle, T. Effects of Δ -9-tetrahydrocannabinol on regional brain acetylcholine, pp. 673-678. In Jenden, D.J. (ed.) Cholinergic Mechanisms and Psychopharmacology. New York: Plenum Press, 1978.
- Domino, E.F. Cannabinoids and the cholinergic system. J. Clin. Pharmacol. 21(Suppl.):249S-255S, 1981.
- Drew, W.G. and Miller, L.L. Cannabis: Neural mechanisms and behavior--A theoretical review. Pharmacology 11:12-32, 1974.
- El-Yousef, M.K., Janowsky, D.S., Davis, J.M., and Rosenblatt, J.E. Induction of severe depression by physostigmine in marijuana intoxicated individuals. Br. J. Addict. 68:321-325, 1973.
- Fehr, K.A., Kalant, H., Leblanc, A.E., and Knox, G.V. Permanent learning impairment after chronic heavy exposure to cannabis or ethanol in the rat, pp. 495-505. In Nahas, G.G. (ed.) Marihuana: Chemistry, Biochemistry, and Cellular Effects. New York: Springer-Verlag, 1976.
- Feeney, D.M., Wagner, H.R., McNamara, M.C., and Weiss, G. Effects of tetrahydrocannabinol on hippocampal evoked afterdischarges in cats. Exp. Neurol. 41:357-365, 1973.
- Feeney, D.M. Marihuana and epilepsy: Paradoxical anticonvulsant and convulsant effects, pp. 643-657. In Nahas, G.G. and Paton,

- W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Feinberg, I., Jones, R., Walker, J.M., et al. Effects of high dosage delta-9-tetrahydrocannabinol on sleep patterns in man. Clin. Pharmacol. Ther. 17:458-466, 1975.
- Feinberg, I., Jones, R., Walker, J.M., et al. Effects of marijuana tetrahydrocannabinol on electroencephalographic sleep patterns. Clin. Pharmacol. Ther. 19:782-794, 1976.
- Fink, M. Effects of acute and chronic inhalation of hashish, marijuana, and delta-9-tetrahydrocannabinol on brain electrical activity in man: Evidence for tissue tolerance. Ann. N.Y. Acad. Sci. 282:387-398, 1976.
- Freemon, F.R. The effect of delta-9-tetrahydrocannabinol on sleep. Psychopharmacologia 35:39-44, 1974.
- Freemon, F.R., Rosenblatt, J.E., and El-Yousef, M.K. Interaction of physostigmine and delta-9-tetrahydrocannabinol in man. Clin. Pharmacol. Ther. 17:121-126, 1975.
- Fried, P.A. Behavioral and electroencephalographic correlates of the chronic use of marijuana--A review. Behav. Biol. 21:163-196, 1977.
- Fujimori, M. and Himwich, H.E. Δ -9-Tetrahydrocannabinol and the sleep-wakefulness cycle in rabbits. Physiol. Behav. 11:291-295, 1973.
- Harper, J.W., Heath, R.G., and Myers, W.A. Effects of cannabis sativa on ultrastructure of the synapse in monkey brain. J. Neurosci. Res. 3:87-93, 1977.
- Heath, R.G. Marihuana and Δ -9-tetrahydrocannabinol: Acute and chronic effects on brain function of monkeys, pp. 345-356. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuana. New York: Raven Press, 1976.
- Heath, R.G., Fitzjarrell, A.T., Garey, R.E., and Myers, W.A. Chronic marihuana smoking: Its effect on function and structure of the primate brain, pp. 713-730. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Heath, R.G., Fitzjarrell, A.T., Fontana, C.J., and Garey, R.E. Cannabis sativa: Effects on brain function and ultrastructure in Rhesus monkeys. Biol. Psychiatry 15:657-690, 1980.
- Herning, R.I., Jones, R.T., and Peltzman, D.J. Changes in human event related potentials with prolonged delta-9-tetrahydrocannabinol (THC) use. Electroencephalogr. Clin. Neurophysiol. 47:556-570, 1979.
- Hosko, M.J., Kochar, M.S., and Wang, R.I.H. Effects of orally administered delta-9-tetrahydrocannabinol in man. Clin. Pharmacol. Ther. 14:344-352, 1973.
- Jones, R.T. and Stone, G.C. Psychological studies of marijuana and alcohol in man. Psychopharmacologia 18:108-117, 1970.
- Karacan, I., Fernandez-Salas, A., Coggins, W.J., et al. Sleep electroencephalographic-electrooculographic characteristics of

- chronic marijuana users: Part I. Ann. N.Y. Acad. Sci. 282:348-374, 1976.
- Karler, R. and Turkanis, S.A. The cannabinoids as potential antiepileptics. J. Clin. Pharmacol. 21(Suppl.):4375-4485, 1981.
- Kuehne, J., Mendelson, J.H., Davis, K.R., and New, P.F.J. Computed tomographic examination of heavy marijuana smokers. JAMA 237:1231-1232, 1977.
- Lemberger, L. Potential therapeutic usefulness of marijuana. Ann. Rev. Pharmacol. Toxicol. 20:151-172, 1980.
- Low, M.D., Klonoff, H., and Marcus, A. The neurophysiological basis of the marijuana experience. Can. Med. Assoc. J. 108:157-164, 1973.
- Luthra, Y.K. and Rosenkrantz, H. Cannabinoids: Neurochemical aspects after oral chronic administration to rats. Toxicol. Appl. Pharmacol. 27:158-168, 1974.
- Luthra, Y.K., Rosenkrantz, H., Heyman, I.A., and Braude, M.C. Differential neurochemistry and temporal pattern in rats treated orally with Δ -9-tetrahydrocannabinol for periods up to six months. Toxicol. Appl. Pharmacol. 32:418-431, 1975.
- Luthra, Y.K., Rosenkrantz, H., and Braude, M.C. Cerebral and cerebellar neurochemical changes and behavioral manifestations in rats chronically exposed to marijuana smoke. Toxicol. Appl. Pharmacol. 35:455-465, 1976.
- Martinez, J.L., Stadnicki, S.W., and Schaeppi, U.N. Delta-9-tetrahydrocannabinol: Effects on EEG and behavior of Rhesus monkeys. Life Sci. 11:643-651, 1972.
- Meldrum, B.S., Fariello, R.G., Puil, E.A., et al. Delta-9-tetrahydrocannabinol and epilepsy in the photosensitive baboon, Papio papio. Epilepsia 15:255-264, 1974.
- Moreton, J.E. and Davis, W.M. Electroencephalographic study of the effects of tetrahydrocannabinols on sleep in the rat. Neuropharmacology 12:897-907, 1973.
- Myers, W.A. and Heath, R.G. Cannabis sativa: Ultrastructural changes in organelles of neurons in brain septal region of monkeys. J. Neurosci. Res. 4:9-17, 1979.
- Pivik, R.T., Zarcone, V., Dement, W.C., and Hollister, L.E. Delta-9-tetrahydrocannabinol and synhexl: Effects on human sleep patterns. Clin. Pharmacol. Ther. 13:426-435, 1972.
- Prankoff, K., Karacan, I., Larson, E.A., et al. Effects of marijuana smoking on the sleep EEG: Preliminary studies. J. Fla. Med. Assoc. 60:28-31, 1973.
- Revuelta, A.V., Moroni, F., Cheney, D.L., and Costa, E. Effect of cannabinoids on the turnover rate of acetylcholine in rat hippocampus, striatum, and cortex. Arch. Pharmacol. 304:107-110, 1978.
- Rodin, E.A., Domino, E.F., and Porzak, J.P. The marijuana-induced "social high." Neurological and electroencephalographic concomitants. JAMA 213:1300-1302, 1970.
- Rubin, V. and Comitas, L. Ganja in Jamaica: A Medical Anthropological Study of Chronic Marijuana Use. The Hague: Mouton and Co., 1975.

- Rumbaugh, C.L., Fang, H.C.H., Wilson, G.H., et al. Cerebral CT findings in drug abuse: Clinical and experimental observations. J. Comput. Assisted Tomography 4:330-334, 1980.
- Sipe, J.C. and Moore, R.Y. The lateral hypothalamic area: An ultrastructural analysis. Cell Tiss. Res. 179:177-196, 1977.
- Stadnicki, S.W., Schaeppi, U., Rosenkrantz, H., and Braude, M.C. Crude marijuana extract: EEG and behavioral effects of chronic oral administration in rhesus monkeys. Psychopharmacologia 37:225-233, 1974.
- Volavka, J., Dornbush, R., Feldstein, S., et al. Marijuana, EEG and behavior. Ann. N.Y. Acad. Sci. 191:206-215, 1971.
- Wikler, A. and Lloyd, B. Effect of smoking marijuana cigarettes on cortical electrical activity. Fed. Proc. 4:141, 1945.

5

EFFECTS OF MARIJUANA ON OTHER BIOLOGICAL SYSTEMS

This chapter covers what little is known about the effects of cannabis on male and female reproduction and endocrine systems, birth defects and teratogenic effects, genetics, the immune system, and body temperature.

MALE REPRODUCTIVE FUNCTION

A variety of studies indicate that marijuana and some of its derivatives have reversible, suppressive effects upon testicular function in animals and men. These have been measured in terms of diminished weights of the prostate gland, seminal vesicles, or testes, and in decreased levels of testosterone (the male hormone) in blood plasma or suppression of spermatogenesis following chronic or acute administration of cannabis or Δ -9-TcC. Appropriate observations have indicated that the effects of cannabinoids on the male reproductive tract and on testicular function were completely reversed 1 month after drug withdrawal.

There is no general agreement as to the cause or magnitude of these effects. The major reasons for this lack of agreement relate to major differences in study design, including species studied (man, monkey, or rodent), route of drug administration, and purity of the drug used.

Human Studies

In 1974, a group of 20 men were studied who had used marijuana at least 4 days a week for a minimum of 6 months without the use of other drugs (Kolodny et al., 1974). Plasma testosterone levels in subjects smoking five to nine marijuana cigarettes per week were significantly lower than controls (however, only 2 had levels out of the normal range, i.e., below 400 ng/dl); all but 1 of the men smoking more than 10 marijuana cigarettes per week had testosterone levels below 400 ng/dl. These results suggest that there was a dose-dependent effect of marijuana on testosterone levels. Plasma levels of lutenizing hormone (LH) and follicle-stimulating hormone

(FSH), gonadotropins that control the growth of the ovaries or testes and their hormonal activities, were in the normal range; however, in men smoking more than 10 marijuana cigarettes per week, the FSH level was significantly lower than for those who smoked 5 to 10 marijuana cigarettes per week. Because only random samples of blood were obtained for gonadotropic measurements, small but significant changes could have been missed. Levels of prolactin, the female hormone involved in lactation and also present in small quantities in men, were all in the low normal range. In addition, the men who smoked more than 10 marijuana cigarettes per week had significantly lower sperm counts than those who smoked the lesser quantity (26 versus 68 million/ml). These individuals obtained marijuana from a variety of sources, and there was no way to determine whether they were taking other drugs that could lower plasma testosterone.

Later in 1974, another study reported that plasma testosterone levels were not suppressed in 27 men studied in a research ward (Mendelson et al., 1974). These individuals smoked marijuana cigarettes supplied by the federal government. For unexplained reasons, the mean testosterone levels in these individuals were greater than 1,000 ng/dl (higher than the normal mean) before and during the smoking periods. This is in marked contrast to the mean value of 742 ng/dl for nonsmokers in the study of Kolodny et al. mentioned above. There was no report of gonadotropic values or semen analysis in the Mendelson Study.

A study of 16 patients on a metabolic ward who smoked NIDA cigarettes (Hembree et al., 1979) showed that 5 to 6 weeks of high-dose (2 percent) marijuana administration (8-20 cigarettes/day) was associated with a decline in sperm count during the fifth and sixth weeks after initiation of drug exposure. This was preceded by a decrease in sperm motility and an increase in abnormal forms of sperm. Once a week during the study five blood samples were obtained at 15-minute intervals for measurement of testosterone, LH, and FSH. No change in these hormone levels was noted throughout the study (although no values were reported). The relationship in time of these samples to the last previous cigarette was not mentioned, therefore the test would not have excluded a transient decline in hormonal levels after each cigarette. However, because hormonal suppression of spermatogenesis takes longer than 4 weeks and usually is not associated with an increase in the number of abnormal forms and a decrease in motility, the authors concluded that the effect upon the seminiferous tubular epithelium was direct rather than by suppression of gonadotropins. This is the only reported study in man that measured the hour-to-hour fluctuations in gonadotropic levels.

Another study (Coggins et al., 1976) evaluated the health status of 84 marijuana smokers who had used the agent three or more times per week for a minimum of 10 years. Testosterone levels were measured in 38 users and 35 nonusers. The mean levels and ranges were virtually identical. This heterogeneous group of men patients studied in Costa Rica was not recruited for the purpose of studying the pituitary-gonadal axis. No gonadotropic levels or semen samples were studied.

Endocrine function studies are briefly mentioned in a paper by Cohen (1976). Subjects were recruited on the basis of heavy marijuana use and were studied in a metabolic ward. They smoked an average of five marijuana cigarettes per day, which was believed to be the equivalent of 103 mg of Δ -9-THC. During acute administration, mean levels of plasma testosterone declined from 754 to 533 ng/dl over a 3-hour period. After 9 weeks of smoking, plasma testosterone levels had declined from 740 to 509 ng. Plasma LH levels were reported to have fallen after the fourth week; however, no absolute values were given. In addition, no standard errors are given for any of the means presented in this paper. Therefore, it is impossible to evaluate the significance of the reported findings.

In Greece, a population of 47 chronic hashish users was studied. Electron microscope studies of the acrosome, the head of the sperm, showed abnormality in some patients (Issidorides, 1979). It is difficult to evaluate the study because no quantitative data were presented.

Animal Studies

All of the studies mentioned below are substantially different from those of human beings because, with one exception, the active agent (usually Δ -9-THC) was administered intraperitoneally at a dose of 2.5 to 25 mg/kg. Based on calculations given by Cohen (1976), 3 to 6 mg/kg/day would be considered a large dose in human beings. Also, human beings self-administer the drug over many hours rather than as a single dose.

In castrated rhesus monkeys, plasma LH and FSH fell acutely following acute administration of Δ -9-THC (Smith et al., 1980). During this suppression period, both gonadotropins could be stimulated by lutenizing hormone-releasing factor (LHRF), which causes the release of LH. The effect of Δ -9-THC was to suppress prolactin release, which, in turn, could be stimulated by thyrotropin-releasing hormone (TRH). Studies in other species have tended to confirm these observations in monkeys.

The results are compatible with the hypothesis that the effect of marijuana and its derivatives is on gonadotropic secretion (Farclerode et al., 1979). Testicular cytochrome P-450 (an enzyme) decreased in the rat following 2 to 9 weeks of treatment. The concentrations of this enzyme, plus a variety of other testicular markers, were restored with FSH and LH therapy. The effect of various cannabinoids has been studied on sperm morphology in the mouse (Zimmerman et al., 1979).

*Rosenkrantz (1981) considers 0.6-3.0 mg/kg by inhalation and 1.8-9.0 mg/kg orally to be large doses in human beings. For the monkey, 1.8-9.0 mg/kg by inhalation and 5-27 mg/kg orally would be considered a high dose. (These concentrations are equivalent to six cigarettes/day.)

Mice were given five daily intraperitoneal injections of Δ -9-THC, cannabidiol, or cannabinal at doses approaching or exceeding the LD₅₀ (the dose necessary to kill 50 percent of the animals). Thirty-five days after the last treatment, animals were killed and sperm were evaluated by scanning electron microscopy. Control animals had 1.5 percent abnormal forms. Animals that received LD₅₀ doses of the various derivatives had 2.4 to 5.0 percent abnormal forms.

Only a few studies have examined the effects of cannabis on spermatogenesis (Huang et al., 1979). Marijuana was administered to rats in a smoke machine. After 30 days of exposure, marijuana smoke lowered the sperm counts in animals significantly, as did cannabinoid-free smoke. By 75 days, however, only the marijuana smoke group maintained a low sperm count. In the marijuana-treated group, there was an increased number of abnormal forms, particularly with an increase in dissociation of sperm heads and tails. In the discussion of this paper, the authors reported elevated serum FSH levels following marijuana exposure, but did not present data. They concluded that marijuana has a direct effect on the testis. A variety of *in vitro* studies support this suggestion (Jakubovic et al., 1977, 1979).

Marijuana and its derivatives also have been shown to be antiandrogenic (antagonistic to male hormones) (Purohit et al., 1980). Several constituents, including Δ -9-THC, can bind to the receptor for androgen. Marijuana also has been demonstrated to be estrogenic (like female sex hormones) *in vivo*, and recent studies suggest that these effects may be mediated via the estrogen receptor. These observations have been disputed by others (reviewed by Purohit et al., 1980). The ability to inhibit or mimic the action of sex steroids provides one mechanism by which these agents can produce their effects. There obviously are many others.

FEMALE REPRODUCTIVE FUNCTION

The effect of cannabis on female reproduction has been studied in rats, mice, rabbits, and monkeys. The work in rhesus monkeys is of particular importance, because of the similarity in the menstrual cycle among primate species, including human beings.

Human Studies

There is only one study reported on the effects of marijuana on reproductive function in women. The work has appeared in print as a report of the proceedings of a 1978 symposium held in Mexico City (Bauman et al., 1979) and as part of the congressional record subsequent to testimony before a Senate committee hearing (Bauman, 1980). These publications do not provide details on methodology or on individual hormone values. Differences between the control and experimental groups, recognized by the investigators, could be of

importance; alcohol use, for example, was more frequent in the marijuana-using group. The study attempted to establish the endocrine (hormonal) profile and menstrual patterns of women who used marijuana on a chronic and frequent basis. Twenty-six women who used it at least three times a week for 6 months were compared with 17 women who had never used the substance. The number of cycles studied for each variable investigated is not clear from the publications. This difficulty notwithstanding, the report reveals no difference in plasma levels of LH and FSH between the two groups and no change in peaks and basal values of the female hormones estradiol or progesterone, the critical hormone levels controlling the process of ovulation. It would be expected that no major difference was found in the incidence of anovulatory cycles between the two groups. By combining anovulation and shortened luteal phase, however, the authors report a statistically significant difference in the marijuana-using group, which could be clinically important in causing subfertility. This evidence is, at best, only suggestive. The observation that testosterone levels in marijuana-using women are elevated is difficult to interpret in terms of clinical significance; apparently, the subjects did not report episodes of acne, abnormal hairiness, or other testosterone-dependent side-effects. According to the authors, serum prolactin levels are lower in marijuana users than in controls. The implications of this observation for fertility, lactation, or the development of breast cancer are not clear.

The absence of other studies on users of marijuana makes it difficult to draw conclusions on the implications of the data cited above. Several of the effects noted are different from the more extensive and experimentally controlled observations in rhesus monkeys and other laboratory animals. This situation calls attention to the urgent need for more comprehensive endocrine and gynecologic investigations of women who use marijuana.

Animal Studies

Administration of crude marijuana extract to rats or mice resulted generally in suppression of ovarian function and in various aspects of estrogen activity, such as uterine metabolism, weight, glycogen content, and levels of RNA and sialic acid (Chakravarty et al., 1975; Dixit et al., 1975).

The administration of crude marijuana extract for 30 days to rats and mice abolished the estrus cycle and caused a significant reduction in the size of the ovaries and in some primordial ova (Dixit et al., 1975). Intraperitoneal administration of Δ -9-THC to rats, appropriately timed, has also been reported to block ovulation (Nir et al., 1973). This effect of Δ -9-THC was exerted by suppressing the characteristic preovulatory surge of plasma LH. Other investigators have reported suppression also of plasma FSH and prolactin when Δ -9-THC is given just before ovulation (Ayalon et al., 1977). The substance was found to depress plasma concentration of LH in ovariectomized rats (Marks, 1973; Tyrey, 1978, 1980) and

rhesus monkeys (Besch et al., 1977). Asch et al. (1979) also showed in the rabbit, a reflex ovulator, that a precoital single-dose of Δ -9-THC blocks the postcoital LH surge and ovulation.

Administration of LHRF was able to bring about the release of LH in Δ -9-THC treated rats and rhesus monkeys (Smith et al., 1979). These results indicate a direct effect of cannabinoids at the level of the hypothalamus, part of brain important in reproductive hormone regulation. The ovulation-blocking effect of the cannabinoids was further investigated by Cordova et al. (1980). Natural and chemically modified cannabinoids blocked ovulation in rats.

Administration of Δ -9-THC to rhesus monkeys during the follicular phase resulted in prolonged periods of amenorrhea (absence or abnormal stoppage of the menstrual flow), absence of midcycle LH surge, and progesterone levels characteristic of anovulation (Asch et al., 1981).

BIRTH DEFECTS AND TERATOGENICITY

Because Δ -9-THC crosses the placenta it is a potential teratogen, an agent that causes defects in the developing embryo. This effect could occur in either of two ways: (1) exposure to cannabis prior to conception could harm the sex cells (the ova and sperm), or (2) the fetus could be harmed directly during organogenesis. In addition, Δ -9-THC can be secreted in breast milk and, therefore, can be toxic postnatally.

Human Studies

The evidence for teratogenicity in human beings is very difficult to interpret. Although there is widespread use of marijuana in young women of reproductive age, there is no evidence yet of any teratogenic effects of high frequency or consistent association with the drug. There are isolated reports of congenital anomalies in the offspring of marijuana users, but there is no evidence that they occurred more often in users than in nonusers and in those cases there was coincident use of other drugs. Subtle development effects in offspring, such as nervous system abnormalities, and reductions in birth weight and height may indeed exist (Finnegan, 1980; Fried, 1980; Hingson et al., in press). Additional carefully designed, prospective studies should provide valuable information in this area.

Animal Studies

Crude marijuana extract and Δ -9-THC are teratogenic at certain doses in animals.*

*Bibliography available upon request from the Institute of Medicine, National Academy of Sciences.

One study reported that subcutaneous injection of pregnant hamsters and rabbits with various doses of crude marijuana extract caused malformations of the brain, spinal cord, forelimb, and liver, as well as edema of the head and spinal region in developing embryos (Gerber and Schramm, 1969). In hamsters, significant embryocidal and growth retardation effects also were noted. It was concluded that doses greater than 200 mg/kg in hamsters and 250 mg/kg in rabbits were teratogenic. Caution in interpreting these findings must be exercised because the teratogenic effects may be caused by any combination of constituents of the cannabis extract.

In a study of mice, the teratological effects of Δ -9-THC were evaluated for doses ranging from 3.0 to 400 mg/kg by various routes of administration--intravenous, subcutaneous, and intragastric (Jones, 1976). Significant fetal growth retardation was induced at higher dose levels and by some routes of administration. For example, a high dose of 400 mg/kg was significantly teratogenic by the intragastric route; 12.1 percent of the live fetuses were malformed.

In a study of female monkeys given an oral dose of 2.4 mg/kg Δ -9-THC for 1 to 4 years, a nonspecific pattern of reproductive difficulties was observed characteristic of "high-risk" pregnancies, including a high rate of offspring loss during pregnancy or in the early postnatal period (Sassenrath et al., 1979).

GENETIC EFFECTS

The potential genetic effects of marijuana are of major concern because of its prevalent use by young people in their reproductive years (see Chapter 2). Although there is a growing amount of evidence that drugs can induce mutations, and an improving ability to use toxicological methods to evaluate agents for their mutagenic potential (such as the Ames test, which detects changes or damages in the genetic material), the available information on the genetic hazards or even on the potential genetic hazards of the use of marijuana is extremely limited.

Mutagenicity

Elsewhere in this report (Chapter 3) the scientific evidence that marijuana smoke and tar are mutagenic has been discussed. Lung explants of mice and human fibroblast cultures exposed to fresh smoke showed abnormalities of cell division, as well as changes in chromosome structure and in DNA synthesis (Leuchtenberger and Leuchtenberger, 1971; Leuchtenberger, et al., 1973a,b). Moreover, extracts and smoke condensates of marijuana are mutagenic when evaluated by the Ames test (Busch et al., 1979; Seid and Wei, 1979; Wehner et al., 1980). Animal studies on rodents painted with marijuana tar, three times weekly for 1 year, resulted in skin papillomas, carcinomas, and fibrosarcomas (Hoffmann et al., 1975).

However, extensive testing with Δ -9-THC using three established tests for mutagenesis failed to detect any mutagenic effect, or any effect as an inhibitor of DNA repair (Legator et al., 1976; Glatt et al., 1979; Zimmerman et al., 1978).

Cytogenetic Effects

The numbers and kinds of chromosomes (structures in a cell nucleus that contain and transmit genetic information carried in DNA) are highly characteristic for a given species. Structural variation and changes in numbers of chromosomes may be evidence for genetic damage produced by drugs and other chemical agents. Unfortunately, the literature on the effects of marijuana on chromosomes is limited and conflicting. Studies suggesting that marijuana probably does not break chromosomes are fairly conclusive. There is less evidence that marijuana may produce aneuploidy (abnormal numbers of chromosomes) in some daughter cells during cell division.

Does marijuana cause chromosome breaks? The weight of the evidence from in vitro cultures of human cells and from in vivo animal and human studies is that neither marijuana nor Δ -9-THC causes chromosome breaks.

In Vitro and Animal Studies

Cultures of human leukocytes, exposed to different concentrations of Δ -9-THC, showed no increase in the incidence of chromosome breaks or gaps when compared to controls (Stenchever and Allen, 1972). Studies of golden hamsters given subcutaneous injections for 10 days of marijuana extract distillate containing 17.1 percent Δ -9-THC (Nicholson et al., 1973), and of beagle dogs trained to smoke high doses of marijuana (3 g/day/week for 30 months), showed no significant differences in chromosome gaps or breaks when compared with control groups (Genest et al., 1976).

Human Studies

Cytogenetic analysis of chromosomes from peripheral blood leukocytes and cultures of subjects exposed to marijuana smoking, marijuana extract, or synthetic Δ -9-THC revealed no increase in chromosome breakage attributable to these compounds (Nichols et al., 1974; Matsuura, 1976; Morishima et al., 1979). Doses ranged from 20 mg Δ -9-THC per day to 12-16 marijuana cigarettes per day. Studies that have reported chromosome breaks or gaps in cell cultures of users of marijuana have largely been carried out on multiple drug users, and the breaks and gaps may be due to other factors associated with a life of heavy drug use (Gilmour et al., 1971; Horha and Obe, 1974). However, in a retrospective study on college students, chromosome breaks were found in blood cultures of 49 light (one or

less exposure per week) and heavy (more than two exposures per week) users of marijuana (Stenchever et al., 1974). One problem in this study is the poor dose characterization. Furthermore, the increase in the numbers of breaks in both light and heavy users of marijuana was not dose-related; the same frequency of breaks was observed in both groups. Although the evidence is inconclusive, it suggests that marijuana does not cause chromosome breaks.

Does marijuana interfere with cell division and chromosome segregation, thereby resulting in abnormal numbers of chromosomes? There is conflicting evidence in the literature. On the one hand, no significant effects of marijuana smoke or Δ -9-THC on chromosome complement have been reported using the micronuclei test in mice or in cytogenetic studies in dogs (Genest et al., 1976; Legator et al., 1976). On the other hand, more extensive studies have demonstrated aneuploidy resulting from in vitro exposure of cells to marijuana as well as in vivo studies of animals and human beings.

In Vitro and Animal Studies

Exposure of mouse lung and adult human lung tissue culture to marijuana smoke in vitro resulted in abnormal cell proliferation and abnormalities in DNA content (Leuchtenberger and Leuchtenberger, 1971; Leuchtenberger, et al., 1973b). Addition of Δ -9-THC and olivetol, a compound with a ring structure similar to cannabinoids, to normal human leukocyte cultures induced hypodiploidy (defined as metaphase nuclei with a chromosome complement of less than 30 chromosomes—a normal human cell contains 46 chromosomes) (Morishima et al., 1976). Hybrid mice treated for 5 consecutive days with Δ -9-THC, cannabinal, and cannabidiol at a dose of 10 mg/kg had a three- to fivefold increase of micronuclei over controls. The number of micronuclei increased with increasing Δ -9-THC dosage. Examination of bone marrow mitosis in these same mice showed a five- to sevenfold increase in chromosome number aberrations during metaphase (Zimmerman and Raj, 1980).

Human Studies

Studies of lymphocytes cultured from human marijuana smokers defined either as "moderate" users (at least one marijuana cigarette per week, range 1-10 for a minimum of two years) or "heavy" users (more than three times per week) all of whom consumed between 12.9 and 15.3 marijuana cigarettes per day during the experiment, turned up a significantly larger number of cells with less than 30 chromosomes than would be found in normal control cultures (Morishima et al., 1979). These positive findings suggest that marijuana may affect chromosome segregation during cell division and result in cells with fewer than the normal number of chromosomes. What these findings mean in terms of risk for abnormalities in offspring or possible disease is not known. Findings in lymphocyte cultures may not be relevant to what is happening in the germ cells (sex cells).

THE IMMUNE SYSTEM

The immune system functions in protecting the body against viruses, bacteria, and other infections. It also plays a major role in preventing the growth and dissemination of cancerous cells.

There have been reports that cannabis is immunogenic, capable of activating components in the immune system. These components include such cells as lymphocytes, some of which produce antibodies in response to invasion by a foreign agent, and macrophages, which can be stimulated by inflammation to ingest invaders.

Human Studies

There have been reports that cannabis interferes with components in the immune system in man. Antibodies will develop in response to marijuana in some people, along with an allergic response, while others develop antibodies without apparent allergic reaction (Liskow et al., 1971; Shapiro et al., 1974, 1976; Lewis and Slavin, 1975). However, the studies reporting these effects were not designed to determine which components of the marijuana are immunogenic and which are allergenic.

Studies of various aspects of the immune system in persons who were chronic users of marijuana have indicated mild decreases in activity of one or another component of the system; however, other investigators have noted no changes outside of the normal range (Gupta et al., 1974; Petersen et al., 1975, 1976; White et al., 1975; Lau et al., 1976; Rachelefsky et al., 1976; Silverstein and Lessin, 1976; Cushman and Khurana, 1977; McDonough et al., 1980). These apparent inconsistencies may stem from the variability in the amount of marijuana consumed among users in different studies and the differences in the immune system assays. Hashish, as distinct from marijuana, was shown to have a slight temporary stimulatory effect on the immune system (Kaklamani et al., 1978; Kalofoutis et al., 1978).

Animal Studies

A number of studies have shown that Δ -9-THC and other cannabinoids induce immunological defects in rodents (Petersen and Lemberger, 1976; Lefkowitz and Klager, 1978; Lefkowitz et al., 1978; Preuss and Lefkowitz, 1978). The doses varied from 5 to 25 mg/kg (intra-peritoneally) to 100 mg/kg (orally). At the higher doses there was a diminution of immune response, as measured by standard immunological assays. Delta-9-THC had the same effects on cells grown in vitro. Other cannabinoids also have been tested for their effects. Cannabinol, Δ -8-THC, and 1-methyl- Δ -8-THC had the same immunosuppressive effects as Δ -9-THC, but cannabidiol had no immunosuppressive effect. Immunizing rabbits with Δ -9-THC resulted in the production of antibodies (Chiarotti et al., 1980).

BODY TEMPERATURE

Regulation of body temperature is a complex process that can be influenced by drugs. In several species of animals, Δ -9-THC produces a lowering of body temperature (hypothermia). The effect is seen when animals are housed at normal room temperatures, and it is greater with colder ambient temperatures (Pertwee and Travendale, 1979). Marijuana apparently causes a decrease in heat production for reasons that are unclear.

In experiments with human subjects, marijuana has produced little or no change in body temperature when given in a cool environment (Beaconsfield et al., 1972; Hanna et al., 1976). In a hot environment (40°C) marijuana caused inhibition of sweating and a consistent rise in body temperature (Jones et al., 1980). Thus, there is evidence that marijuana does interfere with temperature regulation, although there is no currently known clinical significance to this finding.

Cannabis appears to interfere with temperature regulation, but the clinical significance is unknown.

SUMMARY

Male Reproductive Function

In animals, marijuana and its derivatives can acutely lower gonadotropic secretion when administered intraperitoneally. There is also some evidence in animals to suggest that these agents can directly affect the seminiferous tubule. In man, sperm number and motility are decreased during chronic marijuana use. From the available studies, it appears this was due to a direct effect of the cannabinoids either on the seminiferous tubular epithelium or the epididymal sperm. Due to conflicting and incomplete evidence, it is not possible to conclude at the present time whether marijuana smoking has a significant effect upon gonadotropic and testosterone concentrations in humans. Whether the decrease in sperm number or motility has any effect on fertility is not known.

Female Reproductive Function

There is only one study of human beings that attempts to establish the endocrine profile and menstrual patterns of women who used marijuana on a chronic and frequent basis. By combining categories of anovulation and shortened luteal phase, a statistically significant difference was noticed in the marijuana using group. It is not known if this leads to problems with fertility or lactation, or if it leads to cancer of the reproductive organs.

Animal studies have shown that Δ -9-THC lowers the serum gonadotropic levels. It is unknown if there is a direct effect on the reproductive tissues, particularly under prolonged use of cannabis products.

Birth Defects and Teratogenicity

Cannabis is teratogenic at high doses in animals. There is no evidence of obvious teratogenicity or structural defects in the offspring of human users. But the data are not adequate to reveal a long-range functional impairment or a very low level of teratogenicity if one is present. It may be impossible to identify a distinct role for cannabis in the production of subtle effects in offspring, because of the confounding influences of malnutrition, smoking, and alcohol.

Genetic Effects

Marijuana and Δ -9-THC do not appear to break chromosomes, although there is some conflicting evidence on this point. Multiple drug use seems to be correlated with an increase in the numbers of gaps and breaks in the genetic material. Furthermore, marijuana may affect chromosome segregation during cell division, resulting in abnormal numbers of chromosomes in daughter cells. While these conflicting results are worrisome, their clinical significance is not known. Further investigations, especially controlled prospective studies, of human beings are needed.

The Immune System

The data from animal studies suggest that Δ -9-THC and some of its analogues have a mild, transient, immunosuppressant effect in both in vitro and in vivo systems; the effects are mild compared with known immunosuppressant drugs. The studies in human beings are contradictory; some demonstrated mild, immunosuppressive effects, but others, using the same or similar methods, did not find any differences in the immune system between normals and chronic marijuana smokers. At the present time, there have been no human or animal studies that have determined if marijuana smokers are more prone to infections or other diseases. Because of the widespread use of marijuana, even weak immunosuppressive effects are a concern. Since further research may not demonstrate definitive findings, immunologic effects should be studied along with other variables in a larger investigation. If marijuana is to be used on immunosuppressed patients (for example, for antiemetic purposes during cancer chemotherapy), even minor additional suppression might be dangerous.

RECOMMENDATIONS FOR RESEARCH

The committee recommends the following types of studies.

- * Further observations should be made regarding the relation of marijuana use to reproductive defects in human beings, especially

on young users whose reproductive biology is undergoing rapid change. The principal need is for assessment of endocrine profiles and semen analysis in male users versus nonusers, with adequate control of confounding variables--for example, diet, alcohol, other drug use. In women, the principal need is for more data on endocrine and menstrual patterns in users versus nonusers, with particular attention to the length of cycles, the presence or absence of ovulation, and the existence or absence of subfertility. More studies are needed to detect subtle, low-frequency, or cumulative effects on reproductive function in long-term, heavy users.

• Although routine testing of teratogenicity in human beings is not recommended at this time, the collection of precise epidemiologic information on the outcome of human pregnancy in marijuana users is of great importance and must be carefully controlled.

• There are no good animal models for studying the effects of smoking marijuana, but cytogenetic studies in animals after exposure to Δ -9-THC by other routes than smoking would be of some value. The most relevant studies still would be in vivo human studies.

• Marijuana has been found to have mild immunological effects in a variety of test systems, but studies of its influence on the body's immune defense against microorganisms are lacking and need to be conducted.

• Critical experiments are needed to test the hypothesis that Δ -9-THC causes disruption of thermoregulatory effector responses rather than an alteration of the level of thermoregulation.

• Inherited variation in the way some drugs are metabolized is widely recognized. This type of variation must be evaluated in respect to susceptibility to marijuana.

REFERENCES

- Asch, R.H., Fernandez, E.O., Smith, C.G., and Pauerstein, C.J. Precoital single doses of Δ -9-tetrahydrocannabinol block ovulation in the rabbit. Fertil. Steril. 31:331-334, 1979.
- Asch, R.H., Smith, C.G., Siler-Khodr, T.M., and Pauerstein, C.J. Effects of Δ -9-tetrahydrocannabinol during the follicular phase of the Rhesus monkey (Macaca mulatta). J. Clin. Endocrin. Metab. 52:50-55, 1981.
- Ayalon, D., Nir, I., Cordova, T., Bauminger, S., et al. Acute effect of Δ -1-tetrahydrocannabinol on the hypothalamo-pituitary-ovarian axis in the rat. Neuroendocrinology 23:31-42, 1977.
- Bauman, J. Marijuana and the female reproductive system, pp. 85-88. Testimony before the Subcommittee on Criminal Justice of the Committee on the Judiciary, U.S. Senate, "Health Consequences of Marijuana Use," January 16-17, 1980. Washington, DC: U.S. Government Printing Office, 1980.
- Bauman, J.E., Kolodny, R.C., Dornbusch, R.L., and Webster, S.K. Efectos endocrinos del uso cronico de la mariguana en mujeres,

- pp. 85-97. In Simposio Internacional Sobre Actualizacion en Marihuana, Volume 10. Tlalpan, Mexico, July 1979.
- Beaconsfield, P., Ginsburg, J., and Rainsbury, R. Marihuana smoking: Cardiovascular effects in man and possible mechanisms. N. Engl. J. Med. 287:209-212, 1972.
- Besch, N.F., Smith, C.G., Besch, P.K., and Kaufman, R.H. The effect of marihuana (delta-9-tetrahydrocannabinol) on the secretion of lutenizing hormone in ovariectomized Rhesus monkey. Am. J. Obstet. Gynecol. 128:635-642, 1977.
- Busch, F.W., Seid, D.A., and Wei, E.T. Mutagenic activity of marihuana smoke condensates. Cancer Lett. 6:319-324, 1979.
- Chakravarty, I., Sengupta, D., Bhattacharyya, P., and Ghosh, J.J. Effect of treatment with cannabis extract on the water and glycogen contents of the uterus in normal and estradiol-treated prepubertal rats. Toxicol. Appl. Pharmacol. 34:513-516, 1975.
- Chiarotti, M., Giusti, G.V., and Vigevani, F. In vivo and in vitro properties of anti-delta-9-tetrahydrocannabinol antibody. Drug Alcohol Dependence 5:231-233, 1980.
- Coggins, W.J., Swenson, E.W., Dawson, W.W., et al. Health status of chronic heavy cannabis users. Ann. N.Y. Acad. Sci. 282:148-161, 1976.
- Cohen, S. The 94-day cannabis study. Ann. N.Y. Acad. Sci. 282:211-220, 1976.
- Cordova, T., Ayalon, D., Lander, N., et al. The ovulation blocking effect of cannabinoids: Structure-activity relationships. Psychoneuroendocrinology 5:53-62, 1980.
- Cottrell, J.C., Sohn, S.S., and Vogel, W.H. Toxic effects of marihuana tar on mouse skin. Arch. Environ. Health 26:277-278, 1973.
- Cushman, P. and Khurana, R. A controlled cycle of tetrahydrocannabinol smoking: T and B cell rosette formation. Life Sci. 20:971-979, 1977.
- Dixit, V.P., Arya, M., and Lohiya, N.K. The effect of chronically administered cannabis extract on the female genital tract of mice and rats. Endokrinologie 66:365-368, 1975.
- Finnegan, L.P. Pulmonary problems encountered by the infant of the drug-dependent mother. Clin. Chest Med. 1:311-325, 1980.
- Fried, P.A. Marihuana use by pregnant women: Neurobehavioral effects in neonates. Drug Alcohol Dependence 6:415-424, 1980.
- Genest, P., Huy, N.D., and Roy, P.D. Toxicity study of marijuana in dogs: Effects on the mitotic index and chromosomes. Res. Commun. Psychol. Psychiatr. Behav. 1:83-290, 1976.
- Gerber, W.F. and Schramm, L.C. Effect of marihuana extract on fetal hamsters and rabbits. Toxicol. Appl. Pharmacol. 14:276-282, 1969.
- Gilmour, D.G., Bloom, A.D., Lela, K.P., et al. Chromosomal aberrations in users of psychoactive drugs. Arch. Gen. Psychiatry 24:268-272, 1971.
- Glatt, H., Ohlsson, A., Agurell, S., and Oesch, F. Delta-1-tetrahydrocannabinol and 1-alpha, 2-alpha-epoxyhexahydrocannabinol: Mutagenicity investigation in the Ames test. Mutat. Res. 66:329-335, 1979.

- Gupta, S., Grieco, M.H., and Cushman, P. Impairment of rosette-forming T-lymphocytes in chronic marihuana smokers. N. Engl. J. Med. 291:874-877, 1974.
- Hanna, J.M., Strauss, R.H., Itagaki, B., et al. Marijuana smoking and cold tolerance in man. Aviation, Space, and Environmental Medicine 47:634-639, 1976.
- Harclerode, J., Nyquist, S.E., Nazar, B., and Lowe, D. Effects of cannabis on sex hormones and testicular enzymes of the rodent, pp. 395-405. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Hembree, W.C., Nahas, G.G., Zeidenberg, P., and Huang, H.F.S. Changes in human spermatozoa associated with high dose marihuana smoking, pp. 429-439. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Herha, J. and Obe, G. Chromosomal damage in chronic users of cannabis in vivo investigation with two-day leukocyte cultures. Pharmakopsychiat. 7:328-337, 1974.
- Hingson, R., Alpert, J.J., Day, N. et al. Effects of maternal drinking and marijuana use on fetal growth and development. J. Pediatr. (in press).
- Hoffmann, D., Brunnemann, K.D., Gori, G.B., and Wynder, E.L. On the carcinogenicity of marijuana smoke, pp. 63-81. In Runeckles, V.C. (ed.) Recent Advances in Phytochemistry. New York: Plenum Publishing Corporation, 1975.
- Huang, H.F.S., Nahas, G.G., and Hembree, W.C. Effects of marihuana inhalation on spermatogenesis of the rat, pp. 419-427. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Issidorides, M.R. Observations in chronic hashish users: Nuclear aberrations in blood and sperm and abnormal acrosomes in spermatozoa, pp. 377-388. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Jakubovic, A. and McGeer, P.L. Biochemical changes in rat testicular cells in vitro produced by cannabinoids and alcohol: Metabolism and incorporation of labeled glucose, amino acids, and nucleic acid precursors. Toxicol. Appl. Pharmacol. 41:473-486, 1977.
- Jakubovic, A., McGeer, E.G., and McGeer, P.L. Effects of cannabinoids on testosterone and protein synthesis in rat testis Leydig cells in vitro. Molec. Cell. Endocrinol. 15:41-50, 1979.
- Joneja, M.G. A study of teratological effects of intravenous, subcutaneous, and intragastric administration of Δ -9-tetrahydrocannabinol in mice. Toxicol. Appl. Pharmacol. 36:151-162, 1976.
- Jones, E.P., Manno, J.E., Brown, R.D., et al. Tachycardia in rats following intracerebroventricular (ICV) administration of Δ -9-tetrahydrocannabinol (THC). Fed. Proc. Fed. Am. Soc. Exp. Biol. 39:848, 1980.

- Kaklamani, E., Trichopoulos, D., Koutselinis, A., et al. Hashish smoking and T-lymphocytes. Arch. Toxicol. 40:97-101, 1978.
- Kalofoutis, A., Koutselinis, A., Dionyssiou-Asteriou, A., and Miras, C. The significance of lymphocyte lipid changes after smoking hashish. Acta Pharmacol. Toxicol. 43:81-85, 1978.
- Kolodny, R.C., Masters, W.H., Kolodner, R.M., and Toro, G. Depression of plasma testosterone levels after chronic intensive marijuana use. N. Engl. J. Med. 290:872-874, 1974.
- Lau, R.J., Tubergen, D.G., Barr, M., and Domino, E.F. Phytohemagglutinin-induced lymphocyte transformation in humans receiving delta-9-tetrahydrocannabinol. Science 192:805-807, 1976.
- Lefkowitz, S.S. and Klager, K. Effect of delta-9-tetrahydrocannabinol on in vitro sensitization of mouse splenic lymphocytes. Immunol. Commun. 7:557-566, 1978.
- Lefkowitz, S.S., Klager, K., Nephth, D., and Pruess, M. Immunosuppression of mice by delta-9-tetrahydrocannabinol. Res. Commun. Chem. Pathol. Pharmacol. 19:101-107, 1978.
- Legator, M.S., Weber, E., Connor, T., and Stoeckel, M. Failure to detect mutagenic effects of Δ -9-tetrahydrocannabinol in the dominant lethal test, host-mediated assay, blood-urine studies, and cytogenetic evaluation with mice, pp. 699-709. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuana. New York: Raven Press, 1976.
- Leuchtenberger, C. and Leuchtenberger, R. Morphological and cytochemical effects of marijuana cigarette smoke on epithelioid cells of lung explants from mice. Nature 234:227-229, 1971.
- Leuchtenberger, C., Leuchtenberger, R., and Schneider, A. Effects of marijuana and tobacco smoke on human lung physiology. Nature 241:137-139, 1973a.
- Leuchtenberger, C., Leuchtengerger, R., Ritter, U., and Inui, N. Effects of marijuana and tobacco smoke on DNA and chromosomal complement in human lung explants. Nature 242:403-404, 1973b.
- Lewis, C.R. and Slavin, R.G. Allergy to marijuana: A clinical and skin-testing study. J. Allergy Clin. Immunol. 55:131-132, 1975.
- Liskow, B., Liss, J.L., and Parker, C.W. Allergy to marihuana. Ann. Intern. Med. 75:571-573, 1971.
- Marks, B.H. Δ -1-tetrahydrocannabinol and luteinizing hormone secretion. Prog. Brain Res. 39:331-338, 1973.
- Matsuyama, S.S., Jarvik, L.F., Fu, T., and Yen, F. Chromosome studies before and after supervised marijuana smoking, pp. 723-729. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuans. New York: Raven Press, 1976.
- McDonough, R.J., Madden, J.J., Falek, A., et al. Alteration of T and null lymphocyte frequencies in the peripheral blood of human opiate addicts: In vivo evidence for opiate receptor sites on T lymphocytes. J. Immunol. 125:2539-2543, 1980.
- Mendelson, J.H., Kuehnle, J., Ellingboe, J., and Babor, T.F. Plasma testosterone levels before, during and after chronic marihuana smoking. N. Engl. J. Med. 291:1051-1055, 1974.

- Morishima, A., Milstein, M., Henrich, R.T., and Nahas, G.G. Effects of marihuana smoking, cannabinoids, and olivetol on replication of human lymphocytes: Formation of micronuclei, pp. 711-722. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuana. New York: Raven Press, 1976.
- Morishima, A., Henrich, R.T., Jayaraman, J., and Nahas, G.G. Hypoploid metaphases in cultured lymphocytes of marihuana smokers, pp. 371-376. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses, Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Nichols, W.W., Miller, R.C., Heneen, W. et al. Cytogenetic studies on human subjects receiving marihuana and Δ -9-tetrahydrocannabinol. Mutat. Res. 26:413-417, 1974.
- Nicholson, M.T., Pace, H.B., and Davis, W.M. Effects of marihuana and lysergic acid diethylamide on leukocyte chromosomes of the golden hamster. Res. Commun. Chem. Pathol. Pharmacol. 6:427-434, 1973.
- Nir, I., Ayalon, D., Tsafiriri, A. et al. Suppression of the cyclic surge of luteinizing hormone secretion and of ovulation in the rat by Δ -1-tetrahydrocannabinol. Nature 243:470-471, 1973.
- Pertwee, R.G. and Tavendale, R. Effects of Δ -9-tetrahydrocannabinol, 2,4-dinitrophenol and pentolinium tartrate on behavioral thermoregulation in mice. Br. J. Pharmacol. 66:39-50, 1979.
- Petersen, B.H., Lemberger, L., Graham, J., and Dalton, B. Alterations in the cellular-mediated immune responsiveness of chronic marihuana smokers. Psychopharmacol. Commun. 1:67-74, 1975.
- Petersen, B.H., Graham, J., and Lemberger, L. Marihuana, tetrahydrocannabinol and T-cell function. Life Sci. 19:395-400, 1976.
- Petersen, B.H. and Lemberger, L. Effect of delta-9-tetrahydrocannabinol administration of antibody production in mice. Fed. Proc. 35:313, 1976.
- Pruess, M.M. and Lefkowitz, S.S. Influence of maturity on immunosuppression by delta-9-tetrahydrocannabinol. Proc. Soc. Exp. Biol. Med. 158:350-353, 1978.
- Purohit, V., Ahluwalia, B.S., and Vigersky, R.A. Marihuana inhibits dihydrotestosterone binding to the androgen receptor. Endocrinology 107:848-850, 1980.
- Rachelefsky, G.S., Opelz, G., Mickey, M.R., et al. Intact humoral and cell-mediated immunity in chronic marijuana smoking. J. Allergy Clin. Immunol. 58:483-490, 1976.
- Rosenkrantz, H. In Report of an Addiction Research Foundation/World Health Organization (ARF/WHO) Scientific Meeting on Adverse Health of Behavioral Consequences of Marijuana Use. ARF/WHO, Toronto, 1981.
- Sassenrath, E.N., Banovitz, C.A., and Chapman, L.F. Tolerance and reproductive deficit in primates chronically drugged with Δ -9-THC. Pharmacologist 21:201, 1979.
- Seid, D.A. and Wei, E.T. Mutagenic activity of marihuana smoke condensates. Pharmacologist 21:204, 1979.

- Shapiro, C.M., Orlina, A.R., Unger, P., and Billings, A.A. Antibody response to cannabis. JAMA 230:81-82, 1974.
- Shapiro, C.M., Orlina, A.R., Unger, P.J., et al. Marihuana-induced antibody response. J. Lab. Clin. Med. 88:194-201, 1976.
- Silverstein, M.J. and Lessin P. 2,4-dinitrochlorobenzene skin testing in chronic marihuana users, pp. 199-203. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuana New York: Raven Press, 1976.
- Smith, C.G., Besch, N.F., Smith, R.G., and Besch, P.K. Effect of tetrahydrocannabinol on the hypothalamic-pituitary axis in the ovariectomized Rhesus monkey. Fertil. Steril. 31:335-339, 1979.
- Smith, C.G., Besch, N.F., and Asch, R.H. Effects of marihuana on the reproductive system, pp. 273-294. In Thomas, J.A. and Singhal, R.L. (eds.) Advances in Sex Hormone Research Baltimore: Urban and Schwarzenberg, 1980.
- Stenchever, M.A. and Allen, M. The effect of delta-9-tetrahydrocannabinol on the chromosomes of human lymphocytes in vitro. Am. J. Obstet. Gynecol. 114:819-821, 1972.
- Stenchever, M.A., Kunysz, T.J., and Allen, M.A. Chromosome breakage in users of marihuana. Am. J. Obstet. Gynecol. 118:106-113, 1974.
- Tyrey, L. Δ -9-Tetrahydrocannabinol suppression of episodic luteinizing hormone secretion in the ovariectomized rat. Endocrinology 102:1808-1814, 1978.
- Tyre, L. Δ -9-Tetrahydrocannabinol: A potent inhibitor of episodic luteinizing hormone secretion. J. Pharmacol. Exp. Ther. 213:306-308, 1980.
- Wehner, F.C., Van Rensburg, S.J., and Thiel, P.G. Mutagenicity of marijuana and transkei tobacco smoke condensates in the salmonella/microsome assay. Mutat. Res. 77:135-142, 1980.
- White, S.C., Brin, S.C., and Janicki, B.W. Mitogen-induced blastogenic responses of lymphocytes from marihuana smokers. Science 188:71-72, 1975.
- Zimmerman, A.M. and Raj, A.Y. Influence of cannabinoids on somatic cells in vivo. Pharmacology 21:277-287, 1980.
- Zimmerman, A.M., Stich, H., and San, R. Nonmutagenic action of cannabinoids in vitro. Pharmacology 16:333-343, 1978.
- Zimmerman, A.M., Bruce, W.R., and Zimmerman, S. Effects of cannabinoids on sperm morphology. Pharmacology 18:143-148, 1979.

6

BEHAVIORAL AND PSYCHOSOCIAL EFFECTS OF MARIJUANA USE

The mind-altering effects of marijuana underlie its widespread and increasing popularity. Marijuana users who experience effects on mood, perception, and motivation report that they seek the "high" and the "mellowing-out." However, under some circumstances many of these same effects can be considered adverse. Perceptual distortions that are sought by users pose risks for driving cars or using other machines. There is reason to be concerned about effects on learning by students using marijuana in school. Older adults receiving Δ^9 -THC as therapy may be highly intolerant of altered consciousness and perceived loss of control. Thus, it has become a matter of practical as well as scientific interest to learn more about the effects of marijuana on the brain and behavior.

Many psychological and neuropsychological studies have been conducted to investigate specific effects of marijuana on behavior. These include studies of intellectual functions, such as memory, attention, sequential information processing, and decision-making, as well as perceptual and psychomotor functions. There is a methodological challenge in trying to design experiments that will discriminate reliably among these functions and determine precisely which is being affected when a drug produces a particular behavioral outcome. For example, one's ability to process and respond to environmental stimuli represents a chain of events. The sequence begins with a sensation or perception. Drugs can influence the manner, speed, and accuracy with which this input is received. The information must then be stored in memory, even if only very briefly, and then retrieved from memory to be integrated with recalled prior experiences and other sensory inputs. The response from the subject is the result of the integration of new and old information. A drug acting at any point in this chain of events can alter behavioral performance.

Studies of the effects of marijuana on complex behavior must be carefully interpreted, because there are numerous variables that can influence the results. First, there is the drug itself. The dose, type of preparation, route of administration, and speed of administration must be specified. Next, the user--his personality, level of innate ability, motivation to perform, and especially his previous experience with marijuana, are powerful influences on test results. Finally, there is the type of behavioral test and the setting in

which it is performed. Simpler and well-practiced skills are less susceptible to disruption by drug effects than are novel or complex tasks. The studies in the literature vary in their attention to these factors.

Most of these studies have been carried out on male college students who volunteer for marijuana research. Although this age-group (19-25) represents a period of peak use of marijuana, it cannot be assumed that findings from a college population will generalize to other sectors of the youth population. The differing motives of student volunteers seriously confound the interpretation of results in intellectual areas, where it has been established that motivation plays a significant role in determining performance. Some dedicated users want to do well and demonstrate that marijuana has no harmful effect. Others are simply interested in obtaining the drug and enjoying its effects with little interest in the experiment. Additional methodological issues that recur in this body of research include: (a) reliance on self-reports by subjects regarding personal history of frequency and intensity of drug use, (b) occasional reliance on self-reports of drug dose and level of intoxication at the time of the experiment, (c) lack of standardized dosages and methods of administration of Δ -9-THC even when the drug is administered by the investigator, and (d) lack of attention to motives and beliefs of users and nonusers with whom they interact.

A representative sample of studies will be reviewed here, and a summary table of 88 reports of the relationship between marijuana use and behavioral and psychosocial functioning is available from the Institute of Medicine by request.

PERCEPTUAL AND PSYCHOMOTOR FUNCTIONS

Acute Effects

The studies reported here cover the range of commonly used doses* from very low up to 0.250 mg/kg of Δ -9-THC in marijuana cigarettes at a single sitting. These are acute effects--changes that can be seen after a single dose. The effects begin to be seen at about the same dose level at which a "high" is perceived (0.050-0.150 mg/kg Δ -9-THC). Generally the effects are dose-related. In other words, low doses have small effects; higher doses tend to have greater effects.

*Doses are reported in milligrams per kilogram (mg/kg) where provided by the authors or as total doses in milligrams with the route of administration.

Coordination

Marijuana has been found to impair motor coordination at doses commonly used in social settings by both naive and chronic users. The functions studied include: hand steadiness (Mayor's Committee on Marihuana, 1944; Clark et al., 1970; Milstein et al., 1975), body sway (Mayor's Committee on Marihuana, 1944; Kiplinger et al., 1971; Evans et al., 1973), and accuracy of execution movements (Rafaelsen et al., 1973; Milstein et al., 1975; Kvalseth, 1977). Studies have also showed a dose-related increase in impairment of postural stability as measured by increased body sway (Kiplinger et al., 1971).

Reaction Time

Reaction time is defined as the time lag between a signal and the response a subject makes to that signal. Most studies examine the time that it takes a subject to respond to a visual or auditory signal. The effects of marijuana on either speed of initial detection of the signal or speed of response have been inconsistent at doses commonly used in social settings ("low to moderate"). The same subjects are impaired at some times, but not at other times (Mayor's Committee on Marihuana, 1944; Clark et al., 1970; Dornbush et al., 1971; Moskowitz et al., 1972, 1974; Borg et al., 1975; Schaefer et al., 1977; Peeke et al., 1976; Stillman et al., 1977). The meaning of this inconsistency is uncertain, but it probably involves an effect on attention mechanisms. When a subject is intoxicated with marijuana, he is probably less likely to attend to the reaction time task. Perhaps it is when he does pay attention to the task that function on this test is not impaired.

Tracking

Tracking is the term used to describe the act of following a moving stimulus. It is an important component of driving and flying skills. Tracking behavior is highly sensitive to the effects of marijuana. Impairment of tracking occurs even at very low doses (4.5 mg by smoking) in naive subjects (Weil et al., 1968). Studies of experienced users have also demonstrated consistent impairment. The tracking impairment has been found to persist for 4 to 8 hours, well beyond the feeling of intoxication ("high") by one laboratory (Moskowitz and Sharma, 1979; Moskowitz et al., 1981). No other studies have measured the effects of marijuana beyond 2 hours. This finding on the long-lasting effects has very important implications, as will be discussed later when the effects of marijuana on driving are reviewed, and, therefore, such studies should be repeated by other investigators.

While reaction time studies (as noted above) showed inconsistent results, tracking behavior is regularly and significantly diminished by marijuana at doses usually used in social settings. Tracking

tasks differ from reaction time studies, because the subject must continuously pay attention to the task. Since reaction time tests are intermittent, continuous attention is not required, and this may explain why reaction time studies fail to show consistent marijuana effects.

Sensory and Perceptual Functions

Tests that measure a subject's ability to detect a brief flash of light show significant impairment by low to moderate doses (2-3 mg are examples) of smoked marijuana (Sharma and Moskowitz, 1972, 1973, 1974; Moskowitz et al., 1972, 1974; Casswell and Marks, 1973; Jones and Stone, 1970). Sustained attention is required in signal detection tasks, and the relation between this sustained attention requirement and motivation effects has not been explored. Signal detection tasks are prototypes of perceptual demands found in man-machine interactions. The large reductions in signal detection that occur under the influence of marijuana may suggest a substantial risk for users who are operating machines. Other visual functions, such as visual search, that depend on eye movements are not impaired.

Intellectual and Cognitive Functions

The effects of marijuana on such intellectual and cognitive functions as verbal fluency, short-term memory, learning ability, calculation skills, ability to follow complex directions, and time sense have been investigated and are reported below. However, this area of study has been hampered by the lack of standard measures of functioning in the intellectual and cognitive areas tested. Overall, the investigation of marijuana effects on intellectual and cognitive functioning has not followed a logical progression.

Learning and Memory When studying the effects of drugs on learning, it is difficult to control all of the factors that might influence the results; for example, as noted above, how hard a subject tries to perform can make a big difference even in the presence of a sedating drug. Thus, it is not surprising that early studies of marijuana's effects gave inconsistent results.

More recently, several studies have demonstrated that a single moderate dose of marijuana impairs short-term memory. This effect is especially noticeable in the phases of short-term memory that are heavily dependent on attention, such as information-acquisition and storage (Abel, 1970, 1971; Dornbush et al., 1971; Dittrich et al., 1973; Melges et al., 1974; Belmore and Miller, 1980). Examples of the types of impaired tasks would be remembering a sequence of numbers or syllables or memorising and following a sequence of directions.

Physiological changes have been monitored in some of the same studies in which intellectual impairment has been reported. Miller

and Cornett (1978) found that increases in heart rate are produced by marijuana to about the same degree as impairment on intellectual tasks. This linking of a physiological marker with studies of behavioral effects is a useful model for research in this field.

Time Sense Another intellectual function influenced by marijuana is time sense. Under the influence of moderate doses of the drug, most investigators report that subjects consistently overestimate the amount of time that has elapsed. Thus, under the influence of marijuana, a given event is reported to last longer than it actually does last (Clark et al., 1970; Vachon et al., 1974; Tinklenberg et al., 1976a).

State-Dependent Learning State-dependent learning refers to a situation in which material that is learned while under the influence of a drug is remembered best in the state of drug intoxication in which it was originally learned. A series of studies were conducted with oral doses of 2 mg (in a subsequent study this dose was calibrated to 0.3 mg/kg) of Δ -9-THC to investigate the extent to which learning and memory are linked to the state of intoxication (Darley et al., 1973a,b, 1974, 1977). This modest dose of marijuana caused learning to take place more slowly than when the subject was drug-free. Once learned, recall of the learning that occurred during intoxication was best when the subject was again under the influence of marijuana. Although state-dependent learning occurs with marijuana, the quality of learning and recall is impaired because the information or problem-solving skills learned in the marijuana-intoxicated state will be reduced or impaired. These investigators believe that the major deficit is in the attention-storage-phase of learning.

Oral Communication

Marijuana use in low to moderate doses impairs oral communication, especially clarity of sequential dialogue with other persons (Dornbush et al., 1971; Paul and Carson, 1973; Zeidenberg et al., 1973; Crockett et al., 1976; Miller et al., 1977a-d, 1978a,b, 1979; Pfefferbaum et al., 1977; Miller and Cornett, 1978; Natale et al., 1979; Belmore and Miller, 1980). Marijuana at moderate doses disrupts continuity of speech by impairing short-term memory (6-18 seconds duration) (Belmore and Miller, 1980). Communication while intoxicated is also impaired by the intrusion of irrelevant words and ideas into the stream of communication. When a list of words is learned and then the subjects are asked to recall those words without regard to sequence, words that were never in the original list are inserted during recall more often by subjects given Δ -9-THC than by those who were drug-free (Pfefferbaum et al., 1977; Miller and Cornett, 1978; Miller et al., 1978a,b). Zeidenberg et al. (1973) administered 5 mg Δ -9-THC orally and found that, in a social context, phrases became shorter, speech became slower, and there was

greater lag time between the cue to talk and the actual onset of talking. These subjects were also less able to recognize three-letter nonsense syllables to which they had previously been exposed. Further, when experimental subjects were all given the same dose of Δ -9-THC, they reported different subjective levels of intoxication. Those who reported more intoxication showed greater disruption of two-person communication (Paul and Carson, 1973).

Experimental subjects who were asked to tell stories about ambiguous pictures (the Thematic Apperception Test) demonstrated drug impaired organization and integration of stories. The authors reported "a timeless, nonnarrative quality, with greater discontinuity in thought sequence and more frequent inclusion of contradictory ideas" (Roth et al., 1975). When asked to talk for five minutes on any topic, subjects under Δ -9-THC demonstrated decreased variability of language and an increase in personal references, as well as less detailing of items mentioned in the monologue and less critical evaluation of those items (Natale et al., 1979).

Auto Accidents

Simulator Studies

A driving simulator is a laboratory instrument that requires the subject to perform a sample of the behavior required in automobile driving situations. Simulators differ from most of the laboratory studies described above in that complex behavior is required. Although simulators are representative of the multitask character of driving, no one simulator is capable of presenting all aspects of driving simultaneously. The behavior sampled varies across simulators; however, in comparison to car driving situations, the simulator has the advantage of presenting a standard stimulus to all subjects.

Most simulator studies reveal impairment of driving skills following moderately intoxicating doses of marijuana such as 10-15 mg (Crancer et al., 1969; Dott, 1972; Ellingstad et al., 1973; Rafaelsen et al., 1973; Moskowitz et al., 1976; Smiley et al., 1981). These impairments have been reported in simulators that test the perceptual functions as well as those that test motor skills of car control.*

*Another type of simulator study examined marijuana's effect on performance in a flying simulator (Janowsky et al., 1976). Subjects smoked marijuana cigarettes with 0.09 mg/kg Δ -9-THC, a dose of Δ -9-THC commonly used in social settings. Significant impairment of short-term memory was noted. Subjects were unable to recall where they were in the execution of a task. On the simulator they tended to forget where they were in a given flight sequence.

Test Courses

Experimental studies of the effects of marijuana on closed course automobile driving performance show that this skill is impaired by marijuana. Car handling skills were reduced, as shown by objective measures (Klonoff, 1974; Hansteen et al., 1976; Attwood, in press). It should be noted that these studies, involving subjects under the influence of marijuana, examined performance in less complex situations than are actually met in real-life driving situations. However, a closed course has the advantages of standard conditions and safety factors. In real-life driving situations, the perceptual and cognitive demands are considerably more complex. The Klonoff (1974) study of driving performance on city streets indicates that smoked marijuana (5-10 mg Δ -9-THC) impairs judgment and concentration in addition to impairing car handling skills.

Accident Surveys

Experimental evidence of impairments caused by marijuana on psychomotor functions, judgment, and motor skills involved in driving has led to research on the relationship of the use of marijuana and automobile accidents. A strongly positive relationship between use of alcohol and increased driving risk has long been established. The techniques used to establish the relationship of alcohol to accidents might appear to offer an excellent paradigm for comparable marijuana-accident research. However, there have been practical reasons why the roadside survey model of using breath samples obtained from accident drivers and comparing those to breath samples of randomly selected drivers who are passing the accident site in the same direction, the same time of day, and same day of the week has not worked for marijuana studies. Whereas there has been 97 percent cooperation for alcohol breath analysis, marijuana determination requires a blood sample, and only a minority of drivers willingly cooperate. Further, marijuana has a quite different body distribution pattern due to its high fat solubility. Delta-9-THC is not only technically quite difficult to detect in samples of body fluid, but it may be active in the nervous system long after it is not detectable in blood. The detrimental effects on driving skills (Moskowitz and Sharma, 1979; Moskowitz et al., 1981) may even persist 4 to 8 hours beyond the time when the user has had subjective feelings of euphoria or sleepiness.

Several reports of accident surveys have recently been published (Teale et al., 1977; Cimbura et al., 1980; McBay and Owens, 1981), but all suffer from the problems discussed above and particularly from the lack of a reasonable comparison group. For example, one study reported that 16 percent of Boston drivers had Δ -9-THC in their blood (Sterling-Smith, 1975). There was no description of the group who declined to give a blood sample but provided breath or urine samples instead. Also, there is no information as to the frequency of finding Δ -9-THC in the blood of those drivers who have

not had an accident or otherwise come to police attention. In addition, many users of marijuana also use other drugs so that data are available on only a few subjects who only used marijuana.

In an effort to obtain some reference point for the association of marijuana with accidents as compared with other drugs, Warren et al. (in press) reanalyzed the Cimbura et al. (1980) data. Twelve percent of the fatally injured drivers and pedestrians in that study had been found to have Δ -9-THC in their blood. The presence of other drugs was also determined and a culpability index was developed. A culpability index compares the frequency that a drug is found in drivers assigned responsibility for causing a collision with the frequency in individuals from the same sample who had not caused an accident.

Aspirin was found to have a culpability index of 1.0. That is, it was no more frequent in individuals assigned responsibility for a collision than on those who were not. This is of some significance because it serves as an internal check on the technique, agreeing with the a priori assumption that it would be unlikely for aspirin users to be overrepresented among those responsible for accidents. In contrast, subjects with cannabinoids present in the urine were found to have a culpability index of 1.7, the same culpability level found for the presence of alcohol. This indicates an excess of Δ -9-THC-positive drivers in the category responsible for accidents. The presence of antihistamines produced a culpability index of 1.5, and tranquilizers/antidepressants, 1.8.

Given the difficulties in executing epidemiologic studies where it is so difficult to obtain adequate control groups, it would appear that only tentative conclusions about marijuana's role in accidents can be reached. Supportive evidence that marijuana is a contributing cause of accidents comes from surveys of marijuana users who report they receive a higher-than-average number of tickets for driving violations and are involved in a higher-than-average number of accidents (Johnston, 1980). Nevertheless, the problems described above are yet to be solved. But the culpability index model presents a methodology that may be refined and utilized in future studies.

Alcohol-Marijuana Interactions.

Surveys show that marijuana and alcohol are frequently consumed together (Fishburne et al., 1980; Johnston et al., 1980). Thus, it is important to determine what interactions, if any, occur between these two drugs. As both drugs have sedative properties, an additive effect would be expected and has been found in the few systematic investigations of the effects of this combination. One study reported that 0.05 percent blood alcohol level concentration (BAC) increased the impairment produced by 5 mg of smoked Δ -9-THC on tracking behavior (Manno et al., 1971). In a study using two doses of alcohol and two doses of marijuana, even the low dose of alcohol (0.07 percent BAC) and the low dose of Δ -9-THC (1.4 mg) impaired complex tracking in an additive fashion (Hansteen et al., 1976).

Higher doses produced more pronounced decrements. A combination of Δ -9-THC (0.320 mg/kg) and ethanol (a dose that produces a peak blood level of less than 0.08 percent BAC) has also produced an additive effect on the ability to perform on a psychomotor test (Belgrave et al., 1979). This additive effect would be of concern to those operating a motor vehicle.

The issue of alcohol-marijuana interactions is an important one, but currently few data are available. Clearly, more studies of marijuana's interaction with alcohol and other commonly used drugs are needed.

Chronic Effects

Animal Studies

Studies of chronic effects are necessary to determine whether a drug produces changes that persist after administration has stopped. In view of the theoretical possibility of cumulative or persistent marijuana effects, it is surprising that only a few laboratories have conducted experiments involving repeated dosing and testing for residual effects. Mice injected with 10 mg/kg Δ -9-THC for 20-40 days were found to be persistently impaired in new learning 100 days after the injections stopped (Radouco-Thomas et al., 1976). Similarly, rats given 20 mg/kg of Δ -9-THC orally for 180 days had learning still impaired 2 months after the Δ -9-THC treatment stopped (Fehr et al., 1976). This was confirmed by the same group in two subsequent studies (Fehr et al., 1979; Stiglick and Kalant, in press). Another group of investigators, however, could find no residual learning effects in monkeys 20 days after stopping comparable doses of Δ -9-THC (Ferraro and Grilly, 1974).

Human Studies

Clinical reports of memory impairment, lack of concentration, lethargy, etc., in nonintoxicated chronic users of marijuana have led to studies in which psychological testing was administered to users of marijuana and controls. The results of these studies are inconclusive. Several studies show impaired performance in users as compared to controls (Agarwal et al., 1975; Soueif, 1976; Wig and Varma, 1977; Mendhiratta et al., 1978); others found no significant residual effects in the marijuana users (Bowman and Pihl, 1973; Rubin and Comitas, 1975; Satz et al., 1976; Ray et al., 1978; Schaeffer et al., 1981). All of these studies can be criticized on methodological grounds, and the results have been disputed. This is not surprising, because it is technically very difficult to obtain a sample of chronic marijuana users, get them into a truly drug-free condition, test them, and similarly test an appropriate group of controls.

Several groups of investigators (Dornbush et al., 1972; Frank et al., 1976; Harshman et al., 1976; Rossi et al., 1977) examined

chronic marijuana users before and after 21-94 days of chronic intoxication in a research hospital setting. None of the investigators found any psychological changes during postdrug tasting. However, 2 months of use is a relatively short period of time for a change to be detected, and the subjects had already been using marijuana for at least a year prior to entering each study (Fehr et al., 1976).

The available studies of chronic behavioral effects lead to no clear conclusions. Although some animal studies demonstrated a learning deficit that persisted for months after daily marijuana exposure was discontinued, the human studies have such methodological weaknesses that they cannot be interpreted. A prospective concurrent cohort study and a retrospective case-control study of possible outcomes of and risk factors for use of marijuana could add useful information. (See research recommendations at the end of this chapter.)

CLINICAL SYNDROMES

In this section we will discuss both acute and chronic behavioral changes that have been reported in the clinical literature to be associated with the use of marijuana. An association based on case reports does not imply causality. Studies of appropriate control groups are necessary. In general, acute or immediate clinical effects of drugs can be determined scientifically much more readily than chronic or delayed effects. This is as true for marijuana as it is for alcohol and other drugs. Thus, the acute effects of marijuana are based on more solid evidence than are the reported chronic effects.

Acute Effects

The acute clinical effects of marijuana seem to occur on a continuum from mild dysphoria to acute brain syndrome. In the literature, three different syndromes have been described, although there is blurring of the boundaries in this classification and no general agreement as to diagnostic criteria.

Anxiety/Panic Reaction

A major portion of the evidence for this effect comes from reports by marijuana users themselves. Marijuana's popularity notwithstanding, a surprisingly high proportion of users report reactions that they regard as unpleasant or undesirable. For example, 32 percent of regular users reported that while intoxicated they occasionally experienced such symptoms as acute panic, paranoid reaction, hallucinations, and unpleasant distortions in body image (Tart, 1970; Negrete and Kwan, 1972). Another study reported that 16 percent of

regular users reported anxiety, fearfulness, confusion, dependency, or aggressive urges as a "usual occurrence" (Halikas et al., 1971). Similar findings in groups of stable, well-adjusted, moderate users have been found by other investigators (Annis and Smart, 1973; Marcus et al., 1974). First-time users are more likely than are experienced users to report adverse reactions. The frequency of such reactions appears to be higher when the setting for use is not a favorable one; for example, when the user sees the environment as threatening.

These adverse psychological reactions also have been observed in subjects of laboratory experiments with marijuana. Such controlled observations of persons whose immediate prior mental status and whose dosage were known give a basis for concluding that acute adverse psychological reactions can occur under single moderate doses of marijuana. These effects are more likely at higher doses. They usually last no longer than 2 to 4 hours. Acute paranoid reactions under these controlled conditions have been reported (Mendelson and Meyer, 1972; Tassinari et al., 1973; Frank et al., 1976; Melges, 1976). Ingestion, in which titration of dose (dose adjustment as occurs during smoking) is difficult, may be more likely to produce adverse effects than administration by smoking marijuana. However, chronic use and interaction with other psychoactive substances are not required.

As frequently as these adverse reactions are observed and self-reported, medical treatment is rarely sought. For example, a college student health clinic reported only six students per year sought medical treatment for an adverse reaction to marijuana out of a student population of 20,000 (Pillard, 1970). In the general population, a diagnosis of acute cannabis reaction was found in only 10 cases out of 700,000 hospital admissions in the United States (Lundberg et al., 1971). In the U.S. Army, only 18 such cases were treated over a several-year period from a military population of 33,000 (Tennant and Groesbeck, 1972). There are no recent figures showing requests for medical treatment now that the use of marijuana is more intense, widespread, and reaching younger age-groups. However, a unique monitoring of drug causality behavior documenting emergency room encounters conducted by the Drug Enforcement Administration and the National Institute on Drug Abuse (U.S. Department of Health and Human Services, 1979) may in the future provide additional information about the frequency of adverse reactions to use of marijuana.

Dysphoric Reaction

Therapeutic trials have been carried out testing Δ -9-THC as a possible treatment for mood disorders (see Chapter 7). Severe dysphoric reactions characterized by disorientation, catatonialike immobility, acute panic, and heavy sedation have occurred in several patients. The dysphoric symptoms appeared at moderate doses comparable to those used in social settings. They lasted only a few hours and responded to discontinuation of the drug and reassurance of the patients (Kotin et al., 1973; Ablon and Goodwin, 1974).

Similar dysphoric reactions have been reported in cancer patients who were on a therapeutic trial of Δ -9-THC to control the nausea associated with chemotherapy. The symptoms, course, and response to ceasing use of the drug were identical to those described above. Investigators have suggested that the dysphoric response is more likely to occur in older patients not accustomed to drug use for whom the mood-altering effects are unanticipated and unwelcome (Shilling and Stillman, 1980).

Acute Brain Syndrome

Diagnostic criteria for the syndrome now called delirium and previously called acute brain syndrome appear in Diagnostic and Statistical Manual of Mental Disorders, Third Edition, 1980 (DSM III). These include: (a) a clouding of consciousness as manifested by impairment of ability to sustain attention to environmental stimuli, or impairment of ability to sustain goal-directed thinking or goal-directed behavior; (b) a disorder of memory or orientation; (c) perceptual disturbances; and (d) a change in sleep pattern and/or a change in psychomotor activity. The symptoms develop over a short period of time and fluctuate rapidly.

Both the symptom pattern and the course of the acute brain syndrome fit the descriptions of one type of behavior disorder associated with use of marijuana. It has been reported to develop in persons who have a history of prolonged, regular, heavy use of marijuana. It is defined as an "acute" brain syndrome because it comes on during the period of drug use and it gradually disappears after the drug is stopped. The majority of case reports have come from Eastern countries where the cannabis products customarily used have high potency (Spencer, 1970; Chopra and Smith, 1974; Meyer, 1975). It has also been reported in U.S. Army personnel stationed in Viet Nam (Talbot and Teague, 1969) and in Europe (Tennant, 1972), where soldiers had access to very high Δ -9-THC concentrations in cannabis substances. In contrast to the Indian public mental hospital patients who were hospitalized for many weeks, U.S. soldiers recovered in 3 to 11 days and returned to duty. This difference in duration may reflect sociocultural differences in length of in-patient treatment more than a difference in the disorder.

Withdrawal Syndrome

Studies of animals and human subjects given moderate to high doses of marijuana orally or by inhalation several times per day have demonstrated tolerance to many of the effects of marijuana (see Chapter 1). When such use of marijuana is stopped after several days, a withdrawal syndrome occurs. In human subjects, this resembles the typical mild withdrawal symptoms seen after prolonged sedative use (Jones and Benowitz, 1976). Subjects show irritability, agitation, insomnia, and EEG changes (see Chapter 4). These symptoms are self-limiting; they peak at 30 hours and disappear by 90 hours.

There is no clinical evidence that physical dependence plays an important role in persistent use of marijuana. Withdrawal symptoms would not be expected in intermittent users; however, daily round-the-clock users of high-dose marijuana may be expected to show some symptoms of withdrawal soon after stopping regular use.

Chronic Effects

Cannabis Psychosis

Cannabis psychosis refers to a chronic psychotic condition (out of contact with reality) reportedly seen in heavy marijuana users, but extending beyond the period of acute intoxication. Some authors have described a schizophrenialike picture with delusions and hallucinations, and others have stressed the existence of organic mental confusion. Most of the reports have come from observation of hospitalized patients in Asian and African countries (Asuni, 1964; Chopra and Smith, 1974; Thacore and Shukla, 1976). There are no reports in the North American literature. At this time, there is insufficient evidence to say that cannabis psychosis exists as a separate clinical entity (Murphy, 1963; Edwards, 1976).

"Amotivational Syndrome"

Clinicians coined the term "amotivational syndrome" to describe a characteristic set of personality changes seen in some daily users of marijuana (McGlothlin and West, 1968; Smith, 1968). The changes include apathy, loss of ambition, loss of effectiveness, diminished ability to carry out long-term plans, difficulty in concentrating, and a decline in school or work performance. As usually described, these changes are seen in frequent or daily users, and thus they may be considered a form of chronic intoxication. The term "amotivational syndrome" is not an official diagnosis, but there is agreement among many clinicians who treat young people that this constellation of symptoms is common. It may also be seen in nonmarijuana users, and daily use of marijuana is not always associated with loss of motivation.

The evidence presented for the linking of this syndrome with marijuana consists of case reports. For example, Baker and Lucas (1969) described the case of a man whom friends described as previously conscientious, capable, and effective; but after smoking hashish daily for 3 years, he changed into a person for whom use of drugs was a way of life and in whom a serious deterioration of social function was observed. Other reports consist of groups of cases with similar histories (Thurlow, 1971). The symptoms mentioned, in addition to loss of motivation, include falling grades, difficulties in concentration, intermittent confusion, and impaired memory. Some authors report improvement when use of marijuana is stopped (Kolansky and Moore, 1971, 1972).

A variety of other data support such a condition. In a large survey, daily marijuana users were asked about the drug's adverse effects (Johnston et al., 1980). The most common response was "loss of energy" (42 percent). Nearly a third (32 percent) of the daily users thought that marijuana caused them to be less interested in other activities than they had been before, and a third (34 percent) thought that it hurt their school and/or job performance. Another type of evidence comes from comparisons of college students who use marijuana with others who do not. Several such studies (Shean and Fechtmann, 1971; Linn, 1972; Simon, 1974; Finnell and Jones, 1975) found marijuana users had increased levels of psychological disturbance, lower academic performance, and lower performance on scales measuring attitudes toward achievement and purpose in life. But some studies in both the United States and foreign countries have failed to show significant differences between marijuana users and abstainers (Brill and Christie, 1974; Rubin and Comitas, 1975).

Interpretation of the evidence linking marijuana to "amotivational syndrome" is difficult. Such symptoms have been known to occur in the absence of marijuana. Even if there is an association between this syndrome and use of marijuana, that does not prove that marijuana causes the syndrome. Many troubled individuals seek an "escape" into use of drugs; thus, frequent use of marijuana may become one more in a series of counterproductive behaviors for these unhappy people.

The available evidence does not allow a sorting of the various possibilities in the relationship between use of marijuana and the complex of symptoms in the "amotivational syndrome." It appears likely that both self-selection and authentic drug effects contribute to the "motivational" problems seen in some chronic marijuana users (see Chapter 2). Persons who are experiencing loss of motivation, apathy, and the other aforementioned symptoms probably will worsen the situation by taking any sedating drug. They should be warned to avoid frequent use of marijuana, alcohol, and other nonprescribed drugs.

"Flashbacks"

In 1968, Keeler et al. reported four cases of the brief spontaneous recurrence of a mental state similar to that experienced during marijuana intoxication 1 to 21 days after the last drug use. Three of the four subjects complained of hallucinations comparable to flashbacks usually associated with LSD (Horowitz, 1969). Three separate reports of marijuana flashbacks followed (Smith, 1968; Favazza and Domino, 1969; Weil, 1970) and all of these latter subjects had used LSD prior to marijuana. In a survey of 720 servicemen, not a single case of flashback in any subject for whom hashish was the only drug consumed was documented (Tennant and Groesbeck, 1972). But in the same sample, 15 subjects were identified who had LSD flashbacks precipitated by use of marijuana. A larger sample of 2,001 army personnel (Stanton et al., 1976) revealed that use of marijuana had the highest and only statistically

significant association with the precipitation of LSD flashbacks among five classes of abused drugs. Clinical studies also have provided evidence that marijuana precipitates a recurrence of the LSD flashbacks experience (Holsten, 1976; Abraham, 1981).

The existence of flashbacks following use of either LSD or marijuana is entirely based on self-reports, because there are no distinctive physical signs or tests, such as EEG changes, to identify this condition. There is no current pharmacological explanation of the phenomenon, and data regarding dose and time parameters do not exist. Still, the reports by users are reasonably consistent. Thus, there is clinical evidence that use of marijuana by those who have previously used LSD increases the likelihood of recurrence of the LSD experience.

Effects on Preexisting Mental Illness

The only evidence available regarding this issue consists of case reports of patients who had recovered and apparently were doing well until they used marijuana. There is no information on the number of mentally ill patients who have used marijuana without complications.

The available data, therefore, do not prove that marijuana worsens mental illness. Still, there are sufficient numbers of uncontrolled clinical reports showing a temporal association between use of marijuana and return of mental symptoms, so that patients should be warned of this possibility.

Patients with a history of schizophrenia may be particularly sensitive to marijuana's effects. Four schizophrenic patients who were otherwise well controlled with medication suffered serious relapse of their schizophrenic symptoms following use of marijuana (Treffert, 1978). Other cases have been reported (Smith and Mehl, 1970; Weil, 1970; Bernhardson and Gunne, 1972). These all were cases in which marijuana was purchased on the street, so the dose and purity were unknown.

Patients with mood disorders have also been reported to show worsening of mental symptoms after use of marijuana. For example, four cases are known in which marijuana apparently precipitated a relapse of psychotic (hypomanic) behavior (Harding and Knight, 1973). Furthermore, depressed patients treated with Δ -9-THC have been observed to show a high incidence of dysphoric reactions (Ablon and Goodwin, 1974).

Effects Sometimes Reported By Users

Mood Changes

There is a general belief that use of marijuana alters mood. This property is one of the desired effects sought by many users. Investigators have described a number of variables that enter into the mood response to marijuana (Jones, 1971). These include dosage,

past experience, attitude, expectations, and setting. For example, individuals who used marijuana in isolation tended to be relaxed and slightly drowsy; in contrast, when the user was in a group situation, marijuana was associated with euphoria and lack of sedative effect (Jones, 1971). Further evidence that mood changes are not attributable solely to the pharmacological action of marijuana comes from a study that found that elevation in mood occurred immediately before use of marijuana and immediately after, but that mood was not correlated with other indicators of the subjective level of intoxication (Rossi et al., 1978). Instead, mood was correlated significantly with the moods of others, whether or not the other persons were intoxicated.

It appears that preexisting mood can influence the decision to use marijuana. High school students who exhibit symptoms of depression are more likely than are others to begin using marijuana as well as other illicit drugs (Paton et al., 1977). There is some evidence that students use the drug as a self-prescribed remedy for their own mood problems, often reporting that they use marijuana as a means of psychological coping (Johnston et al., 1980; Kaplan, 1980).

A belief that marijuana can be used to alleviate clinical depression is not supported by other studies; including one in which Δ -9-THC was carefully tested as an antidepressant. It was given to depressed patients as an experimental treatment without success (Ablon and Goodwin, 1974) (see Chapter 7).

Interpersonal Behavior

Adolescents and young adults often report that they use marijuana to facilitate interaction in new social situations (Mirin and McKenna, 1975). In a survey of 704 midwestern undergraduate students, most reported that marijuana was a meaningful "tool of social bonding" (Linn, 1971). There seems to be a widespread belief that marijuana smoking has several facilitative effects, including enhanced social effectiveness, closer social bonding, heightened interpersonal sensitivity and empathy, and enhanced sexual pleasure. The subcultural lore on one of these measures of interpersonal behavior--sexual effects--has not been studied systematically either in surveys or in experimental studies. The effects on sex hormones are controversial (see Chapter 5). Studies in experimental situations have failed to show any enhancement of social interaction and, in fact, some decrements were noted (Galanter et al., 1974; Clopton et al., 1979; Janowsky et al., 1979). Data from natural settings rather than experimental settings are not available.

Effects on Aggression

Because marijuana users have been involved in delinquent behavior, a number of investigators have questioned whether use of marijuana enhances aggressiveness in human beings. There are specific concerns

about potential links of use of marijuana to aggression. Both retrospective and experimental studies in human beings have failed to yield evidence that marijuana use leads to increased aggression. Most of these studies suggest quite the contrary effect. Marijuana appears to have a sedative effect, and it may reduce somewhat the intensity of angry feelings and the probability of interpersonal aggressive behavior (McGuire and Megaree, 1974; Tinklenberg, 1974; Salzman et al., 1976; Taylor et al., 1976; Tinklenberg et al., 1976b; Hemphill and Fisher, 1980).

SUMMARY

There is experimental evidence that marijuana seriously impairs psychomotor performance. Strong evidence for impairment has been found in:

- coordination as examined by hand steadiness, body sway, and accuracy of execution of movement;
- tracking performance;
- perceptual tasks;
- vigilance;
- performance on automobile driving and flying simulators; and
- operating automobiles on test roadways.

Less reliable evidence of impairment or reliable evidence of a small degree of impairment was found in reaction time, simple sensory functions, and control of eye movements. Although the effects that marijuana produces on psychomotor functions used in driving are clear, studies linking marijuana to auto accidents are inconclusive. The research is impaired by methodological problems related to the pharmacology of marijuana. One recent study reported that marijuana and alcohol had a similar degree of association with fatal accidents, but more investigation is needed.

Studies also have shown acute effects of marijuana on short-term memory. State-dependent learning also has been shown, in that information or problem-solving skills learned in the intoxicated state will be reduced or impaired in the drug-free state. One laboratory has shown tracking impairment to persist for 4 to 8 hours beyond the feeling of intoxication. Some animal studies demonstrate a learning deficit that persists for months after marijuana exposure has been discontinued, but human studies do not permit secure conclusions.

The acute clinical effects of marijuana are fairly well established, although there is no general agreement as to how to classify them. Anxiety and panic reactions have been reported by users and observed in experimental situations. They are not uncommon, but they rarely require medical attention. When marijuana is used to treat nausea and other conditions, mental effects can occur, which some patients, especially older persons, may regard as unpleasant. These mental effects may require cessation of the treatment.

Marijuana also has been found to produce an acute brain syndrome. This is a more severe mental problem consisting of confusion and loss of contact with reality. It lasts from several hours to several days and appears to be more likely to occur with higher doses.

Chronic effects of any drug are more difficult to assess than are immediate effects. The evidence that marijuana produces a chronic psychosis is not convincing. The possible role of marijuana in causing an amotivational syndrome is a matter of great concern. Apathy, poor school work or work performance, and lack of goals characterize a number of long-term marijuana users. But it has not been possible to determine how much is caused by use of marijuana and how much was antecedent; it seems likely that both factors (drug effect and self-selection) contribute to the motivational problems seen in chronic users of marijuana. Existing studies have produced conflicting results. None of the investigators has looked at effects on the very young daily marijuana user, who is regarded as potentially at high risk for damaging effects because of physiological and psychological immaturity.

There is clinical evidence that marijuana use by former LSD users may precipitate a recurrence of LSD-type hallucinations known as a "flashback." Other clinical evidence raises the possibility that marijuana use can worsen preexisting mental illness.

RECOMMENDATIONS FOR RESEARCH

The committee recommends the following types of studies.

- Systematic research on acute behavioral and psychosocial effects of marijuana should be extended to other age groups. There are virtually no data on prepubertal children, young adolescents, older adults, and aging persons.

- Studies of effects of daily use of marijuana on school children are greatly needed. These effects could include the learning of new material, physical, psychological, and social development, acquisition of coping skills, and tools of daily living.

- Systematic studies of long-term effects of marijuana are increasingly possible now that longitudinal studies have identified representative panels of persons known to be chronic heavy users. These studies should cover interactive effects of marijuana and other drugs on behavioral and psychosocial responses, especially interactions of alcohol and marijuana because of their frequency of associated use.

- Dosage effects should be restudied, taking into account the higher potency cannabis that is in current use. Further study is needed of the timing and depth of inhalation of cigarettes with standard doses of marijuana. More animal studies at varying doses are needed. In view of the long-term retention of marijuana in body tissues, further study is needed to see whether or not chronic users may have improvements of function even in the absence of an acute dose

of marijuana. The factors that influence the persistence of effects following an acute dose are not understood.

- The correlation of changes in a physiological marker, such as increased heart rate, with observations of behavioral effects should be encouraged.

- Many of these recommendations, along with those of other chapters, could be consolidated and carried out as part of a study that is both a prospective cohort study and a retrospective case-control study of possible outcomes and risk factors with marijuana use.

- A cohort of drug-naive junior high school students could be assembled and followed over time to see which students become marijuana users and which remain nonusers. Students would be subjected to physical and psychosocial testing at predetermined time intervals. The two groups would be evaluated in terms of the incidence of specific outcomes and the relative risks associated with these outcomes after appropriate follow-up periods.

In order to identify risk factors for marijuana use, individuals who become marijuana users would be compared to individuals who remain nonusers using a case-control methodology. By combining these two epidemiologic research strategies, the etiology and effects of marijuana use may be studied.

REFERENCES

- Abel, E.L. Marijuana and memory. Nature 227:1151-1152, 1970.
- Abel, E.L. Effects of marijuana on the solution of anagrams, memory, and appetite. Nature 231:260-261, 1971.
- Ablon, S.L. and Goodwin, F.K. High frequency of dysphoric reactions to tetrahydrocannabinol among depressed patients. Am. J. Psychiatry 131:448-453, 1974.
- Abraham, H.D. Visual disturbances in population of LSD users, Paper presented at the 134th Annual Meeting of the American Psychiatric Association, New Orleans, May 14, 1981.
- Agarwal, A.K., Sethi, B.B., and Gupta, S.C. Physical and cognitive effects of chronic bhang (cannabis) intake. Indian J. Psychiatry 17:1-7, 1975.
- Annis, H.M. and Smart, R.G. Adverse reactions and recurrences for marijuana use. Br. J. Addict. 68:315-319, 1973.
- Asuni, T. Socio-psychiatric problems of cannabis in Nigeria. Bull. Narcotics 16:17-28, 1964.
- Attwood, D.A. Cannabis, alcohol and driving: Effects on selected closed-course tasks. In Proceedings of the 8th International Conference on Alcohol, Drugs and Traffic Safety, June 15-19, 1980, Stockholm, Sweden (in press).
- Baker, A.A. and Lucas, E.G. Some hospital admissions associated with cannabis. Lancet 1:148, 1969.
- Belgrave, B.E., Bird, K.D., Chesher, G.B., et al. The effects of (-)-trans- Δ -9-tetrahydrocannabinol, alone and in combination

- with ethanol, on human performance. Psychopharmacology 62:53-60, 1979.
- Belmore, S.M. and Miller, L.L. Levels of processing and acute effects of marijuana on memory. Pharmacol. Biochem. Behav. 13:199-203, 1980.
- Bernhardson, G. and Gunne, L.M. Forty-six cases of psychosis in cannabis abusers. Int. J. Addict. 7:9-16, 1972.
- Borg, J., Gershon, S., and Alpert, M. Dose effects of smoked marihuana on human cognitive and motor functions. Psychopharmacologia 42:211-218, 1975.
- Bowman, M. and Pihl, R.O. Cannabis: Psychological effects of chronic heavy use. A controlled study of intellectual functioning in chronic users of high potency cannabis. Psychopharmacologia 29:159-170, 1973.
- Brill, N.Q. and Christie, R.L. Marihuana use and psychosocial adaptation. Arch. Gen. Psychiatry 31:713-719, 1974.
- Casswell, S. and Marks, D. Cannabis induced impairment of performance of a divided attention task. Nature 241:60-61, 1973.
- Chopra, G.S. and Smith, J.W. Psychotic reactions following cannabis use in East Indians. Arch. Gen. Psychiatry 30:24-27, 1974.
- Cimbura, G., Warren, R.A., Bennett, R.C., et al. Drugs Detected in Fatally Injured Drivers and Pedestrians in the Province of Ontario Ottawa, Ontario: Traffic Injury Research Foundation of Canada, 1980.
- Clark, L.D., Hughes, R., and Nakashima, E.N. Behavioral effects of marijuana: Experimental studies. Arch. Gen. Psychiatry 23:193-198, 1970.
- Clopton, P.L., Janowsky, D.S., Clopton, J.M., et al. Marijuana and the perception of affect. Psychopharmacology 61:203-206, 1979.
- Crancer, A., Dille, J.M., Delay, J.C., et al. Comparison of the effects of marihuana and alcohol on simulated driving performance. Science 164:851-854, 1969.
- Crockett, D., Klonoff, H., and Clark, C. The effects of marijuana on verbalization and thought processes. J. Pers. Assess. 40:582-587, 1976.
- Darley, C.F., Tinklenberg, J.R., Hollister, L.E., and Atkinson, R.C. Marihuana and retrieval from short-term memory. Psychopharmacologia 29:231-238, 1973a.
- Darley, C.F., Tinklenberg, J.R., Roth, W.T., et al. Influence of marihuana on storage and retrieval processes in memory. Mem. Cognit. 1:196-200, 1973b.
- Darley, C.F., Tinklenberg, J.R., Roth, W.T., and Atkinson, R.C. The nature of storage deficits and state-dependent retrieval under marihuana. Psychopharmacologia 37:139-149, 1974.
- Darley, C.F., Tinklenberg, J.R., Roth, W.T., et al. Marijuana effects on long-term memory assessment and retrieval. Psychopharmacology 52:239-241, 1977.
- Diagnostic and Statistical Manual of Mental Disorder, Third Edition (DSM III). Washington, D.C.: American Psychiatric Association, 1980.

- Dott, A.B. Effect of Marijuana on Risk Acceptance in a Simulated Passing Task. DHEW Publication No. (HSM)72-10010. Washington, D.C.: U.S. Government Printing Office, 1972.
- Dittrich, A., Battig, K., and Zeppelin, I.V. Effects of (-)- Δ^9 -tetrahydrocannabinol (Δ^9 -THC) on memory, attention and subjective state. Psychopharmacologia 33:369-376, 1973.
- Dornbush, R.L., Fink, M., and Freedman, A.M. Marijuana, memory, and perception. Am. J. Psychiatry 128:194-197, 1971.
- Dornbush, R.L., Clare, G., Zaks, G., et al. 21-Day administration of marijuana in male volunteers, pp. 115-128. In Lewis, M.F. (ed.) Current Research in Marijuana. New York: Academic Press, 1972.
- Edwards, G. Cannabis and the psychiatric position, pp. 321-340. In Graham, J.D.P. (ed.) Cannabis and Health. New York: Academic Press, 1976.
- Ellingstad, V.S., McFarling, L.H., and Struckman, D.L. Alcohol, Marijuana and Risk Taking. Contract No. DOT-HS-191-2-301. University of South Dakota, Vermillion, 1973.
- Evans, M.A., Martz, R., Brown, D.J., et al. Impairment of performance with low doses of marihuana. Clin. Pharmacol. Ther. 14:936-940, 1973.
- Favazza, A.R. and Domino, E.F. Recurrent LSD experience (flashbacks) triggered by marihuana. Univ. Mich. Med. Cent. J. 35:214-216, 1969.
- Fehr, K.A., Kalant, H., Leblanc, A.E., and Knox, G.V. Permanent learning impairment after chronic heavy exposure to cannabis or ethanol in the rat, pp. 495-505. In Nahas, G.G. (ed.) Marihuana: Chemistry, Biochemistry, and Cellular Effects. New York: Springer-Verlag, 1976.
- Fehr, K.O., Kalant, H., and Knox, G.V. Residual effects of high-dose cannabis treatment on learning muricidal behavior and neurophysiological correlates in rats, pp. 681-691. In Nahas, G.G. and Paton, W.D.M. (eds.) Marihuana: Biological Effects. Analysis, Metabolism, Cellular Responses Reproduction and Brain. Oxford: Pergamon Press, 1979.
- Ferraro, D.P. and Grilly, D.M. Effects of chronic exposure to Δ^9 -tetrahydrocannabinol on delayed matching-to-sample in chimpanzees. Psychopharmacologia 37:127-138, 1974.
- Finnell, W.S. and Jones, J.D. Marijuana, alcohol and academic performance. J. Drug Educ. 5:13-21, 1975.
- Fishburne, P.M., Abelson, H.I., and Cisin, I. National Survey on Drug Abuse: Main Findings: 1979. DHHS Publication No. (ADM)80-976. Washington, D.C.: U.S. Government Printing Office, 1980.
- Frank, I.M., Lessin, P.J., Tyrrell, E.D., et al. Acute and cumulative effects of marihuana smoking in hospitalized subjects: A 36-day study, pp. 673-679. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuana. New York: Raven Press, 1976.
- Galanter, M., Stillman, R., Wyatt, R.J., et al. Marihuana and social behavior. A controlled study. Arch. Gen. Psychiatry 30:518-521, 1974.

- Halikas, J.A., Goodwin, D.W., and Guze, S.B. Marijuana effects: A survey of regular users. JAMA 217:692-694, 1971.
- Hansteen, R.W., Miller, R.D., Loner, L., et al. Effects of cannabis and alcohol on automobile driving and psychomotor tracking. Ann. N.Y. Acad. Sci. 282:240-256, 1976.
- Harding, T. and Knight, P. Marihuana-modified mania. Arch. Gen. Psychiatry 29:635-637, 1973.
- Harshman, R.A., Crawford, H.J., and Hecht, E. Marihuana, cognitive style, and lateralized hemispheric functions, pp. 205-254. In Cohen, S. and Stillman, R.C. (eds.) The Therapeutic Potential of Marihuana. New York: Plenum Medical Book, 1976.
- Hemphill, R.E. and Fisher, W. Drugs, alcohol and violence in 604 male offenders referred for inpatient psychiatric assessment. S. Afr. Med. J. 57:243-247, 1980.
- Holsten, F. Flashbacks: A personal follow-up. Arch. Psychiat. Nervenkr. 222:293-304, 1976.
- Horowitz, M.J. Flashbacks: Recurrent intrusive images after the use of LSD. Am. J. Psychiatry 126:565-569, 1969.
- Janowsky, D.S., Meachom, M.P., Blaine, J.D., et al. Marijuana effects on simulated flying ability. Am. J. Psychiatry 133:384-388, 1976.
- Janowsky, D.S.; Clopton, P.L., Leichner, P.O., et al. Interpersonal effects of marijuana. Arch. Gen. Psychiatry 36:781-785, 1979.
- Johnston, L.D. The daily marijuana user. Paper presented at the meeting of the National Alcohol and Drug Coalition, Washington, D.C., September 18, 1980.
- Johnston, L.D., Bachman, J.G., and O'Malley, P.M. Highlights from Student Drug Use in America, 1975-1980. DHHS Publication No. (ADM)81-1066. Washington, D.C.: U.S. Government Printing Office, 1980.
- Jones, R.T. Marihuana-induced "high": Influence of expectation, setting and previous drug experience. Pharmacol. Rev. 23:359-369, 1971.
- Jones, R.T. and Benowitz, N. The 30-day trip--Clinical studies of cannabis tolerance and dependence, pp. 627-642. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuana. New York: Raven Press, 1976.
- Jones, R.T. and Stone, G.C. Psychological studies of marijuana and alcohol in man. Psychopharmacologia 18:108-117, 1970.
- Kaplan, H.B. Deviant Behavior in Defense of Self. New York: Academic Press, 1980.
- Keeler, M.H., Reifler, C.B., and Liptzin, M.B. Spontaneous recurrence of marihuana effect. Amer. J. Psychiatry 125:384-386, 1968.
- Kiplinger, G.F., Manno, J.E., Rodda, B.E., and Forney, R.D. Dose-response analysis of the effects of tetrahydrocannabinol in man. Clin. Pharmacol. Ther. 12:650-657, 1971.
- Klonoff, H. Effects of marijuana on driving in a restricted area and on city streets: Driving performance and physiological changes, pp. 359-379. In Miller, L.L. (ed.) Marijuana: Effects on Human Behavior. New York: Academic Press, 1974.

- Kolansky, H. and Moore, W.T. Effects of marihuana on adolescents and young adults. JAMA 216:486-492, 1971.
- Kolansky, H. and Moore, W.T. Toxic effects of chronic marihuana use. JAMA 222:35-41, 1972.
- Kotin, J., Post, R.M., and Goodwin, F.K. Delta-9-tetrahydrocannabinol in depressed patients. Arch. Gen. Psychiatry 28:345-348, 1973.
- Kvalseth, T.O. Effects of marijuana on human reaction time and motor control. Percept. Mot. Skills 45:935-939, 1977.
- Linn, L.S. Social identification and the use of marijuana. Int. J. Addict. 6:79-107, 1971.
- Linn, L.S. Psychopathology and experience with marijuana. Br. J. Addict. 67:55-64, 1972.
- Lundberg, G.D., Adelson, J., and Prosnitz, E.E. Marijuana-induced hospitalization. JAMA 215:121, 1971.
- McBay, A.J. and Owens, S.M. Marihuana and driving, pp. 257-263. In Harris, L.S. (ed.) Problems of Drug Dependence, 1980. NIDA Research Monograph No.34. DHHS Publication No. (ADM) 81-1058. Washington, D.C.: U.S. Government Printing Office, 1981.
- McGlothlin, W.H. and West, L.J. The marijuana problem: An overview. Am. J. Psychiatry 125:370-378, 1968.
- McGuire, J.S. and Megaree, E.I. Personality correlates of marijuana use among youthful offenders. J. Consult. Clin. Psychol. 42:124-133, 1974.
- Manno, J.E., Kiplinger, G.F., Scholz, N., et al. The influence of alcohol and marihuana on motor and mental performance. Clin. Pharmacol. Ther. 12:202-211, 1971.
- Marcus, A.M., Klonoff, H., and Low, M. Psychiatric status of the marihuana user. Can. Psychiatr. Assoc. J. 19:31-39, 1974.
- Mayor's Committee on Marihuana. The Marihuana Problem in the City of New York. Sociological, Medical, Psychological and Pharmacological Studies. Lancaster, Pa.: Jaques Cattell Press, 1944.
- Melges, F.T., Tinklenberg, J.R., Deardorff, C.M., et al. Temporal disorganization and delusional-like ideation. Arch. Gen. Psychiatry 30:855-861, 1974.
- Melges, F.T. Tracking difficulties and paranoid ideation during hashish and alcohol intoxication. Am. J. Psychiatry 133:1024-1028, 1976.
- Mendelson, J. and Meyer, R. Behavioral and biological concomitants of chronic marihuana smoking by heavy and casual users. Technical Papers of the First Report of the National Committee on Marihuana and Drug Abuse, pp. 68-246. Washington, D.C.: U.S. Government Printing Office, 1972.
- Mendhiratta, S.S., Wig, N.N., and Varma, S.K. Some psychological correlates of long-term heavy cannabis users. Br. J. Psychiatry 132:482-486, 1978.
- Meyer, R.E. Psychiatric consequences of marijuana use: The state of the evidence, pp. 133-152. In Tinklenberg, J.R. (ed.) Marijuana and Health Hazards: Methodological Issues in Current Research. New York: Academic Press, Inc., 1975.

- Miller, L.L., Cornett, T.L., Brightwell, D.R., et al. Marijuana: Effects on storage and retrieval of prose material. Psychopharmacology 51:311-316, 1977a.
- Miller, L., Cornett, T., Drew, W., et al. Marijuana: Dose-response effects on pulse rate, subjective estimates of potency, pleasantness, and recognition memory. Pharmacology 15:268-275, 1977b.
- Miller, L.L., McFarland, D.J., Cornett, T.L., et al. Marijuana: Effects on free recall and subjective organization of pictures and words. Psychopharmacology 55:257-262, 1977c.
- Miller, L.L., McFarland, D., Cornett, T.L., and Brightwell, D. Marijuana and memory impairment: Effect on free recall and recognition memory. Pharmacol. Biochem. Behav. 7:99-103, 1977d.
- Miller, L.L. and Cornett, T.L. Marijuana: Dose effects on pulse rate, subjective estimates of intoxication, free recall and recognition memory. Pharmacol. Biochem. Behav. 9:573-577, 1978.
- Miller, L., Cornett, T.L., and McFarland, D. Marijuana: An analysis of storage and retrieval deficits in memory with the technique of restricted reminding. Pharmacol. Biochem. Behav. 8:327-332, 1978a.
- Miller, L., Cornett, T.L., and Nallan, G. Marijuana: Effect on nonverbal free recall as a function of field dependence. Psychopharmacology 58:297-301, 1978b.
- Miller, L.L., Cornett, T.L., and Wikler, A. Marijuana: Effects on pulse rate, subjective estimates of intoxication and multiple measures of memory. Life Sci. 25:1325-1330, 1979.
- Milstein, S.L., MacCannell, K., Karz, G., and Clark, S. Marijuana-produced impairments in coordination. J. Nerv. Ment. Dis. 161:26-31, 1975.
- Mirin, S.M. and McKenna, G.J. Combat zone adjustment: The role of marihuana use. Milit. Med. 140:482-485, 1975.
- Moskowitz, H., Sharma, S., and McGlothlin, W. Effect of marihuana upon peripheral vision as a function of the information processing demands in central vision. Percept. Mot. Skills 35:875-882, 1972.
- Moskowitz, H., Shea, R., and Burns, M. Effect of marihuana on the psychological refractory period. Percept. Mot. Skills 38:959-962, 1974.
- Moskowitz, H., Ziedman, K., and Sharma, S. Visual search behavior while viewing driving scenes under the influence of alcohol and marihuana. Hum. Factors 18:417-432, 1976.
- Moskowitz, H. and Sharma, S. The effects of marihuana on skills performance. Report to the National Institute on Drug Abuse, Contract No. 271-76-3316, Southern California Research Institute, Los Angeles, Calif., 1979.
- Moskowitz, H., Sharma, S., and Ziedman, K. Duration of skills performance impairment under marijuana. American Association for Automotive Medicine Proceedings, October 1-3, 1981, pp. 87-96. San Francisco, Calif..
- Murphy, H.B.M. The cannabis habit: A review of recent psychiatric literature. Bull. Narc. 15:15-23, 1963.

- Natale, M., Zeidenberg, P., and Jaffe, J. Delta-9-tetrahydrocannabinol: Acute effects on defensive and primary-process language. Int. J. Addict. 14:877-889, 1979.
- Negrete, J.C. and Kwan, M.W. Relative value of various etiological factors in short lasting, adverse psychological reactions to cannabis smoking. Int. Pharmacopsychiat. 7:249-259, 1972.
- Paton, S., Kessler, R., and Kandel, D. Depressive mood and adolescent illicit drug use: A longitudinal analysis. J. Genet. Psychol. 131:267-289, 1977.
- Paul, M.I. and Carson, I.M. Marihuana and communication. Lancet 2:270-271, 1973.
- Peeke, S.C., Jones, R.T., and Stone, G.C. Effects of practice on marijuana-induced changes in reaction time. Psychopharmacology 48:159-163, 1976.
- Pfefferbaum, A., Darley, C.F., Tinklenberg, J.R., et al. Marijuana and memory intrusions. J. Nerv. Ment. Dis. 165:381-386, 1977.
- Pillard, R.C. Marijuana. N. Engl. J. Med. 283:294-303, 1970.
- Radouco-Thomas, S., Magnan, F., Grove, R.N., et al. Effect of chronic administration of Δ^1 -tetrahydrocannabinol on learning and memory in developing mice, pp. 487-498. In Braude, M.C. and Szara, S. (eds.) Pharmacology of Marihuana. New York: Raven Press, 1976.
- Rafaelsen, L., Christrup, H., Bech, P., and Rafaelsen, O.J. Effects of cannabis and alcohol on psychological tests. Nature 242:117-118, 1973.
- Ray, R., Prabhu, G.G., Mohan, D., et al. The association between chronic cannabis use and cognitive functions. Drug Alcohol Dependence 3:365-368, 1978.
- Rossi, A.M., Kuehnle, J.C., and Mendelson, J.H. Effects of marihuana on reaction time and short-term memory in human volunteers. Pharmacol. Biochem. Behav. 6:73-77, 1977.
- Rossi, A.M., Kuehnle, J.C., and Mendelson, J.H. Marihuana and mood in human volunteers. Pharmacol. Biochem. Behav. 8:447-453, 1978.
- Roth, W.T., Rosenbloom, M.J., Darley, C.F., et al. Marihuana effects on TAT form and content. Psychopharmacologia 43:261-266, 1975.
- Rubin, V. and Comitas, L. Ganja in Jamaica: A Medical Anthropological Study of Chronic Marijuana Use. The Hague: Mouton and Co., 1975.
- Salzman, C., Van Der Kolk, B.A., and Shader, R.I. Marijuana and hostility in a small-group setting. Am. J. Psychiatry 133:1029-1033, 1976.
- Satz, P., Fletcher, J.M., and Sutker, L.S. Neuropsychologic, intellectual, and personality correlates of chronic marijuana use in native Costa Ricans. Ann. N.Y. Acad. Sci. 282:266-306, 1976.
- Schaefer, C.F., Gunn, C.G., and Dubowski, K.M. Dose-related heart-rate, perceptual, and decisional changes in man following marihuana smoking. Percept. Mot. Skills 44:3-16, 1977.
- Schaeffer, J., Andrysiak, T., and Ungerleides, J.T. Cogention and long-term use of gauja (cannabis). Science 213:465-466, 1981.
- Sharma, S. and Moskowitz, H. Effect of marihuana on the visual autokinetic phenomenon. Percept. Mot. Skills 35:891-894, 1972.

- Sharma, S. and Moskowitz, H. Marihuana Dose Study of Vigilance Performance, Proceedings, 81st Annual Conference, American Psychological Association, pp. 1035-1036, 1973.
- Sharma, S. and Moskowitz, H. Effects of two levels of attention demand on vigilance performance under marihuana. Percept. Mot. Skills 38:967-970, 1974.
- Shean, G.D. and Fechtmann, F. Purpose in life scores of student marihuana users. J. Clin. Psychol. 27:112-113, 1971.
- Shilling, D.J. and Stillman, R.C. (moderators). Psychological effects of THC in cancer. Paper session at the 133rd American Psychiatric Association annual meeting, San Francisco, California, May 6, 1980.
- Simon, W.E. Psychological needs, academic achievement and marijuana consumption. J. Clin. Psychol. 30:496-498, 1974.
- Smiley, A., Ziedman, K., and Moskowitz, H. Pharmacokinetics of drug effects on driving performance: driving simulator tests of marihuana alone and in combination with alcohol. Report to the National Institute on Drug Abuse, Contract No. 271-76-3316, 1981.
- Smith, D.E. The acute and chronic toxicity of marijuana. J. Psychadelic Drugs 2:37-47, 1968.
- Smith, D.E. and Mehl, C. An analysis of marijuana toxicity. Clin. Toxicol. 3:101-115, 1970.
- Soueif, M.I. Some determinants of psychological deficits associated with chronic cannabis consumption. Bull. Narc. 28:25-42, 1976.
- Spencer, D.J. Cannabis-induced psychosis. Br. J. Addict. 65:369-372, 1970.
- Stanton, M.D., Mintz, J., and Franklin, R.M. Drug flashbacks. II. Some additional findings. Int. J. Addict. 11:53-69, 1976.
- Sterling-Smith, R.S. Alcohol, marihuana and other drug patterns among pperators involved in fatal motor vehicle accidents, pp. 93-105. In: Israelstam, S. and Lambert, S., eds. Alcohol, Drugs and Traffic Safety. Toronto: Addictive Research Foundation of Ontario, 1975.
- Stiglick, A. and Kalant, H. Learning impairment in the radial-arm maze following prolonged cannabis treatment in rats. Psychopharmacology (in press).
- Stillman, R.C., Wolkowitz, O., Weingartner, H., et al. Marijuana: Differential effects on right and left hemisphere functions in man. Life Sci. 21:1793-1800, 1977.
- Talbott, J.A. and Teague, J.W. Marihuana psychosis: Acute toxic psychosis associated with the use of cannabis derivatives. JAMA 210:299-302, 1969.
- Tart, C.T. Marijuana intoxication: Common experiences. Nature 226:701-704, 1970.
- Tassinari, C.A., Ambrosetto, G., and Gastaut, H. Effects of marijuana and delta-9-THC at high doses in man. Electroencephalogr. Clin. Neurophysiol. 34:760, 1973.
- Taylor, S.P., Vardaris, R.M., Rawtich, A.B., et al. The effects of alcohol and delta-9-tetrahydrocannabinol on human physical aggression. Aggressive Behav. 2:153-161, 1976.

- Teale, J.D., Clough, J.M., King, L.J., et al. The incidence of cannabinoids in fatally injured drivers: An investigation by radioimmunoassay and high pressure liquid chromatography. J. Forens. Sci. Soc. 17:177-183, 1977.
- Tennant, F.S. Drug abuse in the U.S. Army, Europe. JAMA 221:1146-1149, 1972.
- Tennant, F.S. and Groesbeck, C.J. Psychiatric effects of hashish. Arch. Gen. Psychiatry 27:133-136, 1972.
- Thacore, V.R. and Shukla, S.R.P. Cannabis psychosis and paranoid schizophrenia. Arch. Gen. Psychiatry 33:383-386, 1976.
- Thurlow, H.J. On drive state and cannabis: A clinical observation. Can. Psychiatr. Assoc. J. 16:181-182, 1971.
- Tinklenberg, J.R. Marijuana and human aggression, pp. 339-357. In Miller, L.L. (ed.) Marijuana: Effects on Human Behavior. New York: Academic Press, 1974.
- Tinklenberg, J.R., Roth, W.T., and Kopell, B.S. Marijuana and ethanol: Differential effects on time perception, heart rate, and subjective response. Psychopharmacology 49:275-279, 1976a.
- Tinklenberg, J.R., Roth, W.T., Kopell, B.S., and Murphy, P. Cannabis and alcohol effects on assaultiveness in adolescent delinquents. Ann. N.Y. Acad. Sci. 282:85-94, 1976b.
- Treffert, D.A. Marijuana use in schizophrenia: A clear hazard. Am. J. Psychiatry 135:1213-1215, 1978.
- U.S. Department of Health and Human Services, National Institute on Drug Abuse. Project DAWN. Annual Report--1979. Washington, D.C.: U.S. Government Printing Office, 1979.
- Vachon, L., Sulkowski, A., and Rich, E. Marijuana effects on learning, attention and time estimation. Psychopharmacologia 39:1-11, 1974.
- Warren, R., Simpson, H., Cimbura, G., et al. Drug involvement in traffic fatalities in the province of Ontario. Proceedings of the 24th Conference of the American Association for Automotive Medicine (in press).
- Weil, A.T., Zinberg, N.E., and Nelsen, J.M. Clinical and psychological effects of marijuana in man. Science 162:1234-1242, 1968.
- Weil, A.T. Adverse reactions to marijuana, classification and suggested treatment. N. Engl. J. Med. 282:997-1000, 1970.
- Wig, N.N. and Varma, V.K. Patterns of long-term heavy cannabis use in North India and its effects on cognitive functions: A preliminary report. Drug and Alcohol Dependence 2:211-219, 1977.
- Zeidenberg, P., Clark, W.C., Jaffe, J., et al. Effect of oral administration of delta-9-tetrahydrocannabinol on memory, speech, and perception of thermal stimulation: Results with four normal human volunteer subjects. Preliminary report. Compr. Psychiatry 14:549-556, 1973.

J

THERAPEUTIC POTENTIAL AND MEDICAL USES OF MARIJUANA

There has been growing interest in the possibility that cannabis and its derivatives will be valuable for the treatment of several medical and psychiatric conditions. The 97th Congress, for example, introduced a bill (H.R. 4498) "to provide for the therapeutic use of marijuana in situations involving life-threatening or sense-threatening illness and to provide adequate supplies of marijuana for such use."

Most of the putative therapeutic effects of cannabis are believed to be mediated by the central nervous system. These include effects on appetite, nausea and vomiting, epilepsy, muscle spasticity, anxiety, depression, pain, and on glaucoma, asthma, and the symptoms of withdrawal from alcohol and narcotics. The literature on these and other therapeutic actions believed mediated by the central nervous system will be reviewed in this chapter.

In general, the committee finds that cannabis shows promise in some of these areas, although the dose necessary to produce the desired therapeutic effect is often close to one that produces an unacceptable frequency of toxic (undesirable) side-effects. What is perhaps more encouraging than the therapeutic effects observed thus far is that cannabis seems to exert its beneficial effects through mechanisms that differ from those of other available drugs. This raises the possibility that some patients who would not be helped by conventional therapies could be treated effectively with cannabis. A second possibility is that cannabis could be combined with other drugs to achieve a therapeutic goal, but with each drug being used at a lower dose than would be required if either were used alone. As a result, fewer side-effects would be expected to occur. It may be possible to reduce side-effects by synthesizing related molecules that could have a more favorable ratio of desired to undesired actions; this line of investigation should have high priority, because such synthetic derivatives may ultimately have widespread therapeutic use.

GLAUCOMA

Glaucoma is the leading cause of blindness in the United States. The term is used to describe a group of ocular diseases characterized by an increase in intraocular pressure, which damages the optic nerve and leads eventually to loss of vision. The disease affects over two million Americans of age 35 or older. Although there is increasing risk of glaucoma with increasing age, there are forms that develop in infancy. The National Society to Prevent Blindness (1980) also estimates that 300,000 new cases are diagnosed each year.

Treatment of glaucoma depends on the type and cause. It may be pharmacological or surgical. Surgery is useful treatment in relatively few cases; there is a high incidence of failure and serious complications may occur. Available antiglaucoma drugs are effective in regulating intraocular pressure in many patients, and are the mainstay of treatment in the most common form of glaucoma, but there are some adverse side-effects. Some patients are refractory to present forms of treatment and become blind as the disease progresses; for them, there is a particularly urgent need to find effective drugs.

Cannabis (the crude drug), Δ -9-THC (the pure compound), and some other cannabinoid derivatives lower intraocular pressure when administered by various routes, such as inhalation, oral, or intravenous. However, adverse side-effects of cannabis and Δ -9-THC also have been reported. Most patients with glaucoma are elderly, and have a reduced tolerance for many of these side-effects. Even without the adverse side effects, smoking, oral, and intravenous routes of administration are not suitable for the long term. For example, to give adequate control for intraocular pressure, four marijuana cigarettes per day of 2 percent Δ -9-THC would be necessary; this amount is considered heavy usage and could pose a serious health problem in long-term use. Therefore, topical application would be the most salutary route of administration for the patient who needs continuous treatment.

Human Studies

Interest in using cannabis for the treatment of glaucoma was first stimulated by the observation of Hepler et al. (1971, 1972) that intraocular pressure decreased when healthy human subjects smoked cannabis (0.9 percent and 1.5 percent Δ -9-THC content) using an ice-cooled water pipe. (See Green, 1979, for an extensive literature review.)

A study of the acute ocular effects of cannabis in 429 subjects showed there was a dose-related and statistically significant reduction of intraocular pressure following the smoking or ingestion of cannabis containing 1, 2, or 4 percent Δ -9-THC (Hepler et al., 1976a). The amount of pressure decrease was in the range of 30 percent for the cannabis that contained 2 percent Δ -9-THC. Nineteen hospitalized subjects who smoked cannabis of 1 or 2 percent

Δ -9-THC content were observed for 35 days and another 29 subjects were observed as in-patients for a total of 94 days. There was a consistent drop in intraocular pressure in those smoking the 2 percent cannabis and the reduction appeared to last 4 to 5 hours (Hepler et al., 1976a). The authors noted that there did not seem to be much of a cumulative effect on size of pupils or upon intraocular pressure response. Studies by other investigators have confirmed this effect of cannabis and Δ -9-THC in causing reduction of intraocular pressure in humans (Shapiro, 1974; Purnell and Gregg, 1975).

Perez-Reyes et al. in 1976 investigated the effect of intravenous infusion of six cannabinoids in healthy volunteers. Delta-8-THC, Δ -9-THC, 11-hydroxy- Δ -9-THC, cannabinal, cannabidiol, and 8- β -hydroxy- Δ -9-THC were tested on healthy subjects with normal intraocular pressure; Δ -8-THC, Δ -9-THC, and 11-hydroxy-THC caused the greatest reduction in pressure. Of these Δ -8-THC caused the largest decrease in intraocular pressure, with the least number of psychological side-effects.

In a preliminary study of 11 human glaucoma patients who smoked marijuana (1, 2, and 4 percent) or ingested Δ -9-THC (15 mg), intraocular pressure was lowered an average of 30 percent in 7 out of 11 patients (Hepler et al., 1976a). Another study showed that most patients had a decrease in intraocular pressure after ingestion of 15, 20, or 30 mg of Δ -9-THC and after smoking cannabis containing 1, 2, and 4 percent Δ -9-THC (Hepler et al., 1976b).

Ideally, the synthesis of a preparation that could be applied topically to the eye would be most desirable for humans, because this would allow for self-administration. However, initial studies in humans with a topical preparation of Δ -9-THC have not shown a consistent effect on intraocular pressure (Merritt et al., 1981). More work needs to be done on this possibility.

Animal Studies

While animal studies have supported the observation that Δ -9-THC lowers intraocular pressure after oral and topical administration in rabbits (Green et al., 1977a,b; 1978), and after intravenous administration in the cat (Innemees et al., 1979), the reduction in intraocular pressure is not completely understood. It may result in part from a central nervous system effect, and in part through action on the adrenergic system in the eye, probably mediated by the neurotransmitter norepinephrine.

Side-Effects

Marijuana and Δ -9-THC given orally, intravenously, or in cigarettes, to control glaucoma cause systemic side-effects, such as increase in heart rate, decrease in blood pressure, and psychotropic effects. Some of these side-effects are significant enough to pose problems, particularly in patients with glaucoma, who are usually elderly. But

on the other hand, some of these effects may disappear as tolerance (decreased response with repeated use) develops.

Tolerance to the Intraocular Pressure Reducing Effect

No tolerance was detected to the ocular effects of cannabis in rabbits after 1 year's topical instillation of the synthetic cannabinoids SP-1, SP-106, and SP-204 (Green and Kim, 1977; Green et al., 1977b). Hepler et al. (1976b) noted a ceiling effect in humans, in that the smoking of 22 cannabis cigarettes did not result in a significant decrease in eyeball pressure as compared with a subject who smoked only 2 cigarettes. The area of tolerance will need further study, especially if a cannabinoid preparation with a satisfactorily high ratio of therapeutic to side-effects can be found.

Summary

cannabis, Δ -9-THC, other cannabinoid derivatives, and their synthetics, reduce intraocular pressure in humans when smoked, or given intravenously or orally. However, there are systemic side-effects as well as psychotropic effects that are of concern. It is not yet clear whether an effective topical preparation can be developed that will not have these side-effects. Future work should determine whether synthetic cannabinoids or cannabinoid analogues can be found that will be effective in treating glaucoma without causing side-effects.

ANTIEMETIC ACTION

Certain cancer chemotherapeutic agents regularly produce nausea and vomiting after oral or intravenous administration. Those that are most severe in that respect are cisplatin, actinomycin D, adriamycin, cyclophosphamide, methotrexate, and the nitrosoureas. Other anti-cancer compounds may produce nausea less regularly or in less marked fashion.

Because cancer chemotherapy now can produce increased survival in patients with some neoplasms (recurrent or metastatic breast cancer, small cell carcinoma of the lung, ovarian cancer, and others) and substantial cure rates in several (acute lymphoblastic leukemia, Hodgkins disease, germ cell tumors of the testis, etc.) nausea and vomiting that interfere with patients' willingness to continue therapy can be a life-threatening side-effect. Even for those willing to endure the symptoms, they can be extremely unpleasant and debilitating.

Established antiemetics (prochlorperazine and other phenothiazines) are not very effective against drug-induced emesis, and there is a need for new and more reliable antiemetic agents. Metoclopramide, a derivative of procainamide, has recently been shown to be

more effective than prochlorperazine in certain situations and seems promising (Gralla et al., 1981).

The suggestion that cannabis might have some useful antiemetic activity in this setting arose about 1973, when patients receiving intensive chemotherapy for acute leukemia observed that their "social" use of cannabis appeared to reduce their customary nausea and vomiting.

Clinical Investigations

Several controlled studies have been reported. In one of the early ones (Sallan et al., 1975), Δ -9-THC in 15- or 20-mg doses by mouth was compared with a placebo in a randomized double-blind crossover trial in 22 patients whose nausea and vomiting had been shown refractory to other antiemetics. In 14 of 20 courses of treatment, patients obtained "complete or partial relief" with Δ -9-THC; in none of 22 courses did patients report benefit with the placebo. It was observed that the antiemetic effect of Δ -9-THC occurred only in association with the "high," and it was necessary to maintain the "high" in order to maintain the antiemetic effect.

In another controlled trial (Chang et al., 1979), 14 of 15 patients with osteogenic sarcoma treated with high-dose methotrexate had less nausea and vomiting with Δ -9-THC than with the placebo. In that study, patients with other tumors being treated with cytoxan and adriamycin did not respond as well. That report and others like it suggested that the antiemetic effect of Δ -9-THC against those chemotherapeutic agents that are moderate in their emetic potential (e.g., methotrexate) was pronounced, but that Δ -9-THC was less effective against those agents with severe emetic properties. In a similar study (Lucas and Laszlo, 1980), 38 of 53 patients with nausea and vomiting refractory to other antiemetics reported good results with Δ -9-THC. Among the failures were those treated with cisplatin, which has been characterized as one of the most emetic agents used in cancer chemotherapy.

In comparison with prochlorperazine, Δ -9-THC has also been reported to be more effective in preventing nausea and vomiting (Ekert et al., 1979; Sallan et al., 1980).

In a larger study (Frytak et al., 1979), of 116 patients treated with 5-fluouracil and methyl-CCNU, Δ -9-THC was said to be no more effective than prochlorperazine. In that study, in which nearly all patients were older than those in the other reported trials, the majority of patients considered the other side-effects of Δ -9-THC so unpleasant that they preferred either prochlorperazine or the placebo.

Another cannabinoid, a synthetic, nabilone, has been provided to several investigators for evaluation as an antiemetic agent; it has been licensed for use in Canada for treatment of nausea associated with cancer treatment. In the largest clinical study to date (Herman et al., 1979), nabilone was compared with prochlorperazine in a double-blind crossover trial. It was found more effective than

prochlorperazine. The patients in that study preferred nabilone to prochlorperazine; the predominant side-effects were somnolence, dry mouth, and dizziness. Hallucinations occurred in a few patients. Euphoria of the type associated with cannabis was infrequent in that study. However, a study in dogs has revealed previously unrecognized late neurologic effects of nabilone at high doses (Archer et al., 1981). Monkeys and rats did not show similar toxic effects with long-term administration of nabilone (Archer et al., 1981), and further studies will be necessary to clarify the safety of this new agent.

Levonantradol is yet another synthetic cannabinoid, related to Δ -9-THC, which has been shown in preliminary clinical studies to have antiemetic action in patients with refractory chemotherapy-induced emesis (Diasio et al., 1981).

Uncontrolled Use of Δ -9-THC

In response to public and political pressures, the National Cancer Institute, the United States Drug Enforcement Agency, and the Food and Drug Administration have agreed to a program whereby the National Cancer Institute is making Δ -9-THC available through the pharmacies of approximately 500 teaching hospitals and cancer centers to physicians who wish to use Δ -9-THC in treating the nausea and vomiting of patients receiving cancer chemotherapy. This broad, uncontrolled program, in which no data other than the reporting of severe reactions are to be collected, may make it extremely difficult to obtain continuing valid evaluations of the effectiveness of Δ -9-THC in the management of nausea and vomiting due to cancer chemotherapy. Although the extent of use of Δ -9-THC under this program is difficult to evaluate, informal communication with the National Cancer Institute indicates that Δ -9-THC has been supplied in substantial quantities to several hundred hospital pharmacies. The problem is further complicated by the fact that the legislatures of 23 states have authorized the use of cannabis by any physician for the management of nausea and vomiting due to cancer chemotherapy. It is expected that little reliable information will be derived from such use.

Summary

There seems little doubt that Δ -9-THC and other cannabinoids are active against the severe nausea and vomiting produced by cancer chemotherapeutic agents. The extent of this activity, its relation to other antiemetics, and its relation to the other effects of the cannabinoids have not yet been accurately determined.

Cannabis leaf, smoked or eaten, is also antiemetic but its activity has been even less well determined than that of Δ -9-THC. Studies with other synthetic cannabinoids have barely begun and much remains to be learned in this field.

APPETITE STIMULANT

It has been stated by "social" users that the smoking of cannabis increases appetite. On that basis, there have been sporadic attempts to use it in patients with advanced cancer to overcome their customary debilitating weight loss.

In several of the studies in which Δ -9-THC was used as an antiemetic in patients receiving cancer chemotherapy, they were reported to have increased appetite and food intake. At this time, it is not certain whether that increase was due merely to relief of nausea and vomiting or to stimulation of appetite. One comparison of habitual marijuana users and controls matched for age and educational background showed increased caloric intake and weight gain among the users (Greenberg, et al., 1976). Furthermore, a double-blind controlled study (Hollister, 1971) of smokers of cannabis or placebo cigarettes provided with unlimited quantities of a high-caloric beverage indicated an increase in caloric consumption in those using cannabis compared with those using the placebo; however, the variability was very large and there was some question that cannabis could be considered a clinically significant appetite stimulant.

In another study of the psychological effects of Δ -9-THC in patients with advanced cancer, it was observed that Δ -9-THC appeared to stimulate appetite and retard weight loss (Regelson et al., 1976). In that study many patients refused to complete the 2-week trial because of unacceptable side-effects from Δ -9-THC.

The evidence to date suggests that there may be some influence of cannabis on appetite. However, it is not possible to separate that from the effect on nausea and vomiting. Further studies are in progress in cancer patients whose course is not complicated by nausea and vomiting.

ANTICONVULSANT ACTION

A large number of animal studies have been conducted using cannabis as an anticonvulsant. These can be divided into several categories. The first to be discussed will be maximal electroshock seizures (MES)* both in the rat and mouse (Loewe and Goodman, 1947; Sofia et al., 1971; Fujimoto, 1972; Consroe and Man, 1973; Karler et al., 1973; Chesher and Jackson, 1974; Karler et al., 1974; McCaughran et al., 1974; Karler and Turkanis, 1976; Consroe and Wolkin, 1977; Turkanis et al., 1977). In these studies there is a clear dose-response effect in the protection to MES conferred by cannabidiol (CBN) and cannabidiol (CBD). Tolerance to the effect has frequently been reported. However, the tolerance noted with cannabinoids is similar to that seen with phenytoin (DPH). Further, even though tolerance to phenytoin develops with MES, this has not been shown to

*Electrical shock of maximum intensity to cause a major seizure.

be a clinically significant phenomenon. In these studies it is generally found that CBN is less effective against MES and against audiogenic seizures, the latter produced in rodents by loud noise, than CBD. In addition, Turkanis et al. (1977) have emphasized the fact that CBD acts more like DPH than other anticonvulsants and hence would be expected to be effective against major seizures rather than against minor seizures.

There is also extensive literature that CBN and CBD will protect against electrically induced, minimal-(kindling)-seizures (Corcoran et al., 1973; Fried and McIntyre, 1973; Izquierdo et al., 1973; Turkanis et al., 1977, 1979). Reduction of seizures produced by subcortical electrical stimulation in the cat has been reported (Wada et al., 1973). There appears to be much less effect on pentylenetetrazol-induced seizures (Consroe and Man, 1973; Turkanis et al., 1979). Any effect of CBN and CBD on such seizures occurs with maximal toxic doses (Turkanis et al., 1974). Protection against audiogenic seizures (Consroe et al., 1973) and against reflex seizures in the gerbil (Cox et al., 1975) have been reported.

Human studies are largely anecdotal and conflicting. There is one study by Cunha et al. (1980) in which 15 patients suffering from partial complex epilepsy with a temporal focus were randomly divided into two groups. Each patient received, in a double-blind procedure, 200-300 mg of CBD or placebo daily. The drugs were administered for as long as 4 1/2 months. Throughout the study, clinical and laboratory examinations, electroencephalograms, and electrocardiograms were performed at 15- to 30-day intervals. The patients continued their anticonvulsant medications taken before entering the study, on which all them had previously experienced uncontrolled seizures. All patients tolerated CBD well, and there were no signs of toxicity or serious side-effects. Four of the 8 CBD subjects remained nearly free of convulsions during CBD treatment and 3 other patients demonstrated partial improvement in their clinical condition. Cannabidiol was ineffective in 1 patient. The placebo group showed no alteration of seizure frequency. A series of 8 healthy volunteers given CBD showed no effects of the drug.

Summary

There is substantial evidence from animal studies to indicate that cannabinoids are effective in blocking both kindling seizures and MES, and this is particularly true for CBD. MES is a standard testing procedure for evaluation of anticonvulsant drugs. This is strong support for further investigation into the utility of CBD in human epilepsy. The one available carefully controlled human study is in accord with this review.

MUSCLE RELAXANT ACTION

anecdotal reports that cannabis is effective in reducing spasticity. Petro (1980) has reported such a study has been carried out a double-blind study of the effect of cannabis on spasticity (Petro and Ellenberger, 1980). They reported that 10 mg of Δ -9-THC significantly reduced spasticity on clinical measurement and that quadriceps electromyograms showed a decrease in interference pattern in four patients with primarily extensor spasticity. These are preliminary observations, but they suggest that further and more rigorous investigations of the use of cannabinoids in spasticity should be suggested to test their therapeutic effectiveness.

ANTIASTHMATIC EFFECT

Intensive, chronic smoking of concentrated cannabis produces several adverse effects on the airways, including mild bronchoconstriction. But acute smoking of cannabis as well as the ingestion of Δ -9-THC produces bronchodilation in normals and in subjects with chronic, clinically stable bronchial asthma of minimal to moderate severity (Tashkin et al., 1974). These bronchodilator effects were also investigated in individuals in whom an asthmatic attack was induced experimentally by exercise or methacholine (Tashkin et al., 1975). Immediately after the development of bronchospasm, subjects smoked a cigarette containing 500 mg of cannabis assayed at either 1 or 2 percent Δ -9-THC.

Methacholine inhalation promptly caused significant bronchoconstriction (an average decrease in airway conductance of 40-55 percent) and significant hyperinflation (mean increases in thoracic gas volume of 35-43 percent). After placebo smoking or saline inhalation, airway conductance increased only modestly, remaining significantly less than initial control values for 30 to 60 minutes, and thoracic gas volume decreased only gradually, remaining significantly increased for 15 minutes. However, after 2 percent cannabis, and after isoproterenol, there was a prompt return of airway conductance and thoracic gas volume to control values.

Exercise in the asthma-prone individual resulted in average decreases in airway conductance of 30-39 percent and average increase in thoracic gas volume of 25-35 percent. After placebo or saline, there was only a gradual return to control values during 30-60 minutes, but after cannabis, airway conductance and thoracic gas volume returned promptly to preexercise values. Four of the subjects who had previously used cannabis could detect pleasurable sensations after smoking cannabis, which distinguished these effects from those of the placebo cigarette. In that sense these experiments were not strictly blind. The four subjects who had no previous experience with cannabis did not experience any central nervous system effects but did note mild somnolence or light-headedness after cannabis. The results of this study suggest that any bronchial irritant effects of

CORRECTION

**THIS DOCUMENT
HAS BEEN REPHOTOGRAPHED
TO ASSURE LEGIBILITY**

be a clinically significant phenomenon. In these studies it is generally found that CBN is less effective against MES and against audiogenic seizures, the latter produced in rodents by loud noise, than CBD. In addition, Turkanis et al. (1977) have emphasized the fact that CBD acts more like DPH than other anticonvulsants and hence would be expected to be effective against major seizures rather than against minor seizures.

There is also extensive literature that CBN and CBD will protect against electrically induced, minimal-(kindling)-seizures (Corcoran et al., 1973; Fried and McIntyre, 1973; Izquierdo et al., 1973; Turkanis et al., 1977, 1979). Reduction of seizures produced by subcortical electrical stimulation in the cat has been reported (Wada et al., 1973). There appears to be much less effect on pentylenetetrazol-induced seizures (Consroe and Man, 1973; Turkanis et al., 1979). Any effect of CBN and CBD on such seizures occurs with maximal toxic doses (Turkanis et al., 1974). Protection against audiogenic seizures (Consroe et al., 1973) and against reflex seizures in the gerbil (Cox et al., 1975) have been reported.

Human studies are largely anecdotal and conflicting. There is one study by Cunha et al. (1980) in which 15 patients suffering from partial complex epilepsy with a temporal focus were randomly divided into two groups. Each patient received, in a double-blind procedure, 200-300 mg of CBD or placebo daily. The drugs were administered for as long as 4 1/2 months. Throughout the study, clinical and laboratory examinations, electroencephalograms, and electrocardiograms were performed at 15- to 30-day intervals. The patients continued their anticonvulsant medications taken before entering the study, on which all them had previously experienced uncontrolled seizures. All patients tolerated CBD well, and there were no signs of toxicity or serious side-effects. Four of the 8 CBD subjects remained nearly free of convulsions during CBD treatment and 3 other patients demonstrated partial improvement in their clinical condition. Cannabidiol was ineffective in 1 patient. The placebo group showed no alteration of seizure frequency. A series of 8 healthy volunteers given CBD showed no effects of the drug.

Summary

There is substantial evidence from animal studies to indicate that cannabinoids are effective in blocking both kindling seizures and MES, and this is particularly true for CBD. MES is a standard testing procedure for evaluation of anticonvulsant drugs. This is strong support for further investigation into the utility of CBD in human epilepsy. The one available carefully controlled human study is in accord with this review.

MUSCLE RELAXANT ACTION

There are widespread, anecdotal reports that cannabis is effective in relieving muscle spasm or spasticity. Petro (1980) has reported such effects in two cases and has carried out a double-blind study of the administration of Δ -9-THC on spasticity (Petro and Ellenberger, 1981). They reported that 10 mg of Δ -9-THC significantly reduced spasticity by clinical measurement and that quadriceps electromyograms demonstrated a decrease in interference pattern in four patients with primarily extensor spasticity. These are preliminary observations, but they suggest that further and more rigorous investigations of the use of cannabinoids in spasticity should be suggested to test their therapeutic effectiveness.

ANTIASTHMATIC EFFECT

Intensive, chronic smoking of concentrated cannabis produces several adverse effects on the airways, including mild bronchoconstriction. But acute smoking of cannabis as well as the ingestion of Δ -9-THC produces bronchodilation in normals and in subjects with chronic, clinically stable bronchial asthma of minimal to moderate severity (Tashkin et al., 1974). These bronchodilator effects were also investigated in individuals in whom an asthmatic attack was induced experimentally by exercise or methacholine (Tashkin et al., 1975). Immediately after the development of bronchospasm, subjects smoked a cigarette containing 500 mg of cannabis assayed at either 1 or 2 percent Δ -9-THC.

Methacholine inhalation promptly caused significant bronchoconstriction (an average decrease in airway conductance of 40-55 percent) and significant hyperinflation (mean increases in thoracic gas volume of 35-43 percent). After placebo smoking or saline inhalation, airway conductance increased only modestly, remaining significantly less than initial control values for 30 to 60 minutes, and thoracic gas volume decreased only gradually, remaining significantly increased for 15 minutes. However, after 2 percent cannabis, and after isoproterenol, there was a prompt return of airway conductance and thoracic gas volume to control values.

Exercise in the asthma-prone individual resulted in average decreases in airway conductance of 30-39 percent and average increase in thoracic gas volume of 25-35 percent. After placebo or saline, there was only a gradual return to control values during 30-60 minutes, but after cannabis, airway conductance and thoracic gas volume returned promptly to preexercise values. Four of the subjects who had previously used cannabis could detect pleasurable sensations after smoking cannabis, which distinguished these effects from those of the placebo cigarette. In that sense these experiments were not strictly blind. The four subjects who had no previous experience with cannabis did not experience any central nervous system effects but did note mild somolence or light-headedness after cannabis. The results of this study suggest that any bronchial irritant effects of

placebo cannabis smoke were not sufficient to aggravate or perpetuate existing acute bronchospasm to an extent greater than that which might result from the irritant effect of inhaled saline. The results also demonstrate that inhaled Δ -9-THC causes a prompt and complete sustained reversal of methacholine-induced bronchospasm and correction of the associated hyperinflation. These effects were not significantly different from those observed after isoproterenol, although there was a tendency toward a greater degree of bronchial dilation after isoproterenol. Similarly, after inhalation of Δ -9-THC, there was a prompt return of airway conductance and thoracic gas volume during exercise-induced bronchospasm to the preexercise value. After exercise the effects of 10 mg Δ -9-THC was not as efficacious as 1.25 mg isoproterenol.

The way in which Δ -9-THC induces bronchial dilation has not been determined, but previous studies have shown that this effect is not mediated by beta-adrenergic stimulation or inhibition of muscarinic receptors (Shapiro et al., 1973). A vagolytic mechanism is possible, as suggested by other studies carried out on the dog salivary gland (Cavero et al., 1972) and on guinea pig ileum (Gill et al., 1970).

Although ingestion of Δ -9-THC in a sesame oil vehicle has produced bronchodilation in asthmatic patients (Tashkin et al., 1974), less dilation was noted than after smaller doses of Δ -9-THC delivered by smoking. Its significant bronchodilator effect notwithstanding, Δ -9-THC does not appear to be suitable for that therapeutic use, because of its psychotropic effects and possibly other side-effects. However, other cannabinoid compounds such as cannabinal and cannabidiol do not produce the central nervous system effects of tachycardia characteristic of cannabis (Hollister, 1973) and deserve further investigation for possible bronchodilator activity.

ANTI-ANXIETY EFFECT

Users of cannabis have often reported that the drug produces feelings of relaxation and calmness, and some have reported its use to reduce anxiety. A problem with evaluating cannabis as an anti-anxiety drug, however, is that some subjects report increased anxiety or panic after using cannabis (see Chapter 6). For example, Regelson et al. (1976) found less tension and apprehension in cancer patients after cannabis use; but 6 of 50 subjects receiving Δ -9-THC reported such side-effects as severe dizziness, confused thinking, dissociation, and concern over loss of sanity. In normals, Pillard et al. (1974) found no effects of cannabis on experimentally induced anxiety. Nabilone, a synthetic cannabinoid, was found to reduce experimentally induced anxiety in normal volunteers but it was less effective than diazepam (Nakano et al., 1978). Nabilone was found to be more effective than placebo in patients with psychoneurotic anxiety (Fabre et al., 1978).

There are very few studies of cannabis effects on anxiety. There is no indication at this time that cannabis or nabilone are

more effective or reliable than currently available antianxiety medication.

ANTIDEPRESSANT EFFECT

Regelson et al. (1976) reported a significant reduction in self-rated depressive symptoms in cancer patients treated with Δ -9-THC. However, in a carefully controlled trial with four bipolar and four unipolar depressed patients, Kotin et al. (1973) found no anti-depressant activity.

ANALGESIC ACTION

Several animal models have been used to show analgesic effects of cannabis and its analogues (for example, Grunfeld and Edery, 1969; Sofia et al., 1973). Human studies have been conflicting. Milstein et al. (1975) found increase in tolerance to experimentally induced pain after smoking cannabis, while Hill et al. (1974) were unable to detect effects using a different kind of experimental pain. Noyes et al. (1976) found a reduction in pain reports by cancer patients given oral Δ -9-THC; Regelson et al. (1976) also studied cancer patients and found no significant changes in pain after Δ -9-THC.

Those subjects who show analgesic effects of cannabis also show other pharmacological effects such as mental clouding. The literature does not indicate a specific effect of cannabis on pain pathways nor does it suggest that cannabis is likely to be more effective than currently available analgesics.

ALCOHOLISM

Cannabis has been proposed as a treatment for alcoholism (Scher, 1971) based upon case reports and on the observation that cannabis and alcohol were generally not used together. A systematic evaluation (Rosenberg et al., 1978) failed to find cannabis useful in alcoholism. Moreover, recent surveys (see Chapter 2) indicate that currently the abuse of cannabis and alcohol are frequently combined.

OPIATE WITHDRAWAL

Early clinical reports suggested that cannabis might be useful in suppressing the symptoms of opiate withdrawal (Birch, 1889; Thompson and Proctor, 1953). Recently a series of animal studies (Hine et al., 1975a,b; Bhargava, 1976) have found that Δ -9-THC suppresses many of the behavioral manifestations of withdrawal precipitated by naloxone in morphine dependent rodents. This effect is enhanced by cannabidiol (CBD) (Hine et al., 1975a,b), but CBD is not effective alone.

There are no reports of systematic evaluations of cannabis as a treatment of opiate withdrawal in human beings. The animal studies do not present evidence that cannabis is likely to be more effective than currently available treatments for opiate withdrawal.

ANTITUMOR ACTION

There is very little information about the effects of cannabis on neoplasms. In one study (Harris et al., 1976), minor effects were seen on the Lewis Lung Tumor but not in L1210 leukemia. In another study (White et al., 1976), it was found that Δ -9-THC inhibited tumor DNA replication somewhat. In that same study, cannabidiol appeared to have a growth enhancing effect on the Lewis Lung Tumor. These limited studies do not support a view that Δ -9-THC has a useful effect in inhibiting tumor growth.

SUMMARY

Cannabis and its derivatives have shown promise in the treatment of a variety of disorders. The evidence is most impressive in glaucoma, where their mechanism of action appears to be different from the standard drugs; in asthma, where they approach isoproterenol in effectiveness; and in the nausea and vomiting of cancer chemotherapy, where they compare favorably with phenothiazines. Smaller trials have suggested cannabis might also be useful in seizures, spasticity, and other nervous system disorders. Effective doses usually produce psychotropic and cardiovascular effects and can be troublesome, particularly in older patients.

Although marijuana has not been shown unequivocally superior to any existing therapy for any of these conditions, several important aspects of its therapeutic potential should be appreciated. First, its mechanisms of action and its toxicity in several diseases are different from those of drugs now being used to treat those conditions; thus, combined use with other drugs might allow greater therapeutic efficacy without cumulative toxicity. Second, the differences in action suggest new approaches to understanding both the diseases and the drugs used to treat them. Last, there may be an opportunity to synthesize derivatives of marijuana that offer better therapeutic ratios than marijuana itself.

RECOMMENDATIONS FOR RESEARCH

The committee believes that the therapeutic potential of cannabis and its derivatives and synthetic analogues warrants further research along the lines described in this chapter. There also may be significant heuristic benefits to be derived from the study of the biological mechanisms by which these compounds act.