

ALASKA LEGISLATURE COMMITTEE FILES

2007-2008

SHES

124

(DDD) is also a strong protective factor for most secondary disabilities. The services provided by DDD would appear to be both useful and necessary for many clients with FAS/FAE who do not now qualify.

7. Violence against individuals with FAS/FAE occurred at an alarming rate: 72% had experienced physical or sexual abuse, or domestic violence. Being a victim of violence was a strong risk factor for Inappropriate Sexual Behavior, increasing the odds fourfold. Children and adults disabled by FAS/FAE must have better protection and their families may need special training and guidance about Inappropriate Sexual Behavior.
8. Thirty females with FAS/FAE had given birth to a child. Of these, 57% no longer had the child in their care; 40% were drinking during pregnancy; 17% had children diagnosed FAS or FAE; and an additional 13% had children who were suspected by the informants of having FAS/FAE. Special advocacy services for these high risk mothers who themselves have FAS/FAE and special attention to their birth control needs and child care needs should be a top priority.

Recommendations Deriving from the Findings

1. Develop statewide networks of local FAS/FAE Diagnostic Clinics coordinated with local community service providers, to facilitate early diagnosis. This could reduce disrupted schooling, trouble with the law, inappropriate sexual behavior, confinement, alcohol and drug problems, dependent living, and problems with employment, because the organic problems of the child will be recognized from an early age.
2. Develop a coordinated system of parent and citizen education centers, and a system for ongoing in-service training programs for all relevant service providers. These should focus on strategies for improving the quality of life, increasing the duration of stay in each placement, and providing the appropriate life skills and job skills training. This could decrease all secondary disabilities.
3. Develop a state inter-agency network specifically on FAS/FAE including representatives from key state and private agencies and parents groups, to identify an "FAS/FAE Coordinator" within each agency, and to develop methods for detecting, diagnosing, and serving this population. Multiple state agencies, including mental health, schools and special education, the juvenile and criminal justice system, alcohol and drug abuse treatment, and the health department must be involved in this effort. No single agency can be responsive to all the broad based needs of people with FAS/FAE and their families.
4. Fund research on methods to quantify the Central Nervous System impairments associated with FAS/FAE in order to develop clinically-useful diagnostic tools for the neurobehavioral effects of prenatal alcohol exposure. This should facilitate diagnosis of alcohol-affected individuals without the facial dysmorphism and growth deficiency, and permit a speedier response to their service needs before the onset of secondary disabilities.
5. Develop and test new methods (such as those identified in this report) that could be utilized for modifying the eligibility criteria for services from the Division of Developmental Disabilities. Enhancing eligibility for case management, job coaching, and supervised housing should reduce the level of costly secondary disabilities among individuals with FAS/FAE who are unable to live and work independently, but are now unable to get appropriate services because they may not have an IQ below 70 or a full FAS diagnosis.
6. Fund a model long-term residential/job training program for youth and adults with FAS/FAE, and then implement this statewide and nationwide.
7. Mandate the full disclosure of medical/mental health/background history when placing a child in foster or adoptive placements, and provide education and training on FAS/FAE and appropriate support services to families raising such children, including biological families.
8. Prevent future children from being born with FAS/FAE: (1) Expand alcohol/drug inpatient treatment programs for women and their children; (2) Fund a statewide network of Birth to 3 Advocacy programs

for working with the highest risk mothers abusing alcohol/drugs during pregnancy; and (3) Provide advocacy and free long-term birth control options to women with FAS/FAE.

Conclusions

1. People with FAS and FAE have an unacceptable level of secondary disabilities that severely impairs their quality of life and is extremely costly to society.
2. The low level of societal protection and support given to people with FAS and FAE and their families is unacceptable and further compromises their lives. They should be given an appropriate level of societal protection and support. To do this, their primary disabilities must be better understood by families, service providers, and by society at large.
3. The permanent, organic brain damage of people with FAS and FAE is often "hidden" in that it often does not conform to current system guidelines for providing services, such as: a low IQ score, a debilitating physical handicap, serious mental illness, or even an FAS face (and diagnosis).
4. By understanding the devastating secondary disabilities that characterize most individuals with FAS/FAE, and by understanding the intrinsic and extrinsic risk and protective factors that exacerbate or ameliorate these disabilities, we should be able to improve the quality of life for people with FAS and FAE and their families, and to reduce costs to society.
5. The magnitude and cost of these secondary disabilities is huge—when calculated against the estimated figure of 1 to 3 FAS per 1000 births (NIAAA, 1987) and several fold this figure for FAE.

2

Introduction

It is fitting that the International Conference on Overcoming and Preventing Secondary Disabilities in Fetal Alcohol Syndrome and Fetal Alcohol Effects is being held at the University of Washington School of Medicine, September 4-6, 1996. This University has been an important site for fetal alcohol studies for the past 23 years.

In 1973, Kenneth Lyons Jones and David W. Smith, dysmorphologists at the University of Washington Medical School, Department of Pediatrics, identified a "similar pattern of craniofacial, limb, and cardiovascular defects associated with prenatal onset growth deficiency and development" in eight unrelated children of three ethnic groups, all born to chronic alcoholic mothers (Jones, Smith, Ulleland, & Streissguth, 1973). The distinctive pattern of malformations indicated that the damage was of prenatal origin. In a second report, three more infants were identified, and the first necropsy on such a patient "disclosed serious dysmorphogenesis of the brain," which the authors thought might be responsible for some of the functional abnormalities and joint malpositions seen in the syndrome. The naming of this syndrome as Fetal Alcohol Syndrome (FAS) put the emphasis squarely on the presumed etiology and brought international attention to this phenomenon (Jones & Smith, 1973). Among the letters that came to David Smith, was one from Paul Lemoine of Nantes, France, who had published a large study of children born to alcoholic mothers who had similar features and behaviors. Lemoine and colleagues had published their findings in 1968 in a local pediatric journal, but the article had not been translated into English, nor had it appeared in the international medical literature.

From the start, both groups of investigators were fascinated with the unusual behaviors of these children who looked alike (although they were not related) and behaved in a hyperactive and unfocused manner. Through utilization of records from the National Perinatal Collaborative Project (NPCP), Jones and colleagues (1974) were able to demonstrate that 44% of the children of chronic alcoholic mothers identified during pregnancy had "borderline to moderate mental retardation" (defined as an IQ of 79 or below) when examined at 7 years of age. Thirty two percent had enough abnormal features from the physical examination alone to suggest the Fetal Alcohol Syndrome. In a carefully matched comparison group selected from over 50,000 other women and children in the NPCP study, fewer than 10% of the other children had IQ scores below 79, and none had the physical features of FAS.

Clarren and colleagues (1978) presented additional evidence of Central Nervous System (CNS) damage in patients with FAS. "neuropathologic findings on four human neonates exposed to large amounts of ethanol at frequent intervals during gestation... All four brains showed similar malformations stemming from errors in migration of neuronal and glial elements." Two had hydrocephalus and only two of the four were diagnosed as having the Fetal Alcohol Syndrome from external criteria. They concluded: "... the brain alterations may be the only distinct abnormality produced by in utero ethanol exposure." Since 1978 children who had some but not all the features of FAS have been referred to as having Fetal Alcohol Effects (FAE) (Hanson et al., 1978) or Possible Fetal Alcohol Effects (PFAE) (Clarren & Smith, 1978). Recently there have been suggestions that the terminology should be reconsidered (Aase et al., 1995; IOM, 1996).

Within 4 years of the naming of Fetal Alcohol Syndrome, experimental studies of laboratory animals were published demonstrating that alcohol is teratogenic and can produce malformations from prenatal exposure (e.g., Chernoff, 1977; Randall et al., 1977). By 1978, 245 clients with Fetal Alcohol Syndrome had been reported in the medical literature, and FAS was described as the "most frequent known teratogenic cause of mental deficiency in the western world" (Clarren & Smith, 1978). It also was recognized by this time that prenatal alcohol exposure produces a whole spectrum of effects, of which FAS is the most clearly definable.

By the mid 1980's, it was apparent that the physical features of FAS were less unique and characteristic after the onset of adolescence. Clients with this disorder whom we had watched grow up in Seattle had far more life problems than would be expected solely on the basis of their mental retardation or delayed development. Furthermore, as more clients were evaluated, it was clear that there were many children with the full features of FAS who did not technically qualify as mentally retarded, and in many instances, had trouble obtaining appropriate services (Streissguth et al., 1985). It also was clear that many children without the physical manifestations of FAS were also born to alcohol abusing mothers, and that these too had cognitive and adaptive behavior problems similar to children with FAS (Streissguth et al., 1991a).

Efforts to prevent FAS began almost as soon as alcohol was clearly identified as a teratogen, and several public policy decisions were made to enhance efforts to prevent the prenatal effects of alcohol. The Surgeon General of the U.S. (1981) recommended that women not drink alcoholic beverages during pregnancy or when planning a pregnancy. Congress passed legislation in 1989 to mandate warning labels on all alcohol beverage containers sold in the U.S. that included a warning against drinking alcohol during pregnancy.

Understanding the long-term impacts of FAS and FAE on society has taken more time. Understanding the life problems that characterize people with FAS and FAE as they grow and mature is essential to: developing effective interventions, estimating the costs to society, and mandating appropriate public policies.

FAS and FAE have been recognized for the past 20 years as major known causes of developmental disability. Yet, it is only in the past 10 years that the lifelong implications of these disabilities have been recognized. Follow-up studies in four countries have demonstrated the continuing adverse effects of prenatal alcohol exposure into adolescence and adulthood (Aronson, Olegard et al., 1987; Lemoine & Lemoine, 1992; Majewski, 1993; Spohr et al., 1993; 1994; Steinhausen et al., 1993; 1994; Streissguth et al., 1985; 1991).

In 1992, in recognition of this problem, the Centers for Disease Control and Prevention (CDC) through the National Center for Environmental Health, Disabilities Prevention Program, issued a request for proposals. The present research project was undertaken in response to that request. The aim of this project has been to build a prevention information base fundamental to the amelioration of secondary disabilities in clients with FAS and FAE.

For the purposes of this study, primary disabilities are defined as those that reflect the CNS dysfunctions inherent in the FAS or FAE diagnosis. Secondary Disabilities are those that a client is not born with, and that could presumably be ameliorated through better understanding and appropriate interventions.

The study has three main goals:

1. To document the occurrence and range of secondary disabilities that are associated with FAS and FAE.
2. To determine the risk factors associated with these secondary disabilities in order to make recommendations for preventive strategies.
3. To develop a brief Fetal Alcohol Behavior Scale (FABS) so that state/community agencies may identify clients with probable FAS/FAE who may be in need of special services to prevent additional secondary disabilities.

This document is the Final Report of that project.

3.1 Diagnostic Criteria

Fetal Alcohol Syndrome (FAS) is diagnosed when three primary characteristics occur together: growth deficiency, a characteristic pattern of abnormalities primarily observable in the face, and some manifestations of Central Nervous System (CNS) dysfunction. The definition of CNS criteria used here is in keeping with that originally used by Clarren and Smith (1978) and is *not* wholly consistent with the modification suggested by the recent IOM (1996) report for diagnosing FAS. Over the years, a small number of children had been diagnosed PFAS (possible or probable FAS). This term was applied to borderline cases: either the CNS and facial features were classic but the growth was marginal, or the CNS and growth deficiency were classic and the face was "almost" classic. For this report, PFAS and FAS were combined. Fetal Alcohol Effects (FAE) and PFAE (possible or probable FAE) are terms that have been used clinically to apply to individuals who manifest some, but not all of the characteristics of FAS, but were exposed prenatally to significant levels of alcohol. The terms FAE and PFAE, as they have been used by Seattle dysmorphologists since 1974, are consistent with the new diagnostic category of ARND (Alcohol Related Neurodevelopmental Disabilities) suggested by the IOM (1996). For this report, FAE and PFAE were combined.

In the present study, a diagnosis of FAS was assigned to clients who had (1) a clear history of prenatal alcohol exposure; (2) dysmorphic features (primarily observed in the face, such as short palpebral fissures, a pattern of flattened midface, smooth and/or long philtrum, and thin upper lip); (3) growth retardation for height and/or weight below the 10th percentile, and (4) CNS dysfunction (as manifested by microcephaly, developmental delay, hyperactivity, attention and/or memory deficits, learning difficulties, intellectual deficits, motor problems, neurologic signs, and/or seizures). A diagnosis of FAE or PFAE was attributed to those who had a clear history of prenatal alcohol exposure and CNS dysfunction but did not manifest all of the physical features of FAS. All diagnostic evaluations were performed by physicians trained in dysmorphology and genetics.

3.2 Ascertainment

The 661 clients in this study represent a gradually accrued group that began with the first patients diagnosed FAS in 1973 by Jones and Smith and ended with those who came to the University of Washington FAS Diagnostic Clinic between 1993 and 1995. The clients were largely ascertained through clinical referral across a 22-year period, and diagnosed by a small homogeneous group of dysmorphologists who were trained by David W. Smith at the University of Washington Dysmorphology Unit. The sample includes, but is not limited to all those available from the following published studies: 11 clients from the two 1973 Lancet papers describing FAS (Jones et al., 1973; Jones & Smith, 1973); 20 and 17 clients respectively, from the first two FAS follow-up studies (Streissguth, Herman, & Smith, 1978a and b); 8 clients from the 10-year follow-up study of the first 11 clients diagnosed with FAS (Streissguth et al., 1985); Northwest sample only ($n=51$) of the first FAS follow-up study of adolescents and adults (Streissguth et al., 1991a); 40 clients in the test-retest IQ study of adolescents and adults (Streissguth, Randels, & Smith, 1991b); and 24 clients participating in the FAS genotype study (Faustman et al., 1992). The published groups are overlapping, but data for each client are counted only once in the report that follows.

Ascertainment of the 661 clients was through clinical referral for an FAS diagnostic evaluation or through participation in ongoing FAS research projects conducted by the Fetal Alcohol and Drug Unit. Sterling Clarren was the primary diagnostician, diagnosing two-thirds of the clients; David W. Smith, Kenneth L. Jones, and Smith's other dysmorphology fellows diagnosed 7% of the sample. Twenty-four percent of the sample were diagnosed during or before 1990, 15% were diagnosed 1991-1992, and 61% were diagnosed in 1993 or later at the University of Washington FAS Diagnostic Clinic, directed by Sterling Clarren. The clients were primarily from the Pacific Northwest.

The University of Washington FAS Diagnostic Clinic was established with primary funding from CDC in January 1993. Patients are referred to the FAS clinic by physicians, parents, teachers, caseworkers, and other concerned caregivers. Patients are typically referred for cognitive and behavioral problems, and must have a confirmed or suspected history of prenatal alcohol exposure. Only a small number of patients are referred to the clinic based on there being a history

of prenatal alcohol exposure, without major concern for behavioral or cognitive delays. The clinic meets one day per week and can accommodate 4 or 5 patients per day. As many more clients apply than can be seen, priority is given to those whose biologic mothers are living and known to be in the area, in keeping with the goals of the CDC grant that provides basic funding for the FAS clinic (Clarren & Astley, 1997).

3.3 Recruitment

Patients were referred from the FAS Diagnostic Clinic for our Secondary Disabilities Study. Those clients who met diagnostic criteria and who, along with their caregivers, consented to participate were enrolled in the study. The study, which was approved by the University of Washington Human Subjects Review Committee, was explained to the families by our client advocate. Additionally a Confidentiality Certificate was obtained from the Public Health Service to further protect the clients and their families. Three kinds of consents were obtained: (1) for participation in the study, (2) for photographs, and (3) for release of information from schools, hospitals, etc. Clients in our other FAS research projects who met the diagnostic criteria were also notified about the new study and asked to participate. Caregivers were contacted over a four-year period to gather and update information on past and current client status, Secondary Disabilities, and Risk and Protective Factors. Reports of psychological evaluations conducted for research purposes were provided upon request to the clients or their caregivers. In case of a crisis call, clients were referred to appropriate community agencies and professional local services.

3.4 Samples

This final report utilizes two overlapping samples from the 661 clients in this study, one sample for describing the Primary Disabilities ($n=473$) and one for describing the Secondary Disabilities (LHI Sample, $n=115$ —see Table 3.1). All clients were diagnosed either FAS, FAE, PFAS, PFAE, or ARND as described above. Clients from southwest Indian reservations who were included in earlier reports (Streissguth et al., 1988 Manual; Streissguth et al., 1991a; LaDue, Streissguth, & Randels, 1992) are not included in the current report because their recruitment and follow-up were not comparable to those clients in the Northwest. The Primary Disabilities sample includes clients from three years of age and up, the Secondary Disabilities sample includes those 6 years and older. Although the two samples are not entirely nested, the referral source, the diagnosing physician, and the age and date of diagnosis were comparable for the two samples.

Table 3.1 Secondary Disabilities Sample (LHI): enrollment and interview status

Interview status	(n)
Enrolled patients	661
Not eligible for LHI:	
too young (< 6 years old)	82
SW reservation clients	64
enrolled too late/data not processed in time	9
Eligible interviews not obtained:	
family could not be found	52
caretaker refused interview	22
caretaker not available for interview	16
invalid interview	1
Total valid interviews conducted	115

The 91 eligible clients for whom LHIs were not obtained do not differ from the 115 for whom LHIs were obtained in terms of alcohol-related diagnosis, age at diagnosis, frequency of diagnosis prior to 1980, ethnicity, or sex. The loss of these 91 does not seem to present a bias in extent of Primary Disability in that they do not differ significantly from the 115 valid interviews in: IQ (as measured either by mean IQ score or by number of IQ scores at or below 70), achievement (WRAT reading, spelling, arithmetic scaled scores), or adaptive behavior (VABS, Communications, Daily Living Skills, Socialization, and Adaptive Behavior Composite). The interviewed sample appears to be fairly representative of the full set of eligible clients accumulated in this database.

3.5 Primary Disabilities Sample (IQ): Tests Administered

Clients were administered an age-appropriate Wechsler IQ Test: Wechsler Preschool and Primary Scale of Intelligence-Revised (WPPSI-R); Wechsler Intelligence Scale for Children-Revised (WISC-R); and the Wechsler Adult Intelligence Scale-Revised (WAIS-R) (Wechsler, 1967, 1974, 1981). The Wide Range Achievement Test-Revised (WRAT-R) (Jastak & Wilkinson, 1984) was also administered individually to each client, and the Vineland Adaptive Behavior Scale (VABS) (Sparrow et al., 1984) was administered to a caretaker or person who knew the client well, usually at the time the IQ and achievement tests were administered. Testing was carried out primarily at the Fetal Alcohol and Drug Unit (FADU) or in clients' homes or schools.

Every effort was made to schedule testing at a time and location convenient for the client and caregiver, including evenings and road trips to the client's school or home. If clients were unable to be seen for psychological testing, scores from previous IQ and achievement evaluations were used for analysis when available. Reports of psychological evaluations were systematically requested from the schools. All applicable scores were entered into the database and used in analysis. The clients were not asked to come in for psychological evaluations if they had been administered IQ tests within the past year and achievement tests within the past 6 months.

The following psychological test scores were used in the present analyses: From the WPPSI, WISC-R, and WAIS-R: the full scale IQ score, the Verbal Scale IQ score (VIQ), the Performance Scale IQ score (PIQ), and the 11 Subtest Scores were used. WISC-III IQ scores, corrected for comparability with the WISC-R, were used for 11 clients who either could not be seen at our lab for testing or who had been tested at their schools on the WISC-III within the prior year, so were not eligible for retesting. From the WRAT-R: the Standard Scores (SS) for Reading, Spelling, and Arithmetic were used. From the VABS: the Standard Scores from the Adaptive Behavior Composite (ABC) and the Standard Scores for Socialization, Communication, and Daily Living Skills were used.

3.6 Secondary Disabilities Sample: Life History Interview (LHI)

The Life History Interview (LHI) was developed in the third year of the project to evaluate clients of any age and any degree of disability in order to attain maximum coverage of the sample. The focus of the LHI was on the kinds of Secondary Disabilities and Risk and Protective Factors that characterize these clients. The interview was administered by telephone to caretakers/informants of the clients.

The LHI grew out of our experience with deriving secondary disability data from the clinical database that has been accruing during the many years we have been following these clients. Some preliminary secondary disabilities data, coded directly from our clinical records, were presented previously (Streissguth, Kopera-Frye, & Barr, 1994). Higher rates of Secondary Disabilities were obtained from the LHI than from abstracting our clinical database.

The LHI is a comprehensive structured evaluation of ten major areas of possible long-term functional covariates or consequences characteristic of clients diagnosed with FAS/FAE: (1) household and family environment; (2) independent living and financial management; (3) education; (4) employment; (5) physical abuse, sexual abuse and domestic violence; (6) physical, social and sexual development; (7) behavior management and mental health issues; (8) alcohol and drug use; (9) legal status and criminal justice involvement; and (10) companionship and parenting. These areas of concern were explored in terms of past and current client status, secondary disabilities, and possible risk and protective factors.

Interviewers phoned ahead of time to schedule a convenient time for the interviewee and asked to interview the person who knew the client best. Adoptive mothers were the most frequent informants, 33%; followed by biologic mothers, 17%; foster mothers, 12%; stepmothers, 6%; fathers (all types), 7%; legal guardians, 7%; grandmothers, 4%; other relatives, 1%; caseworkers, 0%; spouses or partners, 1%; residential caretakers, 1%; and by others, 2%. A small proportion of the clients, 2%, had no one available who knew them well, so were interviewed themselves for the LHI. Self-report data were only used when they appeared valid in terms of existing data in the clinical database. A total of 415 valid interviews were obtained and coded during a 4-month interval in 1995.

The 37-page LHI contains over 450 separate questions organized to provide the interviewer with clear visual guides for accurate coding. Most questions required the interviewer to code a choice according to a prespecified list of responses. Other questions permitted open-ended responses which were written in verbatim and coded later. Validity ratings by the interviewer follow each section of the LHI. A section was coded "valid" when the responses appeared to be genuine and focused on the questions being posed. A section was coded "questionable" if the informant did not know the client well, seemed guarded, seemed confused and/or contradicted himself/herself, seemed to be biased, seemed hostile, did not understand or speak English very well, or seemed mentally handicapped. All sections with questionable validity codes were reviewed prior to data analysis.

Seven interviewers were trained in administration and coding procedures by the project director, who was regularly available to address queries. Each completed interview was reviewed by the team of interviewers, and coding consensus was reached for any items under contention. Finally, all coded interviews were reviewed by the project director before they were submitted for data entry.

The LHI took, on average, 70 minutes to administer (range, 18 minutes to 3½ hours). Interviews took longer to administer when they involved older clients or those with a greater number of Secondary Disabilities. Interview length was not related to sex, race, or fetal alcohol-related diagnosis of the client.

Most caregivers felt that the LHI covered major areas of the client's functioning, in terms of prevalent problems, possible environmental buffers, and assessment of service needs. Many caregivers stated they felt that the data generated from this interview, once disseminated to the service agencies and community advocates, would help them provide better care for their children with FAS/FAE.

3.7 Quantifying Data Across the Life Span


Data presented in the following chapters were collected across the life span of each client. A given client may have been diagnosed FAS (or FAE) at age 3 years, given an IQ test at age 5 years, and had a caretaker provide information for the LHI when the client was 16 years old. In addition, questions on the LHI assess the onset for many outcomes (age of onset of alcohol/drug problems, age of onset of problems in school). Thus, a given client may contribute data to any of several age groups depending on the topic at hand.

Figure 3.1 presents the cumulative age distributions for ages at which three key evaluation landmarks were made: diagnosis, IQ testing, and LHI. One at a time, follow the dotted horizontal line (at 50%, or median) to each of the three curves and drop a vertical line to the "age" axis. It is seen that the median ages at diagnosis, IQ testing, and LHI are approximately 10, 11, and 14 years, respectively. The dashed line for LHI is to the right of the other lines indicating that this was generally the most recent data collected.

The eight Secondary Disabilities are defined by any history of the secondary disability since birth. Thus, a comment that 59% of clients 21 years and older were reported as having a history of Disrupted School Experience does not mean that problems in school happened after age 21 years; the disruptions may have occurred at much younger ages. Ages at interview correspond to age covered by the history of secondary disability as follows:

- History of secondary disability for clients 21-51 years old at interview covers ages: birth-age at LHI (birth to 51 years)
- History of secondary disability for clients 12-20 years old at interview covers ages: birth-age at LHI (birth to 20 years)
- History of secondary disability for clients 6-11 years old at interview covers ages: birth-age at LHI (birth to 11 years)

3.1 Client ages at evaluation landmarks (LHI sample, n=415)



3.8 Strengths and Limitations of Study Design and Analysis

The strengths of this study design are: It attempts complete coverage of a large and established roster of clients, recruited in a similar manner and examined in a uniform setting. There is no apparent bias in data collection or cooperation. The clients span a broad spectrum of ages, socio-economic backgrounds, and rearing conditions. The instrument (the Life History Interview—LHI) in the hands of a skilled interviewer, seems robust over a huge range of informants and across the full age range of clients, 6 through 51 years, providing extensive coverage of life experience. The LHI has a rich and redundant coding scheme covering many different kinds of behaviors and difficulties.

The limitations of the study design are: It does not involve a representative sample of any defined population. It contains no measures of dose except that all clients were born to mothers who abused alcohol. The diagnostic categorization, although carried out by a small number of dysmorphologists with the same training, is subject to temporal changes in interpretation and referral patterns. Effective utilization of the LHI requires gifted interviewers, trained by those with extensive experience with FAS across the lifespan, and the prior establishment of trust and rapport with families. Nevertheless, the LHI data are only as valid as the knowledge and capabilities of the informants. Obviously there was no random assignment of clients, caretakers, or interventions.

The strengths of the data analysis are: It represents an effective distillation of the vicissitudes of hundreds of lives into a small number of major channels of secondary disability. The sample is large enough to subset informatively in many different and interesting ways. Statistical analysis easily translates into English sentences. Findings are strong and plausible in respect of aspects of the client and also of the client's environment. Many risk/protective factors common to most secondary disabilities have been detected. The statistics are quite simple; the graphics are powerful. The findings lead easily to policy implications and recommendations.

The limitations of the data analysis are: There is no real possibility of assessing causation. The secondary disabilities are intercorrelated as are the risk/protective factors. There is no apparent concern for hypothesis-testing of the academic flavor. The absence of intervention research on this population precludes assessment of the causal role of interventions.

4

Client Characteristics

Two samples are described in this report: The Primary Disabilities Sample (determined by the 473 clients on which a valid IQ test was obtained) and the Secondary Disabilities Sample (determined by the 415 clients on whom a Life History Interview (LHI) was obtained). The two samples are comparable demographically with respect to sex, ethnicity, and diagnosis. Each sample is approximately one-third FAS and two-thirds FAE. The Secondary Disabilities Sample excludes clients below age 6, but includes more clients who are over 21 years.

The two samples have 378 clients in common. Thirty-seven individuals have an LHI but no IQ exam, and 95 have an IQ exam but no LHI. As IQ data from past examinations on our unit had to be used for old clients who could not be brought in for examination, 73 of the 473 IQ scores are evaluations conducted when the clients were 3-6 years of age.

Table 4.1 presents demographic characteristics of the Secondary Disabilities Sample.

Table 4.1. Demographic Characteristics of the Secondary Disabilities (LHI) Sample (N=415).

Demographic Characteristics	N	(%)
SEX		
Male	236	(57%)
Female	179	(43%)
ETHNICITY		
White	248	(60%)
Native American	103	(25%)
Black	30	(7%)
Hispanic	27	(6%)
Asian	5	(1%)
Other	2	(<1%)
PATIENT AGE AT TIME OF INTERVIEW*		
6-11 Years	162	(39%)
12-20 Years	163	(39%)
Over 21 Years	90	(22%)

The 415 clients in the Secondary Disabilities Sample ranged in age from 6 to 51 years at the time the interview was administered to their caretakers or informants. Overall, their median age was 14.2 years. The data in this report are divided into three age groups as shown below. Table 4.2 shows the breakdown of these age groups by sex and diagnosis, Table 4.3 by age at diagnosis and year of diagnosis.

Table 4.2 The Secondary Disabilities (LHI) Sample (n=415) by sex, diagnosis, and age at interview

	6-11 years		12-20 years		21-51 years		Total	
Median age	8.7 years		15.8 years		25.7 years		14.2 years	
(n)=	(162)		(163)		(90)		(415)	

Diagnosis	Male		Female		Male		Female	
	Male	Female	Male	Female	Male	Female	Male	Female
EAS	24	22	34	22	26	27	84	71
EAE	67	49	68	39	17	20	152	108
Total	91	71	102	61	43	47	236	179

Table 4.3. The Secondary Disabilities (LHI) Sample by age of diagnosis, year of diagnosis, and age at interview

Age at LHI	(n)	6-11 years N=162	12-20 years N=163	21-51 years N=90
Age at Diagnosis				
Birth to 5	(94)	65 (40%)	12 (7%)	17 (19%)
6-11	(145)	17 (10%)	41 (27%)	4 (4%)
12-20	(150)	0 (0%)	107 (82%)	25 (20%)
21 and over	(66)	0 (0%)	0 (0%)	66 (81%)
Year of Diagnosis				
1970-1975	(10)	0 (0%)	3 (2%)	13 (17%)
1976-1980	(43)	0 (0%)	5 (3%)	10 (14%)
1981-1985	(48)	2 (1%)	3 (2%)	11 (12%)
1986-1990	(79)	15 (9%)	12 (13%)	12 (28%)
1991-1992	(62)	25 (16%)	27 (17%)	10 (11%)
1993-1997	(253)	32 (17%)	111 (62%)	52 (30%)

Table 4.4 shows the current living situation of these 415 clients by age at interview. About half of these clients lived with biologic or adoptive parents at the time the interview was conducted.


Table 4.4. Current Living Situation of 415 Clients with FAS/FAE at time of interview

Living Situation	TOTAL	Client age at interview:					
		6-11 years		12-20 years		21 years and over	
		FAS	FAE	FAS	FAE	FAS	FAE
Biological Parents	9	2	1	1	3	1	1
Biological Mother	51	4	13	6	19	2	7
Biological Father	34	3	14	3	10	3	1
Adoptive Relatives	10	1	4	1	4	0	0
Non-Adoptive Relatives	22	9	4	3	3	0	3
Adoptive Parents	94	12	39	16	20	5	2
Foster Parents	55	8	23	9	12	3	0
Legal Guardian	25	7	11	2	4	1	0
Friends	7	0	0	1	5	0	1
Spouse only	5	0	0	0	0	3	2
Spouse and Children	3	0	0	0	0	2	1
Partner Only	9	0	0	1	3	2	3
Partner and Children	3	0	0	0	1	2	0
Single Parent with Children	5	0	0	0	0	2	3
Live-in Work	1	0	0	0	1	0	0
Armed Forces	2	0	0	1	0	1	0
Alone	19	0	0	0	0	10	9
With or Near Attendant	1	0	0	0	0	1	0
Rent Subsidy Program	1	0	0	0	0	1	0
Assisted Living	2	0	0	0	0	2	0
Shared Living	6	0	0	0	2	3	1
Therapeutic Foster Home	8	0	2	0	5	1	0
Group Home	6	0	2	0	1	3	0
Adult Family Home	4	0	0	1	1	1	1
Residence School	1	0	0	1	0	0	0
Residence Treatment	8	0	2	3	1	1	1
Residence Corrections	0	0	0	1	4	1	0
Juvenile Corrections	4	0	0	3	1	0	0
Unstable or Homeless	8	0	0	2	4	2	0
Don't Know	0	0	1	1	3	0	1

The general level of intellectual functioning for the clients in the LHI sample is in the "borderline" range, their median IQ is 86. The group's average reading and spelling levels are consistent with their intellectual level: median WRAT-R Reading SS, 82; median Spelling SS, 80. But Arithmetic is lower (median WRAT-R Arithmetic SS, 75). The median for the Vineland Adaptive Behavior Scales (VABS) composite score is 62. The averages for the three summary subdomain scores in Communication, Daily Living Skills, and Socialization are 67, 69, and 66 respectively.

Figure 4.1 shows three relevant ages of events for the LHI sample: the ages at which the clients were diagnosed; the ages at which the IQ test used in this report was administered; and the ages of the clients when the interview was conducted with their caretakers.

4.1 Percent of clients getting diagnosed, tested and the LHI within age ranges, n=415



5


Primary Disabilities

General intelligence, mastery of reading, spelling, arithmetic, and general level of adaptive functioning are considered here as measures of "primary disabilities," representing the CNS manifestations of FAS, the birth defect with which these clients were born. Most of these clients have been raised in environments in which they had the opportunities to learn basic academic skills and adaptive behaviors. Rather than trying to "adjust" such data for adverse early environmental circumstances, we evaluate the effects of such circumstances in the section on risk and protective factors.

IQ data were available on 473 clients in the Primary Disabilities Sample: 178 have a diagnosis of FAS and 295 a diagnosis of FAE. Clients with FAS have a mean IQ score of 79, with a range from 29 to 120. Clients with FAE have a mean IQ of 90, with a range from 12 to 142. Twenty-seven percent of the clients with FAS and 9% of the clients with FAE have an IQ of 70 or below.

Figure 5.1 depicts the frequency distribution of the two sets of IQ scores superimposed on the normative bell curve of IQ. Both the FAS sample and the FAE sample deviate to the low side of the standard IQ distribution. The 11-point IQ discrepancy between clients with the full FAS and those with FAE replicates one we have previously reported (Streissguth et al., 1991a).

5.1 IQ distributions in the Primary Disabilities Sample: FAS and FAE



The Wechsler IQ tests contain 11 subtests, which are often interpreted clinically as a profile. We "centered" those subtest profiles by subtracting the client's mean subtest score from each of the 11 raw scores, to derive a set of 11 centered scores varying about zero as shown in Figure 5.2.

5.2 Profile of IQ subtests: FAS and FAE





Figure 5.2 shows the group means of the "centered" subtest scores for clients with FAS versus FAE. Three subtests on which the FAS/FAE clients do poorly relative to their other subtest scores are Arithmetic, Digit Span and Information. This is a similar pattern to that seen in learning disabled children whose arithmetic, coding or vocabulary, information, and digit span are often the lowest subtests (Vargo, Grosser, & Spalford, 1995). Two subtests on which the FAS/FAE clients do best relative to their other subtest scores are Object Assembly and Picture Completion. This specific pattern of subtest strengths and weaknesses is similar to that associated with prenatal alcohol exposure in the 500 subjects in the Seattle Longitudinal Prospective Study on Alcohol and Pregnancy (Streissguth et al., 1993, Figure 7.1, page 159).

A "strength of profile" score was calculated for each client which reflects the degree to which an individual's subtest profile parallels the group subtest profile shown in Figure 5.2. This "strength of profile" score appears as item 18 on Table 6.1 of the most important Risk and Protective Factors. A high score equals high congruence, a low score, little congruence. This pattern profile covariance is computed as a weighted sum of 11 centered subtest scores. The weight for each score is the sample mean for that score. The "strength of profile" score is not correlated with IQ or age.


We compared academic achievement to what might be anticipated from overall IQ level. Figure 5.3 shows that clients with FAS are functioning close to their IQ level for Reading and Spelling, but are two-thirds of a standard deviation low in Arithmetic. For FAE, average Arithmetic scores are nearly a full standard deviation lower than would be expected from IQ levels; the other two achievement scores are relatively lower as well. These data replicate the findings of the previous study restricted to adolescents and adults (Streissguth et al., 1991a).

Figure 5.3 shows that both groups of clients have low adaptive behavior scores relative to their respective IQ levels. The relative deficit in these scores, in fact, is considerably greater than shortfalls in achievement, regardless of diagnosis. Figure 5.4 clarifies all these comparisons in a different graphic style.

5.3 IQ, WRAT, VABS: FAS and FAE



5.4 Relationship of achievement and adaptive behavior to IQ, by diagnosis




In Figure 5.4, each mean score for the FAS group has been connected by line to the corresponding mean score for the FAE group. The darkest line (at the top) is IQ, which is higher for both FAS and FAE clients than are the Achievement and Adaptive Behavior scores for this same group of clients.

The lines for Arithmetic and Communications are approximately parallel to the IQ line. This indicates that although Arithmetic and Communications are lower than IQ, the discrepancy is about the same for FAS clients as it is for clients with FAE.

The five darker lines (Reading, Spelling, Daily Living Skills, Socialization, and the Adaptive Behavior Composite) are relatively flatter. This indicates that although all 5 mean scores are lower than IQ within their respective samples, the discrepancy appears to be greater for FAE clients than for FAS clients.

Figure 5.5 presents data from only the Secondary Disabilities (LHD) Sample on which IQ was available (n= 378). This figure shows that the IQ scores for the clients at each LHD age grouping were comparable for subgroups defined by sex and diagnosis. The distribution of IQ scores is presented with a boxplot for each of the 12 groups: age (3) by sex (2) by diagnosis (2).

5.5 LHD Sample: IQ distributions by sex, diagnosis, and age at interview




The box in a boxplot spans the middle half of the data from 25th to 75th percentiles; the dark horizontal line within a box indicates the median IQ score; the whiskers extending vertically from each box reach to the lowest and highest IQ; the open boxes on lower and upper whiskers indicate the 10th and 90th IQ percentiles respectively.

Figure 5.6 describes the discrepancies between PIQ (Performance IQ) scores and VIQ (Verbal IQ) scores, and how these discrepancies for the FAS/FAE clients differ from the normal population. Notice that more individuals fall outside the dotted line at the top of the figure compared to the number outside of the lower dotted line. This indicates that more FAS/FAE individuals have a PIQ greater than VIQ by 20 or more points, compared to the converse. Actually 8% of the clients fall above the line compared to 2.4% below the line. According to Sattler (1988), only 5% of a population would be expected to have a PIQ greater than VIQ by this magnitude.

Notice also the two heavy dashed curves illustrating that for individuals with IQ's up to about 100, clients were more likely to deviate in the direction of PIQ greater than VIQ rather than the converse.

The large black triangle represents the mean scores of all individuals with FAS. Here we see that the mean PIQ is 84 while the mean VIQ is 79 for an average discrepancy of 5 points. The large white triangle represents the mean scores of all individuals with FAE. There is also a 5 point discrepancy between the mean PIQ of 93 and the mean VIQ of 88.

5.6 The two components of IQ by diagnosis





Risk and Protective Factors: Overview & Definitions

Our main goal in this research project was to study the prevalence in this sample of the deleterious life outcomes we have called Secondary Disabilities. These are examined in Chapters 7 through 17. Of course, risks of these outcomes are not the same for all clients. In this Chapter, we describe our rationale and methodology for quantifying these risks. Table 6.1 contains our final list of 21 Risk and Protective Factors and their definitions; Figure 6.1 depicts the relationship of these 21 Risk and Protective Factors to our six main Secondary Disabilities, and Figure 6.2 displays the 8 by 8 matrix of all eight Secondary Disabilities against each other. Of course, we cannot maintain that these relationships are causal or independent.

Definition of the Risk and Protective Factors

We considered two types of Risk and Protective Factors: Intrinsic and Extrinsic. *Intrinsic* Factors include attributes of the client that directly characterize his/her biology or extent of putative brain damage and include age, sex, diagnosis, IQ, IQ subtest profile, and ABC score. *Extrinsic* Factors represent environmental influences that have the potential of affecting life outcomes; these can be gleaned from the Life History Interview (LHI). Extrinsic Risk and Protective Factors were subjectively evolved from clinical experience, for example, fraction of life in a stable and nurturant home, living in non-alcoholic families, not being abused or neglected, receiving special help as needed. Some of the Risk and Protective Factors were available from the test and demographic data in our database, the rest were developed into questions on the LHI, dispersed appropriately within the various sections of the interview. Variables like "fraction of life" came from questions like: "During what ages did client live in a stable and nurturant household?" The quality of "nurturance" is subjective, but the fraction of life is intended as an explicit quantification. Typically the informant would think back over the history of living situations and list for the interviewer the inclusive periods during which the situation was judged "nurturant"; hence, the dates are probably fairly accurate. (Not all Risk and Protective Factors fit under intrinsic or extrinsic, e.g. the FABS is neither.)

After administering 315 LHIs, we developed a scoring system that included two types of scores. Some were median splits, in which the top half of the distribution was compared with the lower half on a given item, such as "fraction of life lived with alcohol and drug abusers"; others were simple thresholds—binary scores that applied only to a proportion of the subjects, such as "violence against client." For some scores like this, with broader coverage, separate components (domestic violence, physical abuse, and sexual abuse) were also examined individually as risk factors. A likely group of 35 such candidates was examined in relation to the Secondary Disabilities discussed in Chapter 7.

After examination of the data, we retained the set of 21 Risk and Protective Factors depicted in Table 6.1, which presents the definition of each, the type of score obtained, distributions for all dichotomies, and the source of the information from which the data derive. For items deriving from the LHI, the specific item numbers are documented in Table 6.1.

Table 6.1. Definitions and derivations of risk/protective factors as reported by the caretaker, physician and psychometrist with distribution for clients 12 years and over.

Definition	Type	Derivation of binary protective/ risk factors**	%**	Source of info
1. Stable and nurturant household	E	Median split on fraction of life: P: 72 - 100% of life R: 0 - 71% of life	49% 51%	LHI
2. Age of diagnosis of FAS/FAE	E	Age at diagnosis split at 6 years P: under 6 years R: 6 years and over	11% 89%	Dysmorphology Exam
3. Violence against client	E	Yes/no if ever in life patient experienced domestic violence, sexual abuse, and/or other physical abuse: P: no violence experienced R: violence was experienced	28% 72%	LHI
4. Average number of years per living situation till age 18	E	Median split on number of years HH: P: 2.8 - 18.0 yrs per household R: <2.8 yrs per household	62% 38%	LHI
5. Quality of home environment during during later childhood (8-12 years)	E	Median split on a count of positive qualities of home: P: 10 - 12 good qualities R: 0 - 9 good qualities	65% 35%	LHI
6. DDID eligibility status	E	Indicator of whether the family both applied for and was eligible for DDID: P: applied for and was eligible R: applied for, but was not eligible	56% 44%	LHI
7. Living with parental drug abusers	E	Median split on fraction of life the proband has lived with alcohol drug abusers: P: 0 - 29% of life R: 30 - 100% of life	51% 49%	LHI
8. Alcohol related diagnosis	E	Indicator of FAS rather than FAE: P: FAS or PEAS R: FAE, PEAF, ARND	45% 55%	General Information Form
9. Basic needs are met	E	Median split on fraction of life that basic needs are met: P: 0 - 12% of life R: 13 - 100% of life	40% 60%	LHI
10. Fetal Alcohol Behaviors Scale (FABS)	E	Median split on number of fetal alcohol behaviors according to caretaker: P: 0 - 20 of 30 behaviors endorsed R: 21 - 30 of 30 behaviors endorsed	66% 34%	Personal Behavior Checklist
11. Intelligence (measured IQ)	E	all < 70 P: 70 - 75 R: IQ > 75	66% 34%	FABII (parent) WISC-III (school) FABS (parent)

Definition	Type	*Derivation of binary protective/ risk factors**	%***	Source of info (LHI items)
Table Continues				
12. Driver's license attainment (≥16 years of age only)	E	Whether or not patient ≥16 years has has a driver's license: P: yes R: no	31% 69%	LHI
13. Sex	I	Sex: P: female R: male	43% 57%	General Information Form
14. VABS ABC Standard Score	I	Median split on the Adaptive Behavior Composite Standard Score P: VABS ABCss ≥ 65 R: VABS ABCss < 65	58% 62%	VABS Interview (FADU)
15. Difference between IQ and ABCs	I	Comparison of IQ & VABS ABCss P: IQ & ABCss are within 15 pts R: more than 15 pts different	32% 68%	Testing & interviewing (FADU)
16. Has there always been at least one caretaker after age 18 years	E	Yes/No P: never after age 18 years R: sometimes—always	64% 36%	LHI
17. Difference between performance and verbal IQ	I	Performance IQ (PIQ) minus verbal IQ (VIQ) P: PIQ more than 15 points higher R: PIQ not more than 15 points higher	17% 83%	FADU testing
18. Similarity of IQ subtest profile to other FAS/FAE patients	I	Median split on the IQ subtest strength of profile score P: weak IQ profile (not similar) R: strong IQ profile (similar)	49% 51%	FADU testing
19. SSI eligibility status	E	Indicator of whether the patient both applied for and was eligible for SSI P: applied for, but was ineligible R: applied for and was eligible	28% 72%	LHI
20. Age of Secondary Disability assessment	E	Status as adult (≥21 years) P: under 21 years old R: 21 years and older	64% 36%	from dates
21. Always 2 caretakers available before 18 yrs	E	yes/no P: not always 2 caretakers R: always 2 caretakers available	57% 43%	LHI

* E = Extrinsic, I = Intrinsic

**Item polarities are assigned according to the median plotted in figure 6.1, which pertains *only* to the first six secondary Disabilities. For Risk and Protective Factors for the last two Secondary Disabilities (Dependent Living and Problems with Employment) see Figures 14.2 and 15.2, respectively.

P = protective end; R = risk end

***Percents apply only to clients 12 years and older

Odds Ratios

Through most of this report, we will express effects of Risk and Protective Factors in terms of *odds ratios*. (Some readers may recognize this approach as a series of univariate logistic regressions.) An example will make the computation clear. Suppose we are interested in the role of low IQ (less than or equal to 70) as a predictor of some binary outcome, in this case, "trouble with the law," one of the summary Secondary Disability scores introduced in the next chapter. The subsample to which this outcome score seems most pertinent is the set of clients 12 and older. For this subsample, the 2 x 2 table is as follows:

		Trouble	
		no	yes
IQ	>70	72	122
	≤70	20	18

The odds of no Trouble for IQ>70 are 72/122. The odds for IQ≤70 are 20/18. The *odds ratio* for the effect of the IQ dichotomy on this outcome is thus $(20/18)/(72/122)$, or 1.88, about a 2:1 *protective effect*. The clients of lower IQ have *relatively fewer* problems with the law; that is, they do better. For a different cut of "low IQ" at 85, the table is:

41	88
51	52

The odds ratio $(51/52)/(41/88)$, or 2.10, stands for just about the same "protective effect." Ratios less than 1 indicate "risk" rather than protection. The method of multiple logistic regression would attempt to "adjust" all such odds ratios for associations among the separate factors. We have no reason to believe that any of the assumptions of these analyses are met, and so we do not pursue this refinement here.

For each Secondary Disability in the subsequent chapters, each risk or protective factor is associated with an odds ratio of this sort. For IQ ≤ 70, for instance, the ratios are, for Mental Health Problems, $(2/36)/(9/187)$ or 1.15 (clearly a problem of small cell counts here), for Disrupted School Experience, $(19/17)/(73/122)$, or 1.87, for Confinement, $(22/16)/(95/100)$, or 1.45, for Inappropriate Sexual Behavior, $(17/21)/(99/92)$, or 0.75 (a slight escalation of risk), for Alcohol/Drug Problems, $(28/9)/(123/73)$, or 1.85.


As a sorted list, these odds ratios are (.75, 1.15, 1.45, 1.85, 1.87, 1.88) with median $(1.45 + 1.85)/2 = 1.65$. Thus, "IQ ≤ 70" is plotted at height 1.65 in Figure 6.1 with the separate odds ratios around it. Each later chapter begins with a graph of similar information as a "profile" across all the risk or protective factors as they pertain to that outcome, one at a time.

The entire set of odds ratios for these 21 Risk and Protective Factors is presented in Figure 6.1. The horizontal scale in this figure is a log scale, so that reciprocals (e.g., 3 and 1/3) are at equal distances from the centerline in opposite directions. These mirrored points represent equivalent odds ratios of either risk (left) or protection (right). The "Protective" Factors, which lie in the top half of the page are the converse of the "Risk" Factors, which lie in the bottom half of the page. Each row on the figure represents a single Risk/Protective Factor. The one at the top of the page, "percent of life in a stable and nurturant household," is the strongest Protective Factor, on average, against the first six Secondary Disabilities described in Chapter 7. The median impact of this one Protective Factor on the six Secondary Disabilities is represented by a black diamond. Risk/Protective Factors that have the least impact (odds ratios closest to 1.0, whether risky or protective) on these six Secondary Disabilities appear at the midpoint of the page, at the end of the list of 21. Arrows indicate scores that could not be calculated for one of the diamonds because there was a zero in either the numerator or the denominator of the odds ratio. The vertical column of decimal numbers to the left of the Risk/Protective Factor names on Figure 6.1 is the median of the six odds ratios observed for that Factor.

The white diamonds on each row represent the relationships of that Risk/Protective Factor to each of the six Secondary Disabilities studied in Chapters 8 through 13. Each of these chapters has a graph with one set of diamonds, reflecting how the 21 Risk/Protective Factors are related to the Secondary Disability discussed in that chapter. For simplification, these graphs only depict the Protective side; one can imagine the Risk side to be the reflection. These separate graphs appear in each chapter as its second figure. Each white diamond found on a graph can be found at exactly the same spot in Figure 6.1. The ordering of Risk/Protective Factors have been retained across chapters. The Risk and Protective Factors are ordered by median odds ratio seen in Figure 6.1, from most protective at the top through most risky at the bottom.

In Figure 6.1, two types of Risk and Protective Factors can be detected. Recall that the vertical line at the center differentiates the Risk side of the distribution (at the left) from the Protective side (right). Any Factor for which all the open diamonds are to the same side of center line is a "Universal" Factor. The Universal Protective Factors are those that are consistently protective for each of the six Secondary Disabilities. Except as noted in Table 6.1, the sample for each Risk and Protective Factor was divided at the median, which is the number appearing below. These Universal Protective Factors include: living in a nurturant and stable home at least 72% of life; being diagnosed before the age of 6 years; never having experienced violence; getting to stay longer than an average of 2.8 years in each household; experiencing a good quality of home (10 of 12 "good" qualities); being FAS (rather than FAE); and having basic needs met at least 13% of life. Of course, many of these items overlap substantially among themselves; we are making no claim that any of these Risk or Protective Factors are independent of others. Other Risk/Protective Factors are called "Specific" because they are Protective Factors for some Secondary Disabilities yet may be Risk Factors for others.

6.1 Risk/Protective Factors across 6 Secondary Disabilities among clients 12 years and older, max n = 253



7

Secondary Disabilities: Definitions & Overview

Secondary disabilities are those that a client is not born with, and that could presumably be ameliorated (either fully or partially) through better understanding and appropriate interventions. The project examined altogether eight Secondary Disabilities. The first six pertain to clients of all ages and are easily quantifiable. The latter two, Dependent Living (DPL) and Problems with Employment (PWE), pertain only to adult clients (21 and over), restricting the sample size to 90 compared to 415 for the other Secondary Disabilities. DPL and PWE are by nature more subjective and required more complex and qualitative definitions. In the graphs that follow in this chapter, the two latter Secondary Disabilities are set apart from the others to draw attention to this difference. All data on Secondary Disabilities derive from the Life History interview (LHI). This means that the accuracy of the information is dependent on the knowledge of the informant, which varied depending on the age and living circumstances of the client.

Figure 7.1 shows the prevalence of all eight Secondary Disabilities across the entire sample of 408-415 clients. The remaining figures in this chapter focus primarily on clients 12 years of age and older, the age at which most of these Secondary Disabilities were manifest.

7.1 Prevalence of Secondary Disabilities across the Life Span



Secondary Disabilities Definitions

Mental Health Problems (MHP) was coded "yes" if the client was described as (a) ever having any of a list of mental health problems, or (b) gone to a psychiatrist, psychotherapist, or counselor for mental health problems, or (c) ever having been a client in a psychiatric or a mental health hospital (Chapter 8)

Disrupted School Experience (DSE) was coded "yes" if the client was described as ever being suspended or expelled from school, or as having dropped out of school (Chapter 9)

Trouble With the Law (TWL) was coded "yes" if the client was described as having been charged or convicted or in trouble with authorities for any one or more of these main categories of criminal behavior: crimes against persons (excluding sex-related), crimes against property, possession or selling of illegal substances, sex crimes, driving violations, parole violations, skipping bail, or escape, or other crimes (primarily victimless, such as running away from home) (Chapter 10)

Confinement (CNF) was coded "yes" if the client was described as ever having been incarcerated for a crime or having received inpatient treatment for mental health problems or for alcohol or drug abuse problems. (Chapter 11)

Inappropriate Sexual Behavior (ISB) was coded "yes" if the client had ever been sentenced to sexual offenders treatment or had ever been reported as "repeatedly" having one or more problems with sexuality. (Chapter 12)

Alcohol and/or Drug Problems (ADP) was coded "yes" if the client was reported to have had alcohol abuse problems, drug abuse problems, and/or alcohol/drug treatment, whether inpatient or outpatient. (Chapter 13)

Dependent Living (DPL) was coded "yes" for clients 21 years old and over and not independent. Independence requires either:

- (a) —that the client can handle 12 daily living activities without help
or
- (b) —that the client does not have another person organizing his or her life,
 - does not live with a caregiver,
 - needs help with at most 3 of the 12 critical daily activities, and
 - can pay at least some of his or her own expenses. (Chapter 14)

Problems With Employment (PWE) was coded "yes" if a client is 21 years old and over and not effectively employed. Effectively employed is defined as meeting any one of the following three criteria:

1. Having no other cash support except his/her own wages
2. Weekly earnings of \$280 or more
3. Being currently employed 20 hours or more per week, and
 - Having none or only mild job problems (that is, no repeated problems and a maximum of two "minimal problems;" minimal problem is a score of "1" on LHI questions that follow: trouble getting hired, trouble holding a job, being fired from a job, and losing a job without understanding why),
and
 - Having had three or fewer jobs in the past two years. (Chapter 15)

Secondary Disabilities Overview

Figure 7.2 shows prevalences of the six major Secondary Disabilities by three age groups. Prevalence in adults does not exceed prevalence in adolescents except for Confinement and Alcohol and Drug Abuse.

7.2 History of Secondary Disabilities by three age groups at interview






Figure 7.3 through 7.5 consider only those 253 clients 12 years and over. Figure 7.3 compares the prevalence of Secondary Disabilities for males and females. Males have higher rates of Disrupted School Experience, Trouble With the Law, and Confinement.

7.3 History of Secondary Disabilities among clients ≥ 12 years old by sex



There are mild but systematic differences between clients with FAS and those with FAE in terms of the prevalence of Secondary Disabilities (see Figure 7.4). In all matters except independence and employment, clients with FAE (although often thought to be more "mildly affected") have higher rates of the Secondary Disabilities.

7.4 History of Secondary Disabilities among clients ≥ 12 years old by diagnosis



Low IQ seems to have a similar pattern of "protective" effect as a diagnosis of FAS (with which it is formally tied through the CNS criterion of FAS). For the 38 clients technically classifiable as mentally retarded, there are lower rates of Alcohol and Drug Problems, Disrupted School Experience, Trouble With the Law, and Confinement. All of them are in dependent living situations, and 94% have employment problems (Figure 7.5). An IQ above 85 (average for this full sample) is clearly not "protective" and may actually be associated with additional legal troubles and additional confinements.

7.5 History of Secondary Disabilities among clients ≥ 12 years old by IQ level





Figure 7.5 shows that a diagnosis before age 6 is related to lower prevalence of all secondary disabilities except perhaps Mental Health Problems.

7.6 History of Secondary Disabilities among clients ≥ 12 years old by age at diagnosis




8

Mental Health Problems (MHP)

Mental Health Problems (MHP) is a binary summary of reported mental health concerns, problems, or treatment for clients of any age. It is coded 1 for those clients who had one or more of a long list of possible mental health problems, had ever gone to a psychotherapist or a counselor for a mental health problem, or had ever been a client in a psychiatric or mental hospital. At 94% of the full Secondary Disabilities sample (Figure 8.1), Mental Health Problems is by far the most prevalent Secondary Disability, accounting for twice as many clients as Disrupted School Experience or Trouble with the Law, the next most frequent problems. Because of the overwhelming prevalence of Mental Health Problems, hardly any clients were, in fact, "protected."

8.1 History of Mental Health Problems (MHP) by sex, diagnosis and age at interview (n=415)



For those who were protected, Figure 8.2 shows the strongest protective factors against MHP: (1) DDD services eligibility, (2) below median FARS score, (3) higher than the median number of years per living situation, (4) living in a home of above median quality (during childhood ages 8-12), and (5) living in a stable and nurturant household above median proportion of life (72-100%). Because the base rate of Mental Health Problems in this sample is so close to 100%, these odds ratios are more subject to sampling error than the others in Figure 6.1.

8.2 Risk/Protective Factors for History of Mental Health Problems among clients 12 years and older at interview, max n = 253




Figure 8.3 shows that most clients of any age with Mental Health Problems receive treatment for these problems.

8.3 History of Mental Health Problems: Components of the Secondary Disability by age at interview (n=415)




Figure 8.4 shows the rate of Mental Health Problems by age. The most frequent mental health problems for children and adolescents are attention deficit problems (61%), followed by depression and suicide threats. Over half the adults were reported to have depression problems, 43% have made suicide threats, and 23% have made suicide attempts; 29% of adults have psychotic symptoms.

8.4 History of mental health problems by age at interview (n=415)






Figure 8.5 shows that among clients who were reported to have a mental health problem, the first sign of mental health problems with onset at age five or under, was attention deficit problems (51%), followed by behavior problems (25%) including conduct problems (6%) and assaultive behaviors (7%). By contrast, if the onset of mental health problems was after age 21, the most commonly reported Mental Health Problem was depression (62%), or psychotic behavior (12%).

8.5 First sign of a mental health problem by age of onset among clients with MHP (n=310)




The most frequent reason reported for obtaining outpatient mental health treatment is attention problems, followed by depression. For inpatient treatment, the primary reasons are suicide threats and depression (Figure 8.6). One in three of all hospitalized clients have been hospitalized for over two months, including 3% for over one year (Figure 8.7).

8.6 Mental Health Problems: Prevalence of therapy and hospitalization by reason for treatment



8.7 Length of stay in a mental health hospital (n=67)






Disrupted School Experience (DSE)


Disrupted School Experience (DSE) is a binary summary of reported disruptions of schooling for clients of any age. It is set to 1 for those clients who ever were suspended from school, expelled from school, or dropped out of school. Figure 9.1 shows that 60% of clients 12 years and older have had Disrupted School Experience.

9.1 History of Disrupted School Experience (DSE) by sex, diagnosis and age at interview (n=411)




The important protective factors against DSE (items in Figure 9.2 that lower the odds of DSE) include: an early diagnosis, longer duration of living situations, longer duration in a stable and nurturant home, living in a good quality home, and having a driver's license. In addition, having applied for DDD (Division of Developmental Disabilities) services and found to be eligible was a specific protective factor against DSE.

9.2 Risk/Protective Factors for History of Disrupted School Experience among clients 12 years and older at interview, max n = 250




The rate of DSE is higher after childhood when one in two adolescents have been suspended, one in four expelled, and more than one in four have dropped out of school (Figure 9.3).

9.3 Disrupted School Experience: Components of the Secondary Disability by age at interview (n=411)




The most frequent learning problems (Figure 9.4) are attention problems (70%) and repeatedly incomplete school work (55-60%). Nearly half the adolescents and adults had failed a grade in school. The most frequent behavior problems ever experienced (Figure 9.5) were repeated problems in getting along with peers (60%) and repeatedly being disruptive in class (55-60%). Clients with DSE had about twice as many of these problems as compared to clients without DSE (Figure 9.6).

9.4 Learning problems by age at interview (n=395-407)




9.5 Repeated behavior problems by age at interview (n=403-408)



Figures 9.6 through 9.8 show that DSE is correlated with learning problems and behavior problems. Although rates of repeated poor attention are high whether or not students also experience DSE, rates of failing grades and repeatedly failing classes are much higher among those who also had DSE. Being suspended, expelled from school or dropping out is an unfortunate concomitant for students with problems doing the school work and/or have behaviors that are disruptive or aggressive. School truancy is a frequent behavior problem (48%). Clearly children who are not at school for any reason are unable to benefit from the structure and support that the school offers for the time they are away from school.

9.6 Distributions of number of problems observed repeatedly in school among patients 12 years and older



9.7 Learning problems by presence or absence of DSE (n= 235-244)




Figure 9.8 shows that all repeated behavior problems display higher rates among clients with DSE.

9.8 Repeated behavior problems by presence or absence of DSE (n= 240-245)






Figure 9.9 shows that 40% of the clients with FAS/FAE were known to have been in special education.

9.9 Types of early intervention programs attended (N=369-404)




Where schools put emphasis on remediation is of interest, given that over 80% of clients over age 21 are in dependent living situations. It may be more effective for schools to focus more on practical daily living skills training. We found that 65% of the clients with FAS/FAE had received remedial help in Reading and Arithmetic, but fewer than one in three had received life skills training, and only one in five had received occupational or physical therapy (Figure 9.10).

9.10 Types of remedial interventions received at school (N=393-397)



Compared to Washington State statistics (Figure 9.11), clients with FAS/FAE graduate less frequently from high school than other students (counting all types of graduation including special education, GED, etc.). Even after age 21 years, only 30% have gone to any type of post-secondary school education. The category membership of a person in the LJI sample is determined hierarchically from left to right for Figure 9.11. Thus, a 16 year old who is both in school and has been suspended twice is coded as being among the 94% who are "in school." The 5% of 12-17 year olds graphed as DSE are those who are not still in school.

9.11 Educational status in full study sample and in WA state



Notes:

"in school" means currently attending *and* does not yet have a high school diploma. This includes a few clients who may be attending Community college/GED programs.

"graduated" means the client does have a high school diploma (any kind), but does not have any higher education experience.

"higher education" means the client *has* have a high school diploma and has received (or is receiving) some higher education.

"DSE" refers to clients who are not in school and never graduated, and are DSE positive.

We found no clients in the sample who have a high school diploma (any type) and neither graduated, nor higher education.


10

Trouble With the Law (TWL)

Trouble with the Law (TWL) is a binary summary of reported interactions with police, authorities, or the judicial system for clients of any age. It is set to 1 for those clients who ever were in trouble with the police or authorities, or were ever arrested for, charged with, or convicted of any of the seven types of criminal behavior delineated in Figure 10.3.

TWL is reported for 14% of 6-11 year olds, and higher rates in adolescence and adulthood (61% and 58%, respectively). Figure 10.1 shows the prevalence of TWL by sex, diagnosis, and age at interview.

10.1 History of Trouble With the Law (TWL) by sex, diagnosis and age at interview (n=412)



The most effective factor protecting against TWL is eligibility for DDD (Division of Developmental Disabilities) services (Figure 10.2). Of clients 12 years and over, 62 had applied for DDD (applying, by itself, is neither a risk nor a protective factor): 35 were found eligible and 27 were not. Of the 27 "not eligible," 21 (or 78%) were coded yes for TWL. Of the 35 found eligible for DDD services, 16 (46%) were coded yes for TWL. (No data on the relative timing of these events is available.) Whereas DDD provides services (case managers, job placement and coaching, and shelter), SSI provides only money; eligibility for SSI (Supplemental Security Income) was *not* a protective factor against TWL.

10.2 Risk/Protective Factors for History of Trouble with the Law among clients 12 years and older at interview, max n = 251




Figure 10.3 shows that crimes against persons are the most frequently reported in every age group. This category encompasses theft (including shoplifting), burglary, assault, and murder.

10.3 History of Trouble with the Law: Components of the Secondary Disability by age at interview (n=407-412)





Figure 10.4 shows that shoplifting/theft is most often the first type of crime reported.

10.4 Nature of first trouble with the law among clients ≥ 12 years old with TWL by diagnosis (n=151)




About one in three clients 12 years and older committed a first crime between 9 and 14 years of age. Very few clients had their first TWL after age 20 (Figure 10.5). The rate of TWL was mildly lower in the FAS subgroup, but distributions of first offenses for the two diagnoses were similar (Figure 10.4).

10.5 Age at first trouble with the law by diagnosis among clients ≥ 12 years old at interview




Females were less likely to be charged after the police were called, and less likely to be convicted than were males (Figure 10.6).

10.6 Trouble with the law: arrests, charges, convictions by sex among clients ≥ 12 years old at interview




Figures 10.7 and 10.8 show that among clients who have had trouble with the law, the most prevalent types of sentencing are juvenile justice and juvenile detention, and the most prevalent sentencing alternatives are probation and community service.

10.7 Trouble with the law: type of sentence among clients ≥ 12 years old at interview with TWL (n=151)




10.8 Trouble with the law: type of sentencing alternatives among clients ≥ 12 years old at interview with TWL (n=150)




Figures 10.9 and 10.10 examine TWL in relation to two other Secondary Disabilities: Disrupted School Experience (DSE) and Alcohol and Drug Problems (ADP). Staying in school is related to a lower rate of TWL by a factor of 2.1. Absence of Alcohol and Drug Problems (ADP) is also a factor against TWL, decreasing the rate by almost half.

10.9 History of TWL and its components by Disrupted School Experiences among clients ≥ 12 years old at interview (n=248)



10.10 History of TWL and its components by Alcohol and Drug Problems among clients ≥ 12 years old at interview (n=250)




11

Confinement (CNF)


Confinement (CNF) is a binary score summarizing restrictions of personal freedom—the most serious consequence of any secondary disability. CNF is set to 1 for those clients who have ever been in a mental hospital (23% of the clients over 12 years of age), have ever been in an alcohol/drug inpatient treatment program (15%), or have ever been incarcerated for a crime (35%).

11.1 History of Confinement (CNF) by sex, diagnosis and age at interview (n=412)




The protective factors against Confinement are the same as the "universal" set for the first six Secondary Disabilities as a group (notice how smooth the profile is in Figure 11.2). This score seems to extract the essence of the three most serious specific secondary disabilities that we have already examined. What protects the client against confinement is what protects him or her against the other secondary disabilities, namely a stable and nurturant household, an early diagnosis, not being a victim of violence, relatively few distinct living situations, and the like.

11.2 Risk/Protective Factors for History of Confinement among clients 12 years and older at interview, max n = 251



Confinement was reported by caregivers of clients under 12 years of age at the time of interview only for mental health problems (Figure 11.3), but Figure 11.4 shows that age at first confinement was often younger than 12 years.

11.3 History of Confinement: Components of the Secondary Disability by age at interview (n=410-415)



11.4 Age of first confinement by type of confinement among clients ≥ 12 years old at the time of interview





Figure 11.5 indicates that among the clients who were confined, many were confined for more than one reason. For example, clients who had been in mental health inpatient treatment, had also been in alcohol and drug inpatient treatment (22%) or had also been incarcerated (40%). Furthermore, 13 clients were reported to have experienced confinement of all three types.

11.5 Types of confinements in combination with each other among clients ≥ 12 years old at interview




12

Inappropriate Sexual Behavior (ISB)

Inappropriate Sexual Behavior (ISB) is a binary score summarizing problems with sexual behavior among clients of any age. ISB is coded 1 for clients who were ever sentenced to a sexual offender's treatment program or who were reported to have "repeated" problems with one or more of the 10 inappropriate sexual behaviors identified in Figure 12.3.

Figure 12.1 shows prevalence of ISB by sex, age, and diagnosis. There do not seem to be important differences in rates of ISB by these three client characteristics.

12.1 History of Inappropriate Sexual Behaviors (ISB) by sex, diagnosis and age at interview (n=408)



The strongest risk factor against ISB is violence against client (Figure 12.2). Being a victim of violence increases the odds of sexually inappropriate behavior by fourfold. Inappropriate Sexual Behavior is the only one of the six main Secondary Disabilities (Figure 6.1) for which low IQ is not a protective factor. (As we will see in Chapters 14 and 15, IQ < 70 is also not protective for the two adult problems.) The reason that low IQ does not protect against ISB is that low IQ is a *risk* factor for three components of ISB, namely masturbating in public (odds ratio 6.5), inappropriate advances (odds ratio 2.8), and inappropriate touching (odds ratio 2.6).

12.2 Risk/Protective Factors for History of Inappropriate Sexual Behavior among clients 12 years and older at interview, max n = 247






Figure 12.3 itemizes 10 inappropriate sexual behaviors and their prevalence in this sample. Repeated sexual advances toward others are the most prevalent, at 18% of the sample overall.

12.3 History of Inappropriate Sexual Behaviors reported as repeated behaviors (n=388-408)




Figures 12.4 and 12.5 show that female clients who exhibited inappropriate sexual behaviors, were more likely to have experienced sexual victimization, and to receive therapy to treat their ISB. Whereas male clients who exhibited inappropriate sexual behaviors were more likely to get involved in legal trouble

12.4 Prevalence of sexual victimization, therapy for ISB, and trouble with the law for sexual behavior by six common inappropriate sexual behaviors: Among females



12.5 Prevalence of sexual victimization, therapy for ISB, and trouble with the law for sexual behavior by six common inappropriate sexual behaviors: Among males




13

Alcohol and Drug Problems (ADP)

Alcohol and Drug Problems (ADP) is a binary score summarizing problems with alcohol or drug habits among clients of any age. ADP is coded 1 for clients who have ever had alcohol abuse problems, drug abuse problems, or have ever been in alcohol or drug abuse treatment (either outpatient or inpatient).

Figure 13.1 shows that report of ADP is not different for male versus female clients, but is related to age at interview and to diagnosis.

13.1 History of Alcohol/Drug Problems (ADP) by sex, diagnosis and age at interview



The obvious specific risk factor for ADP is living with alcohol abusers and drug users. Other important risk factors are from the "general" RPF set: late diagnosis of FAS/FAE, and violent behavior against clients (Figure 13.2).

13.2 Risk/Protective Factors for History of Alcohol/Drug Problems among clients 12 years and older at interview, max n = 252





Figure 13.3 shows that among clients who were still younger than 12 years at interview, there was no Alcohol and Drug Problems reported. However, as will be seen in subsequent figures, problems often began at ages younger than 12 years among clients 12 years and older at the time of the interview (Figure 13.4). Overall, ADP-related rates among adolescents and adults were 33% for alcohol problems, 23% for street drug problems, and 23% for treatment. Overall ADP was noted for 35% of the clients 12 years and older.

13.3 Alcohol/Drug Problems: Components of the Secondary Disability by age at interview



The 57 clients with street drug abuse problems are almost entirely "nested" within the 81 clients with alcohol problems. The same is true for use of treatment programs. Among clients with both alcohol and street drug problems, alcohol abuse began on average about 2 years before the abuse of street drugs. Sixty-five percent of clients 12 years and older who had abused alcohol later abused street drugs. Also, among clients engaging in treatment for both alcohol and street drug abuse, the age at first treatment was the same for alcohol as for drug use, as if the multiple addictions were often needed to ensure action.

Possibly quicker treatment for alcohol abuse could lower the rate of later street drug abuse.

13.4 Age of onset of alcohol problems, by sex and diagnosis among clients with alcohol problems






Figure 13.5 shows that 70% of clients who have been in treatment had their first alcohol treatment before age 21

13.5 Age of first alcohol treatment among alcohol abusers, by sex and diagnosis




Figures 13.6 and 13.7 show the choices of treatment used by clients with alcohol abuse problems and drug abuse problems by sex of client. There is not much difference in use of the different treatment types by sex. There also does not seem to be an overall difference in rate of utilization of different treatment types. Inpatient treatment is as highly utilized as other types of treatment.

13.6 Alcohol problems and treatment type among clients \geq 12 years old by sex




13.7 Illicit drug problems and treatment type among clients \geq 12 years old by sex




Figures 13.8 and 13.9 depict the four stages of alcohol and drug use respectively: "no use"/"using, but not abusing" / "abusing, but no treatment yet"/"abusing and treatment received." Female FAS clients have the highest rate of abstinence. Among those who become alcohol abusers, they have the lowest rate of treatment (40% compared to 65% for alcohol abusers as a group). Male FAS clients have the highest rate of use. Among those who become alcohol abusers, they also have the highest rate of treatment.

13.8 Stages of alcohol use among clients ≥ 12 years old by sex and diagnosis




A very similar pattern is observed for street drug use (Figure 13.9). Female FAS clients have the highest rate of street drug avoidance, but among those who become drug abusers, they have the lowest rate of treatment (33% compared to 58% for all street drug abusers as a group). Male FAS clients have the highest rate of street drug use.

13.9 Stages of illicit drug use among clients ≥ 12 years old by sex and diagnosis



Many clients have avoided alcohol use completely. The reasons caretakers report for the clients abstaining from alcohol are as presented in Figure 13.10.

13.10 Reasons for abstinence among clients ≥ 12 years old reported not to consume alcohol at present (n=163)




14

Dependent Living (DPL) Over 21 Years

Dependent Living (DPL) could only reasonably be defined for the 90 people who are 21 years and older. Two sets of criteria were developed for independence; one could qualify as "independent" by either one. The first focuses on independence in 12 daily living activities (regardless of whether or not support people are available); the second focuses more on *not* having a support person organizing their lives, *not* living with a caregiver, and being able to handle at least 9 of the 12 daily living activities, including paying own expenses. No one was counted as an independent adult who does not manage his or her own money (See Figure 14.3 for this list of 12 daily activities). Figure 14.1 shows that 83% of adults live dependently.

14.1 Prevalence of Dependent Living (DPL), by sex and diagnosis



Only one of our "universal" RPF's is protective against DPL diagnosis before age six. Several RPF's specific to DPL are apparent in Figure 14.2, including a VABS ABC score of 65 or above, IQ above 70, a VABS ABC being within 15 points of IQ, a low FABS score (below the median), and being female. The environmental RPF's include having a driver's license and not having (presumably, not needing) two caretakers.

Dependent living and problems with employment each characterized approximately 80% of the adults with FAS/FAE. 70% had both. Only 8% (7 clients) had neither.

14.2 Risk/Protective Factors for Dependent Living among clients 21 years and older at interview, max n = 89




Figure 14.3 lists 12 daily activities that people sometimes or frequently need help with. The daily tasks requiring help by the largest numbers of clients 21 years and older include managing money and making decisions.

14.3 Daily life activities among clients \geq 21 years old (n=90)




Figure 14.4 shows that female FAE clients are most likely to be able to do all 12 daily activities without help, but all 20 are over 21 years old, and only 5 (25%) of them are reported to be capable of doing these tasks.

14.4 Number of daily activities among clients \geq 21 years old, by sex and diagnosis





Figure 14.5 shows that even as adults, people with FAS and FAE experience social problems which significantly compromise their daily functioning. Hence, it is not surprising that over 80% of these adults have difficulty living independently and need intensive help and supervision with managing their daily activities.

14.5 Current caregiver reports of social problems "sometimes" or "frequently" experienced among clients ≥ 21 years old (n=76-88)



15

Problems With Employment (PWE)

Problems With Employment (PWE) were endemic and variable in this group of clients. We ended up defining a group who were *relatively* problem-free. The clients in this group have little in common except that they are all employed. Beyond that, we have combined three more specific groups, overlapping, but not nested, defined as follows:

Group 1. No financial support except own earnings.

Group 2. Earns \$280 a week or more.

Group 3. This category was added based on the characteristics of several specific individuals who seemed to be effectively employed but didn't meet any of the preceding criteria. These were people who were not in a sheltered workshop, were working at least half time, and had no more than 3 jobs in the past 2 years. In addition, from the list of problems under Group 1, they had no repeated problems and only 1 or 2 problems of any kind.

PWE is defined by all clients 21 years and older who are *not* relatively problem-free. Figure 15.1 shows that 79% of clients 21 years and older have PWE.

15.1 History of Problems With Employment (PWE) among clients \geq 21 years old, by sex and diagnosis (n=90)




Figure 15.2 shows that a VABS Adaptive Behavior Composite higher than the median is a protective factor for successful employment. There are also two clear intrinsic risk factors against successful employment: being FAS (as opposed to FAE) and having a low IQ. No person with FAS or FAE who had an IQ 70 or below met our criteria for successful employment, even though low IQ was not explicit in the definition. In addition, one specific protective factor emerged (having a driver's license), and some of the "universal" set of protective factors emerged: more than the median number of years per living situation followed closely by never experiencing violence, an early diagnosis, and living in a nurturant and stable home for over 72% of life.

15.2 Risk/Protective Factors for Problems With Employment among clients 21 years and older at interview, max n = 90




Figure 15.3 shows that holding a job is a bigger problem than getting a job.

15.3 Problems acquiring and maintaining employment by clients ≥ 21 years old (n=78-83)




Figure 15.4 itemizes many problems that clients have on the job. At least one, "poor task comprehension" (57%), may reflect poor job placement. Educating personnel directors and job placement services staff to special characteristics of clients with FAS or FAE may be mutually beneficial both to the clients and employers.

15.4 Type of employment problems experienced while on a job (n=80-83)

